

# CS499 Homework 7 (First Draft)

## Interstellar

### Exercise 7.1

(1) Since  $e$  is in the minimum spanning tree, we split the minimum spanning tree into two components by deleting  $e$ . Let the vertices in the two components consist  $S$  and  $V \setminus S$  respectively. Since there is no circle in a tree, obviously  $e$  is the only edge which is good and cross this cut, which means no edge from  $X$  crosses this cut.

(2) Suppose  $e$  is not the minimum weight edge crossing this cut, assume there is an edge  $e'$  which has less weight and crosses this cut.  $e'$  can replace  $e$  and consists a spanning tree with less weight. This means  $e$  is not in the minimum spanning tree, which means  $e$  is not good, which contradicts the condition.

### Exercises 7.4




Figure 1:

**Exercises 7.5** Obviously, if two vertices are connected in  $T_c$ , they are connected in  $G_c$ , since  $T_c$  is in  $G_c$ .

Suppose  $u, v$  are connected in  $G_c$ , but not connected in  $T_c$ . Let two connected components in  $T_c$  contain  $u$  and  $v$  respectively be  $A$  and  $B$ . Let  $e$  be an edge in  $G_c$  that connect  $A$  and  $B$ . Using definition,  $w(e) \leq c$ . Since  $A$  and  $B$  are not connected in  $T_c$ , there must be an edge  $e'$  in  $T$  that connects  $A$  and  $B$ , and  $w(e') > c$ . So,  $e' \notin e$ . Obviously  $T$  which contains  $e'$  is not the minimum spanning tree, since  $e'$  can be replaced by  $e$  with less weight. This contradicts the condition. So, if two vertices are connected in  $G_c$ , they are connected in  $T_c$ .

### Exercise 7.8

As the picture shows , for  $\forall c, m_c(T) = m_c(T')$ .




Figure 2:

### Exercise 7.9

Sort by the weight of  $T'$ 's edges and  $T$ 's edges , we have  $(a_1, a_2, a_3, \dots, a_{n-1})$  ,  $(b_1, b_2, b_3, \dots, b_{n-1})$  . Suppose  $a_i \neq b_i$  ,  $\forall k < i, a_k = b_k$  and  $w(a_i) \geq w(b_i)$  , there are two situations:

(1) edge  $b_i$  exists in the  $T$  , then we can find  $j(j > i)$  and  $a_j = b_i$  . Because  $w(b_i) = w(a_j) \geq w(a_i) \geq w(b_i)$  ,  $w(a_i) = w(b_i) = w(a_j)$  . So we can exchange  $a_i$  and  $a_j$  and new sequence is still ordered .  $T$ 's and  $T'$ 's position is the same edge.

(2) edge  $b_i$  doesn't exist in the  $T$  , then we add  $b_i$  to  $T$  to form a cycle . Because  $T$  is a minimum spanning tree ,  $w(\text{edge in the cycle}) \leq w(b_i)$  . And we can find  $a_j(j > i)$  and  $a_j$  doesn't exist in the  $T'$  and  $a_j$  in the cycle) . Because  $w(b_i) \geq w(a_j) \geq w(a_i) \geq w(b_i)$  ,  $w(b_i) = w(a_i) = w(a_j)$  . So we can change  $a_j$  with  $b_i$  . Turn to the situation (1).

So we know the ordered edge weight list of any two minimum spanning trees is the same.

Obviously,  $m_c(T) = m_c(T')$ .

### Exercise 7.10

108 Suppose there are two minimum spanning tree , sort by the weight of  $T'$ 's edges and  $T'$ 's edges , we  
 109 have  $(a_1, a_2, a_3, \dots, a_{n-1})$  ,  $(b_1, b_2, b_3, \dots, b_{n-1})$ .  $\exists i, a_i \neq b_i$ , based on the 7.9, the ordered edge  
 110 weight list of any two minimum spanning trees is the same, so  $w(a_i) = w(b_i)$ . But no two edges of  
 111  $G$  have the same weight , so there is contradiction . So  $G$  has exactly one minimum spanning tree!

112 **Question:**

113  
 114 **1.** In Exercise 7.11 & 7.12, how can we compute the number of functions with a core of size  $k$ ?  
 115  $(1 \leq k \leq n)$   
 116  
 117  
 118  
 119  
 120  
 121  
 122  
 123  
 124  
 125  
 126  
 127  
 128  
 129  
 130  
 131  
 132  
 133  
 134  
 135  
 136  
 137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145  
 146  
 147  
 148  
 149  
 150  
 151  
 152  
 153  
 154  
 155  
 156  
 157  
 158  
 159  
 160  
 161