

CS499 Homework 9 (First Draft)

Interstellar

Exercise 9.1

We define $f_1 : N \rightarrow N$

$$f_1(0) = 0, f_1(1) = 1, \dots, f_1(n) = n.$$

We define $f_2 : N \rightarrow N^2$ based on this graph:

	0	1	2	3	4	5	...
0	(0,0)	(0,1)	(0,2)	(0,3)	(0,4)	(0,5)	...
1	(1,0)	(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	...
2	(2,0)	(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	...
3	(3,0)	(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	...
...

Figure 1:

$$f_2(0) = (0,0), f_2(1) = (0,1), f_2(2) = (1,0) \dots$$

We define $f_3 : N \rightarrow N^3$ based on this graph:

	0	1	2	3	4	5	...
$f_2(0)=(0,0)$	(0,0,0)	(0,0,1)	(0,0,2)	(0,0,3)	(0,0,4)	(0,0,5)	...
$f_2(1)=(0,1)$	(0,1,0)	(0,1,1)	(0,1,2)	(0,1,3)	(0,1,4)	(0,1,5)	...
$f_2(2)=(1,0)$	(1,0,0)	(1,0,1)	(1,0,2)	(1,0,3)	(1,0,4)	(1,0,5)	...
$f_2(3)=(2,0)$	(2,0,0)	(2,0,1)	(2,0,2)	(2,0,3)	(2,0,4)	(2,0,5)	...
...

Figure 2:

$f_2(0) = (0, 0, 0), f_2(1) = (0, 0, 1), f_2(2) = (0, 1, 0) \dots$

And so on, we can define $f_k, k \in N$. Now we can define a bijection $N \rightarrow N^*$ base on this graph:

	0	1	2	3	4	5	...
f_1	$f_1(0)$	$f_1(1)$	$f_1(2)$	$f_1(3)$	$f_1(4)$	$f_1(5)$...
f_2	$f_2(0)$	$f_2(1)$	$f_2(2)$	$f_2(3)$	$f_2(4)$	$f_2(5)$...
f_3	$f_3(0)$	$f_3(1)$	$f_3(2)$	$f_3(3)$	$f_3(4)$	$f_3(5)$...
f_4	$f_4(0)$	$f_4(1)$	$f_4(2)$	$f_4(3)$	$f_4(4)$	$f_4(5)$...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots

Figure 3:

We have $0 \rightarrow f_1(0), 1 \rightarrow f_1(1), 2 \rightarrow f_2(0) \dots$. This is a bijection $N \rightarrow N^*$.

Exercise 9.2

We can define a bijection from $\{0, 1\}^N$ to N as follows.

Given $A = (a_1, a_2, \dots, a_n), a_i \in \{0, 1\}$ for $1 \leq i \leq n$, let $f(A) = 2^n + a_1 2^{n-1} + a_2 2^{n-2} + \dots + a_n 2^0 - 1$. For every element A in $\{0, 1\}^N$, there is one $f(A)$ in N . For every element $a \in N$, according to the binary code, there is one $f^{-1}(a) \in \{0, 1\}^N$. So, f is a bijection from $\{0, 1\}^N$ to N . Therefore, $\{0, 1\}^N \cong N$.

Using the given fact that $N \times N \cong N$, it is clear that $\{0, 1\}^N \cong \{0, 1\}^N \times \{0, 1\}^N$.

Using the given fact that $R \cong \{0, 1\}^N$, it is clear that $R \cong R \times R$.

Exercises 9.3

Using the given fact that $R \cong \{0, 1\}^N$ and the fact we proved in 9.2 that $\{0, 1\}^N \cong N$, we only need to prove that $N \cong N^N$. N^N is actually $N^* - \{\in\}$. Using the method of Hilbert's Hotel, it is obvious that $N^N \cong N^*$. We have proved in 9.1 that $N \cong N^*$. Therefore, $R \cong R^N$.

Exercise 9.5

$000 \dots, 100 \dots, 1100 \dots, 11100 \dots$ According to this rule, the first n bits of the n_{th} sequence are 1, and the remaining bits are 0. Obviously, these sequences constitute a countably and infinite chain.

Exercise 9.6

$100 \dots, 0100 \dots, 00100 \dots, 000100 \dots$ According to this rule, the n_{th} bit of the n_{th} sequence is 1, and the remaining bits are 0. Obviously, these sequences constitute a countably and infinite antichain. **Exercise 9.8**

We can form a bijection f from $0, 1^N$ to set A , which is a subset of $0, 1^N$ as follow: Assume a string s is an element of $0, 1^N$ and $t = f(s)$, the first k digit of s_1 (we call it s_k) determines the $(2^k + 1) - th$ to $2^{k+1} - th$ digits of t (we call them $t_{2^k+1} to t_{2^{k+1}}$ as the following rule:

Consider the first k digits as a binary number a_k , then t_{2^k+1} to $t_{2^k+a_k}$ are 1 and the $t_{2^k+a_k+1}$ to $t_{2^{k+1}}$ are 0. Specially, we define that the first 2 digits of t are always '00'.

Here is an example:

$$s = 1011 \dots$$

$$f(s) = 00, 10, 1100, 11111000, 111111111100000, \dots$$

Obviously, f is a bijection and A is uncountable. Also, any two elements t_1, t_2 of A is comparable. Assume two elements s_1, s_2 is different and their first different digit is the k -th digit. The k -th

108 element of s_1 is 1. Then for any m such that $m \geq k$, the binary number of the first m digit of s_1 is
109 greater than that of s_2 , which leads to the conclusion that the according string of t_1 is "greater" than
110 t_2 . Thus A is the chain required.
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161