

# CS499 Homework 9 (First Draft)

## Interstellar

### Exercise 9.1

We define  $f_1 : N \rightarrow N$

$$f_1(0) = 0, f_1(1) = 1, \dots, f_1(n) = n.$$

We define  $f_2 : N \rightarrow N^2$  based on this graph:

	0	1	2	3	4	5	...
0	(0,0)	(0,1)	(0,2)	(0,3)	(0,4)	(0,5)	...
1	(1,0)	(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	...
2	(2,0)	(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	...
3	(3,0)	(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	...
...	...	...	...	...	...	...	...

Figure 1:

$$f_2(0) = (0,0), f_2(1) = (0,1), f_2(2) = (1,0) \dots$$

We define  $f_3 : N \rightarrow N^3$  based on this graph:

	0	1	2	3	4	5	...
$f_2(0)=(0,0)$	(0,0,0)	(0,0,1)	(0,0,2)	(0,0,3)	(0,0,4)	(0,0,5)	...
$f_2(1)=(0,1)$	(0,1,0)	(0,1,1)	(0,1,2)	(0,1,3)	(0,1,4)	(0,1,5)	...
$f_2(2)=(1,0)$	(1,0,0)	(1,0,1)	(1,0,2)	(1,0,3)	(1,0,4)	(1,0,5)	...
$f_2(3)=(2,0)$	(2,0,0)	(2,0,1)	(2,0,2)	(2,0,3)	(2,0,4)	(2,0,5)	...
...	...	...	...	...	...	...	...

Figure 2:

$f_2(0) = (0, 0, 0), f_2(1) = (0, 0, 1), f_2(2) = (0, 1, 0) \dots$

And so on, we can define  $f_k, k \in N$ . Now we can define a bijection  $N \rightarrow N^*$  base on this graph:

	0	1	2	3	4	5	...
$f_1$	$f_1(0)$	$f_1(1)$	$f_1(2)$	$f_1(3)$	$f_1(4)$	$f_1(5)$	...
$f_2$	$f_2(0)$	$f_2(1)$	$f_2(2)$	$f_2(3)$	$f_2(4)$	$f_2(5)$	...
$f_3$	$f_3(0)$	$f_3(1)$	$f_3(2)$	$f_3(3)$	$f_3(4)$	$f_3(5)$	...
$f_4$	$f_4(0)$	$f_4(1)$	$f_4(2)$	$f_4(3)$	$f_4(4)$	$f_4(5)$	...
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$

Figure 3:

We have  $0 \rightarrow f_1(0), 1 \rightarrow f_1(1), 2 \rightarrow f_2(0) \dots$ . This is a bijection  $N \rightarrow N^*$ .

### Exercise 9.2

We can define a bijection from  $\{0, 1\}^N$  to  $\{0, 1\}^N \times \{0, 1\}^N$  as follows.

Given  $A = (a_1 a_2 a_3 a_4 \dots, b_1 b_2 b_3 b_4 \dots)$ , we define  $f(A) = a_1 b_1 a_2 b_2 a_3 b_3 a_4 b_4 \dots$ . To be more precisely,

$$f(A)[i] = \begin{cases} A[1][\frac{i+1}{2}], & i \text{ is odd number} \\ A[2][\frac{i}{2}], & i \text{ is even number} \end{cases}$$

Obviously, for each  $A \in \{0, 1\}^N \times \{0, 1\}^N$ , there is only one  $f(A) \in \{0, 1\}^N$ . For each  $B \in \{0, 1\}^N$ , there is only one  $B = f^{-1}(B) \in \{0, 1\}^N \times \{0, 1\}^N$ . Therefore,  $f$  is a bijection and  $\{0, 1\}^N \cong \{0, 1\}^N \times \{0, 1\}^N$ .

Using the fact that  $R \cong \{0, 1\}^N$ , we can get  $R \cong R \times R$ .

### Exercises 9.3

We use the Cantor's method to prove that. For any  $A \in (\{0, 1\}^N)^N$ , we define that  $f(A)$  is the  $\{0, 1\}^N$  sequence we get by following the blue line as follows.



162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215

Here is an example:

$$s = 1011.....$$

$$f(s) = 00, 10, 1100, 11111000, 1111111111100000, .....$$

Obviously,  $f$  is a bijection and  $A$  is uncountable. Also, any two elements  $t_1, t_2$  of  $A$  is comparable. Assume two elements  $s_1, s_2$  is different and their first different digit is the  $k$ -th digit. The  $k$ -th element of  $s_1$  is 1. Then for any  $m$  such that  $m \geq k$ , the binary number of the first  $m$  digit of  $s_1$  is greater than that of  $s_2$ , which leads to the conclusion that the according string of  $t_1$  is "greater" than  $t_2$ . Thus  $A$  is the chain required.