

APPENDIX A

THE MINIMUM ERROR OF THE RANK PREDICTION WITHOUT PRIOR KNOWLEDGE

Given a set of ranks that uniformly spread across a limited range, what is the minimum prediction error we can obtain without any prior knowledge except for the range? Assume that the range is $[0, K]$ and the set's size is infinite. The mean absolute error (MAE) and mean squared error (MSE) can be derived as follows.

Without knowledge of the ground truth value, one may *randomly* take values from the range $[0, K]$ as the predicted ranks. In this case, the errors are:

$$E_r^{MAE} = \frac{1}{K} \int_0^K \frac{1}{K} \left[\int_0^{\hat{p}} (\hat{p} - p) dp + \int_{\hat{p}}^K (p - \hat{p}) dp \right] d\hat{p} = \frac{K}{3}, \quad (1)$$

$$E_r^{MSE} = \frac{1}{K} \int_0^K \frac{1}{K} \int_0^K (\hat{p} - p)^2 dp d\hat{p} = \frac{K^2}{6}, \quad (2)$$

where p and \hat{p} are the ground truth and predicted rank, respectively.

In addition to random prediction, one may use a fixed value as the predicted rank. In this case, the errors are:

$$E_f^{MAE} = \frac{1}{K} \left[\int_0^{\hat{p}} (\hat{p} - p) dp + \int_{\hat{p}}^K (p - \hat{p}) dp \right] = \frac{\hat{p}^2}{K} - \hat{p} + \frac{K}{2}, \quad (3)$$

$$E_f^{MSE} = \frac{1}{K} \int_0^K (\hat{p} - p)^2 dp = K\hat{p}^2 - K^2\hat{p} + \frac{K^3}{3}. \quad (4)$$

It is easy to derive their minimum with respect to \hat{p} :

$$\min E_f^{MAE} = \frac{K}{4}, \text{ when } \hat{p} = \frac{K}{2}, \quad (5)$$

$$\min E_f^{MSE} = \frac{K^3}{12}, \text{ when } \hat{p} = \frac{K}{2}. \quad (6)$$

We set $K = 100$ in our experiments; therefore, the minimum MAE should be the $\min E_f^{MAE} = 100/4 = 25$, and the minimum MSE should be the $E_r^{MSE} = 100^2/6 = 1666.66$. The former can be used as a baseline to compare the MAE obtained by the APN, i.e., MAE in Tables 1, 2, and 3. The latter is where the 1666.66 in Eq. 15 comes from.