Syllabus Download R and RStudio Readings Courseware **Course Info** Discussion **R Tutorials Contact Us Progress** Community Office Hours

Question 2

A study was conducted to compare the resting pulse rates of college smokers and non-smokers. The data for a randomly selected group is summarized in the table below. Pulse rates were normally distributed within each group.

Group	Sample Size (n)	Average Pulse Rate (bpm)	Standard Deviation of Scores
Smokers	26	80	5
Non-Smokers	32	74	6

(1 point possible)

2a. What is the appropriate method for analyzing this data?

1 of 6 02/27/2015 07:23 PM

Q	Paired T-	UT.7.01x Courseware edX -Test dent T-Test ent T-Test	https://courses.edx.org/courses/UTAustinX/UT.7.01x/3T20	14/courseware/ac18c
Helb	Hide Answer	You have used 0 of 1 submissions		
(1 point possible) 2b. What is the alternative hypothesis for this test if the researchers expect smoking to raise pulse respect to the properties of the researchers expect smoking to raise pulse respect to the properties of the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect smoking to raise pulse respect to the researchers expect to the			s expect smoking to raise pulse rates?	
	Hide Answer	You have used 0 of 1 submissions		
2	(1 point possible 2c. How many do	le) legrees of freedom should we use for this test if we a	are to estimate rather than use a calculator?	02/27/2015 07:23 PM

Help

Answer:	25
----------------	----

Hide Answer

You have used 0 of 1 submissions

(1 point possible)

2d. What is **t-critical**, assuming α =0.05? (Round to 3 decimal places.) Use your answer to 2c. to help.

Answer: 1.708

Hide Answer

You have used 0 of 1 submissions

(1 point possible)

2e. Calculate the **standard error**. (Round to 2 decimal places.)

Hide Answer

You have used 0 of 1 submissions

(1 point possible)

2f. Calculate the **test statistic**. (Round to 2 decimal places, and use rounded values from previous answers.)

Help



Answer: 4.15

Hide Answer

You have used 0 of 1 submissions

(1 point possible)

2g. Is there evidence to suggest that the pulse rate of smokers is higher on average than the pulse rate of non-smokers?



Hide Answer

You have used 0 of 1 submissions

(1 point possible)

4 of 2g. How would the p-value be reported in your conclusion?

p < 0.05



Hide Answer

You have used 0 of 1 submissions



EdX offers interactive online classes and MOOCs from the world's best universities. Online courses from MITx, HarvardX, BerkeleyX, UTx and many other universities. Topics include biology, business, chemistry, computer science, economics, finance, electronics, engineering, food and nutrition, history, humanities, law, literature, math, medicine, music, philosophy, physics, science, statistics and more. EdX is a non-profit online initiative created by founding partners Harvard and MIT.

© 2015 edX Inc.

EdX Open edX, and the edX and Open edX logos are registered trademarks or trademarks of edX Inc.

About edX

About

News

Contact

FAQ

edX Blog

Donate to edX

lobs at edX

Follow Us



Twitter



Facebook



Meetup



in LinkedIn



S+ Google+

02/27/2015 07:23 PM

https://courses.edx.org/courses/UTAustinX/UT.7.01x/3T2014/courseware/ac18c...

Help

6 of 6 02/27/2015 07:23 PM