

## Ignorance priors and transformation groups

← Back to Chapters

### Comments on 12.4

I find the discussion in 12.4 somewhat off the mark. “Statistical decision theory and bayesian analysis” by James O. Berger is a better reference. The summary below is largely based on it, particularly sections 3.3.2 and 6.6.

The word “invariance” presupposes a group action. The most natural setting is that in which a group  $G$  acts on the space  $X$  in which we get data.

---

Example

1:  
(location-

scale

in

1D)

Take

$X =$

$\mathbb{A}^1$

the

affine

line,

and

$G$  the

group

of

(orientation-

preserving)

affine

trans-

for-

ma-

tions

of  $X$ .

As is

com-

mon

(for

ex-

am-

ple in

com-

puter

graph-

ics),

we

repre-

sent

ele-

ments

of  $X$

as

vec-

tors

$\vec{x} =$

$\begin{pmatrix} x \\ 1 \end{pmatrix}$

(the

line

$y = 1$

in-

side

$\mathbb{R}^2$ )

and

then

---

is in-  
deed  
affine-  
linear.  
The  
prod-  
uct  
in  $G$   
is  
then:

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a' & b' \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} aa' & ab' + b \\ 0 & 1 \end{pmatrix}$$

This  
is a  
non-  
commutative  
group.  
We  
have  
the  
fol-  
low-  
ing  
pair  
of  
group  
ho-  
mo-  
mor-  
phisms  
(known  
as a  
short  
exact  
sequence):

$$\mathbb{R} \hookrightarrow G \twoheadrightarrow \mathbb{R}_+$$

---

Here  
 $\mathbb{R}$  is  
the  
real  
num-  
bers  
un-  
der  
addi-  
tion,  
and  
 $\hookrightarrow$   
sends  
a  
num-  
ber  
to a  
trans-  
la-  
tion  
by  
that  
amount;  
on  
the  
other  
hand  
 $\mathbb{R}_+$  is  
posi-  
tive  
reals  
un-  
der  
mul-  
tipli-  
ca-  
tion,  
and  
 $\rightarrow$   
sends  
an  
affine  
map  
to its  
stretch-  
ing  
factor.

---

These  
 maps  
 do  
 not  
 de-  
 pend  
 on  
 any  
 par-  
 ticu-  
 lar  
 way  
 of  
 repre-  
 sent-  
 ing  
 $X$   
 and  
 $G$ .  
 On  
 the  
 other  
 hand,  
 there  
 are  
 “back-  
 ward  
 maps”  
 $\mathbb{R}_+ \rightarrow$   
 $G$   
 and  
 $G \rightarrow$   
 $\mathbb{R}$ ,  
 but  
 those  
 are  
 de-  
 pend  
 on  
 addi-  
 tional  
 choices,  
 like  
 the  
 choice  
 to  
 repre-  
 sent  
 $X$   
 and  
 $G$  in  
 terms  
 of  
 vec-  
 tors

---

Example  
 2:  
 (Location-  
 scale  
 in  
 higher  
 di-  
 men-  
 sions)  
 When  
 $X =$   
 $\mathbb{A}^n$   
 the  
 location-  
 scale  
 group  
 $G$  is  
 the  
 sub-  
 group  
 of  
 those  
 affine  
 trans-  
 forms  
 of  $X$   
 whose  
 lin-  
 ear  
 part  
 is a  
 pure  
 rescal-  
 ing  
 by a  
 (posi-  
 tive)  
 fac-  
 tor.  
 We  
 then  
 have  
 $\mathbb{R}^n \hookrightarrow$   
 $G \rightarrow$   
 $\mathbb{R}_+$ .

---

---

We are interested in distribution of the data, i.e. in probability distributions over  $X$ . Thus we consider a collection of  $\mathcal{P}$  of distributions over  $X$ .

Defintion: The collection  $\mathcal{P}$  is said to be invariant under the action of  $G$  if for any  $p \in \mathcal{P}$  and any  $g \in G$  the pushforward distribution  $g_*p$  is also in  $\mathcal{P}$ .

Since  $(g \cdot h)_*p = g_*(h_*p)$ , this means that  $G$  acts on  $\mathcal{P}$  as well. If  $\mathcal{P}$  is a parametric family, parametrized by space  $\Theta$  then we conclude that  $G$  acts on  $\Theta$ .

---

Example 1 continued:

- a) Let  $\mathcal{P}$  be the collection of all normal distributions on  $\mathbb{A}^1$ . Note that in order to talk about the collection  $\mathcal{P}$  specifying the origin is not necessary. This collection is a invariant under the action of  $G$ . If we pick an origin, then we can use mean  $\mu$  and standard deviation  $\sigma$  as parameters for  $\mathcal{P}$ . We can also use representation of  $G$  by matrices that we have discussed above. Then  $\Theta = \mathbb{R} \times \mathbb{R}_+$  is the parameter space and  $G$  acts on by sending  $(\mu, \sigma)$  to  $(a\mu + b, a\sigma)$ .

(Together with  $x$  being sent to  $ax + b$  this appears as 12.30 in Jaynes.)

- b) Let  $\mathcal{P}$  be the collection of all normal distributions on  $\mathbb{A}^1$ ; after choice of the origin this is the family with pdfs  $\frac{1}{\pi\gamma \left[1 + \left(\frac{x-x_0}{\gamma}\right)^2\right]}$ . We no longer have mean or standard deviation available as a parameters, but we do have location  $x_0$  and scale  $\gamma$ . They transform under  $G$  by the same formulas as  $(\mu, \sigma)$  did before.
- c) Let  $\mathcal{P}$  be the collection of all mixures of normal distributions on  $\mathbb{A}^1$ . After picking the origin, this is the collection of distributions which can be written as  $p = \sum_i^m w_i \mathcal{N}(\mu_i, \sigma_i^2)$  for some  $m \in \mathbb{N}$  and  $w_i > 0$  with  $\sum w_i = 1$ . This collection is invariant under  $G$ . When  $m$  is fixed the subfamily  $\mathcal{P}_m$  it is parametric, with parameters being  $3m$  dimensional vectors  $(\mu_i, \sigma_i, w_i)$ . If  $m$  is not fixed, however, the collection  $\mathcal{P}$  is not parametric in the usual sense of the word.

Example 3: Consider the family  $\mathcal{P}$  of all Gamma distributions on  $X = \mathbb{R}_+$ , with pdfs  $\Gamma_{(k,\theta)}(x) = \frac{1}{\Gamma(k)\theta} \left(\frac{x}{\theta}\right)^{k-1} \exp(-\frac{x}{\theta})$ . Here  $G = \mathbb{R}_+$  acts on  $X$  by multiplication,  $\mathcal{P}$  is invariant, and the action of  $a \in G$  sends  $(k, \theta)$  to  $(k, a\theta)$ .

---