

Pattern Recognition practical 2

Maikel Withagen (s1867733)

Steven Bosch (s1861948)

September 23, 2015

1 Covariance matrix

1.1

Using the code given in the appendix we get the following mean vector:

$$\begin{bmatrix} 5.8000 \\ 5.0000 \\ 6.2000 \end{bmatrix} \quad (1)$$

And the following covariance matrix is yielded:

$$\begin{bmatrix} 3.2000 & 0.2500 & -0.4500 \\ 0.2500 & 2.5000 & -3.7500 \\ -0.4500 & -3.7500 & 5.7000 \end{bmatrix} \quad (2)$$

1.2

We computed the following probability densities:

```
1 >> mvnpdf([5;5;6], mean1, cov1)
2 ans =    0.0543
3 >> mvnpdf([3;5;7], mean1, cov1)
4 ans =    6.1287e-04
5 >> mvnpdf([4;6.5;1], mean1, cov1)
6 ans =    7.0300e-29
```

2 Covariance matrix, analytically

Every element in the covariance matrix is determined by:

$$\sigma_{ij} = \frac{1}{n-1} \sum_{n=1}^N (x_{in} - \mu_i)(x_{jn} - \mu_j) \quad (3)$$

Since $n = 2$ for all the covariance matrices calculated below, we leave that factor out (since $\frac{1}{2-1} = 1$).

2.1

Element 1,1:

$$\sigma_{1,1} = (a - \frac{a+c}{2})^2 + (c - \frac{a+c}{2})^2 \quad (4)$$

$$= (\frac{a-c}{2})^2 + (\frac{c-a}{2})^2 \quad (5)$$

$$= \frac{1}{4}(a-c)^2 + \frac{1}{4}(c-a)^2 \quad (6)$$

$$= \frac{1}{2}(a-c)^2 \quad (7)$$

Similarly for $\sigma_{2,2}$ we get $\frac{1}{2}(b-d)^2$. For the other two elements, $cov(1,2)$ and $cov(2,1)$ which are the same, we get the following:

$$\sigma(1,2) = ((a - \frac{a+c}{2})(b - \frac{b+d}{2}) + (c - \frac{a+c}{2})(d - \frac{b+d}{2})) \quad (8)$$

$$= (\frac{a-c}{2})(\frac{b-d}{2}) + (\frac{c-a}{2})(\frac{d-b}{2}) \quad (9)$$

$$= \frac{(a-c)(b-d) + (c-a)(d-b)}{4} \quad (10)$$

$$= \frac{2ab - 2ad - 2bc + 2cd}{4} \quad (11)$$

$$= \frac{1}{2}(a-c)(b-d) \quad (12)$$

This results in the following covariance matrix:

$$\begin{bmatrix} \frac{1}{2}(a-c)^2 & \frac{1}{2}(a-c)(b-d) \\ \frac{1}{2}(a-c)(b-d) & \frac{1}{2}(b-d)^2 \end{bmatrix} \quad (13)$$

2.2

Element 1,1:

$$\sigma_{1,1} = (a + k - \frac{a+c+2k}{2})^2 + (c + k - \frac{a+c+2k}{2})^2 \quad (14)$$

$$= (\frac{a-c}{2})^2 + (\frac{c-a}{2})^2 \quad (15)$$

$$= \frac{1}{4}(a-c)^2 + \frac{1}{4}(c-a)^2 \quad (16)$$

$$= \frac{1}{2}(a-c)^2 \quad (17)$$

As we can see, the ks are all removed in the simplification, due to writing sums like $a+k$ as the fraction $\frac{2a+2k}{2}$ and subtracting fractions with $2k$ in the numerator, resulting in the ks being removed. Again similarly for $\sigma_{2,2}$ we get $\frac{1}{2}(b-d)^2$. For the other two elements, $cov(1,2)$ and $cov(2,1)$ we get the following (the ks are removed in the simplification process again):

$$\sigma(1,2) = ((a+k - \frac{a+c+2k}{2})(b+k - \frac{b+d+2k}{2}) + (c+k - \frac{a+c+2k}{2})(d+k - \frac{b+d+2k}{2})) \quad (18)$$

$$= (\frac{a-c}{2})(\frac{b-d}{2}) + (\frac{c-a}{2})(\frac{d-b}{2}) \quad (19)$$

$$= \frac{(a-c)(b-d) + (c-a)(d-b)}{4} \quad (20)$$

$$= \frac{2ab - 2ad - 2bc + 2cd}{4} \quad (21)$$

$$= \frac{1}{2}(a-c)(b-d) \quad (22)$$

Thus in the end we get the same matrix as in the previous section:

$$\begin{bmatrix} \frac{1}{2}(a-c)^2 & \frac{1}{2}(a-c)(b-d) \\ \frac{1}{2}(a-c)(b-d) & \frac{1}{2}(b-d)^2 \end{bmatrix} \quad (23)$$

2.3

Element 1,1:

$$\sigma_{1,1} = (ak - \frac{ak+ck}{2})^2 + (ck - \frac{ak+ck}{2})^2 \quad (24)$$

$$= (\frac{ak-ck}{2})^2 + (\frac{ck-ak}{2})^2 \quad (25)$$

$$= \frac{1}{4}(ak-ck)^2 + \frac{1}{4}(ck-ak)^2 \quad (26)$$

$$= \frac{1}{2}(ak-ck)^2 \quad (27)$$

$$= \frac{1}{2}k^2(a-c)^2 \quad (28)$$

Similarly for $\sigma_{2,2}$ we get $\frac{1}{2}k^2(b-d)^2$.

For the other two elements, $cov(1,2)$ and $cov(2,1)$ we get the following:

$$\sigma(1,2) = ((ak - \frac{ak+ck}{2})(bk - \frac{bk+dk}{2}) + (ck - \frac{ak+ck}{2})(dk - \frac{bk+dk}{2})) \quad (29)$$

$$= (\frac{ak-ck}{2})(\frac{bk-dk}{2}) + (\frac{ck-ak}{2})(\frac{dk-bk}{2}) \quad (30)$$

$$= \frac{k^2(a-c)(b-d) + k^2(c-a)(d-b)}{4} \quad (31)$$

$$= \frac{2k^2(ab - ad - bc + cd)}{4} \quad (32)$$

$$= \frac{1}{2}k^2(a-c)(b-d) \quad (33)$$

$$\begin{bmatrix} \frac{1}{2}k^2(a-c)^2 & \frac{1}{2}k^2(a-c)(b-d) \\ \frac{1}{2}k^2(a-c)(b-d) & \frac{1}{2}k^2(b-d)^2 \end{bmatrix} \quad (34)$$

3 2D Gaussian pdf, Mahalanobis distance

3.1

Using the code given in the appendix, we generated the following plot:




Figure 1: Two-dimensional Gaussian pdf plotted on a mesh.

3.2 2D Gaussian pdf, Mahalanobis distance

We determined the Mahalanobis distances using the following definition:

$$r^2 = (x - \mu)\Sigma^{-1}(x - \mu)' \quad (35)$$

This yielded the following distances for the given points:

```

1 >> ([10 10]-mean)*inv(cov)*([10 10]-mean). '
2 ans =      67
3 >> ([0 0]-mean)*inv(cov)*([0 0]-mean). '
4 ans =      17
5 >> ([3 4]-mean)*inv(cov)*([3 4]-mean). '
6 ans =       0
7 >> ([6 8]-mean)*inv(cov)*([6 8]-mean). '
8 ans =      17

```

3.3 Independent identically distributed random binary variables

The code in the appendix yields the following plot:

4 2D Gaussian pdf, Mahalanobis distance

4.1

Using the code given in the appendix, we generated the following plot:




Figure 2: Two-dimensional Gaussian pdf plotted on a mesh.

This plot resembles the binomial distribution with a sequence of 100 experiments and a probability of 0.5. Each person that tosses a coin 100 times can be viewed as a random draw from the binomial distribution with a probability P of 0.5 because of the 50/50 chances of the coin flip. The 1000000 people together form 1000000 draws from this binomial distribution, and by plotting them, an approximation of the distribution is displayed.

By tossing a coin 100 times (N) with a 50% chance to go forward (P), the theoretical mean would be $N \cdot P$ or $100 \cdot 0.5 = 50$. The variance would be $N \cdot P(1-P) = 100 \cdot 0.5(1-0.5) = 25$.

4.2 Multivariate normal density, discriminant functions, minimum error rate classification, unequal priors, dichotomizer

4.3 Naive Bayesian rule

Following the Naive Bayes rules as shown on the slides (so calculating the ratio):

a “We offer our dear customers a wide selection of classy watches”

$$\frac{P(\text{spam} | \text{Customers}, \text{Watches})}{P(\text{non-spam} | \text{Customers}, \text{Watches})} = \frac{p(\text{customers} | \text{spam})}{p(\text{customers} | \text{non-spam})} \frac{p(\text{watches} | \text{spam})}{p(\text{watches} | \text{non-spam})} \frac{p(\text{spam})}{p(\text{non-spam})} \quad (36)$$

$$= \frac{0.005 * 0.0003 * 0.9}{0.0001 * 0.000004 * 0.1} \quad (37)$$

$$= 33750 \quad (38)$$

So with a ratio of 33750:1 (spam:non-spam) it is very likely that the message can be classified as spam (a probability of approximately 0.99997037037).

b “Did you have fun on vacation? I sure did!”

$$\frac{P(\text{spam}|\text{Fun}, \text{Vacation})}{P(\text{non-spam}|\text{Fun}, \text{Vacation})} = \frac{p(\text{fun}|\text{spam})}{p(\text{fun}|\text{non-spam})} \frac{p(\text{vacation}|\text{spam})}{p(\text{vacation}|\text{non-spam})} \frac{p(\text{spam})}{p(\text{non-spam})} \quad (39)$$

$$= \frac{0.00015 * 0.00025 * 0.9}{0.0007 * 0.00014 * 0.1} \quad (40)$$

$$= 3.4439 \quad (41)$$

So with a ratio of 3.4439:1 (spam:non-spam) it is likely that the message can be classified as spam (with a probability of approximately 0.7096315224).

Appendix

Code for assignment 1

```
1 v1 = [4,5,6];
2 v2 = [6,3,9];
3 v3 = [8,7,3];
4 v4 = [7,4,8];
5 v5 = [4,6,5];
6
7 m1 = [v1;v2;v3;v4;v5];
8
9 mean1 = [mean(m1(:,1)); mean(m1(:,2)); mean(m1(:,3))];
10
11 cov1 = cov(m1);
12
13 mean1
14 cov1
```

Code for assignment 3

```
1 % Generate a two-dimensional Gaussian pdf with a mean [3 4] and covariance matrix
2 % 1 0
3 % 0 2
4 mean = [3 4]
5 cov = [1 0; 0 2]
6 dist = gmdistribution(mean,cov)
7
8 inputmatrix = [0 0];
9
10 counter = 1;
11 for i = -10:10
12     for j = -10:10
13         inputmatrix(counter,:) = [i j];
14
15         counter = counter + 1;
16     end
17 end
```

```
18 % 1. Plot this function on [-10 10] x [-10 10] Using the mesh function.
19 output = reshape(pdf(dist, inputmatrix),[21,21])
20 mesh(-10:10,-10:10,output) %
```

Code for assignment 4

```
1 output = zeros(100,1);
2 for i=1:1000000
3     step = 0;
4     for j = 1:100
5         if rand > 0.5
6             % Move forward
7             step = step + 1;
8         end
9     end
10    output(step) = output(step) + 1;
11 end
12
13 plot(output)
```