

# Geometric Modeling

*Notes based on a course given by Boris Thibert for M1 Applied Mathematics,  
Université Grenoble Alpes.*

Malik Hacini

Last updated: **October 30, 2025**

## Contents

Chapter 1: Interpolation .....	2
Lagrange interpolation .....	2
Runge Phenomenon .....	2
Interpolation with splines .....	3
Minimization result .....	3
Chapter 2: Differential geometry of curves .....	4
Representation of curves .....	4
Generalities on parametrized curves .....	4
Reminder : .....	4
Metric properties of curves .....	4
Length of a rectifiable curve .....	4
Arc-length .....	5
Notion of regular curve .....	5
Plane curves .....	6
Curvature .....	7
Osculating circle and center of curvature .....	8
Total curvature .....	8
Space curves .....	8
Principal normal vector and curvature .....	8
Chapter 3: Differential geometry of surfaces .....	11
Generalities .....	12
Tangent spaces .....	12
Metric properties .....	13
First fundamental form .....	13
Length of a curve on a surface .....	14
Curvature .....	15
Differentiable map between surfaces .....	15
Gauss map and second fundamental form .....	15

## Chapter 1: Interpolation

### Lagrange interpolation

#### Theorem 1.1

Let  $a = t_0 < \dots < t_n = b, y_0, \dots, y_n \in \mathbb{R}$

There exists a unique polynomial  $L_n$  of degree  $\leq n$  such that:

$$\forall i \in \{0, n\}, L_n(t_i) = y_i \quad [1]$$

These polynomials are called Lagrange interpolation polynomials. A simple expression is :

$$L_n(t) = \sum_{i=0}^n y_i P_i(t), \quad [2]$$

$$\text{where } P_i(t) = \prod_{j \neq i} \frac{t - t_j}{t_i - t_j}$$

**Proof:** We denote  $E = \mathbb{R}_n[X]$  the space of polynomials of degree  $\leq n$ .

- $E$  is a  $\mathbb{R}$ -vector space
- Let

$$\begin{aligned} \varphi : E &\longrightarrow \mathbb{R}^{n+1} \\ P &\longmapsto (P(t_0), \dots, P(t_n)) \end{aligned} \quad [3]$$

$\varphi$  is a linear map between vector spaces of some dimension. To prove the result, we need  $\varphi$  to be one-to-one map.

Let  $P \in \ker \varphi$

Then :

- $P$  is of degree  $\leq n$
- $P$  vanishes at  $n+1$  points

then by the fundamental theorem of algebra,  $P \equiv 0$

### Runge Phenomenon

We observe in practice the Runge Phenomenon. Let

$$\begin{aligned} f : [-1, 1] &\rightarrow \mathbb{R} \\ x &\mapsto \frac{1}{1 + 25x^2} \end{aligned} \quad [4]$$

$t_i$  uniformly distributed in  $[-1, 1]$

We observe  $\|L_n - f\|_{\infty} \xrightarrow{n \rightarrow +\infty} +\infty$

We have an approximation result :

#### Theorem 1.2

:

Let  $f : [a, b] \rightarrow \mathbb{R} \in \mathcal{C}^{n+1}$

$L_n$  the Lagrange polynomial associated to  $a = t_0 < \dots < t_n = b$

Then  $\|f - L_n\| \leq \frac{1}{(n+1)!} \|q_{n+1}\|_{\infty} \|f^{(n+1)}\|_{\infty}$

when  $q_{n+1}(t) = \prod_{i=0}^n (t - t_i)$

Runge phenomenon :  $\|f^{(n+1)}\|_{\infty}$  goes to infinity quickly

**Proof:**

We introduce the error  $g = f - L_n$

Let  $t \in [0, b] \setminus \{t_i\}$

$$\begin{aligned} k(u) &:= g(u) - q_{n+1}(u) \frac{g(t)}{q_{n+1}(t)} \\ k(t_i) &= 0 \text{ for } i = 0, \dots, n \\ k(t) &= 0 \end{aligned} \quad [5]$$

$k(t)$  vanishes at  $(n+2)$  points

so  $k'(t)$  vanishes at  $(n+1)$  points (by Rolle)

so  $k^{n+1}$  vanishes at 1 point  $\xi$

$$\Rightarrow 0 = k^{n+1}(\xi) = g^{n+1}(\xi) - q_{n+1}^{n+1}(\xi) \frac{g(t)}{q_{n+1}(t)}$$

$$g^{n+1} = f^{(n+1)}(\xi) \text{ and } q_{n+1}^{n+1} = (n+1)!$$

$$\Rightarrow |g(t)| = \frac{1}{(n+1)!} q_{n+1}(t) f^{(n+1)}(\xi) \leq \| \dots \|_{\infty}$$

**Interpolation with splines**

## Theorem 1.3

Let  $y_0, \dots, y_n \in \mathbb{R}$

$a = t_0, \dots, t_n = b$  and  $\alpha, \beta \in \mathbb{R}$

Then there exists a unique function  $S$

such that

- $S|_{[t_i, t_{i+1}]}$  is a polynomial of degree  $\leq 3$
- $S$  is  $\mathcal{C}^2$
- $S(t_i) = y_i$
- $S'(a) = \alpha$  and  $S'(b) = \beta$

order of continuity is one less than the order of polynomial (see order 0 case)

This function  $S$  is called cubic splines

**Minimization result**

## Theorem 1.4

Let  $y_0, \dots, y_n \in \mathbb{R} \dots$

Then the associated spline  $S$  satisfies :

$$S = \arg \min \int_a^b f''(t)^2 dt \quad [6]$$

So splines are the solution to the minimization problem of reducing the energy ( $L^2$  norm) of the second derivative.

**Skecth of proof :** Let  $g \in E$  and  $e = f - S$  the error with  $S$  spline

Step 1 :

Show

$$\forall h : [a, b] \rightarrow \mathcal{C}^0, \text{ linear on } [t_i, t_{i+1}],$$

$$\int_a^b e''(x)h(x)dx = 0 \quad [7]$$

Step 2:

$$\begin{aligned} \int_a^b f''^2 &= \int (e + S)^2 \\ &= \int e''^2 + 2 \int e''S'' + \int S''^2 \\ &= \int e''^2 + \int S''^2 \end{aligned} \quad [8]$$

$$\Rightarrow \forall f \in E \int f''^2 \geq \int S''^2$$

We need to show uniqueness of the minimizer

If  $\int f''^2 = \int S''^2$  then  $\int e''^2 = 0$

## Chapter 2: Differential geometry of curves

### Representation of curves

- Drawings
- Parametrized curves :

$$\gamma : [a, b] \rightarrow \mathbb{R}^d \quad [9]$$

- Implicit representation:

$$\mathcal{C} = \{(x, y), x^2 + y^2 = R^2\} = f^{-1}(0) \text{ where } f : t \rightarrow x^2 + y^2 - R^2 \quad [10]$$

### Generalities on parametrized curves

#### Reminder :

Let  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}^2$  of class  $\mathcal{C}^n$ , and  $t_0 \in ]a, b[$ . Performing a Taylor expansion around  $t_0$  yields :

$$f(t) = f(t_0) + (t - t_0)f'(t_0) + \frac{(t - t_0)^2}{2!}f''(t_0) + \dots + \frac{(t - t_0)^n}{n!}f^{(n)}(t_0) + o((t - t_0)^n) \quad [11]$$

Geometrically, we have  $\overrightarrow{f(t_0)f(t)} = f(t) - f(t_0) = ((t - t_0))f'(t_0)$ . Thus, when it does not vanish,  $f'(t_0)$  is tangent to  $\mathcal{C} = f([a, b])$  at  $f(t_0)$ .

Now, let  $p$  be the smallest  $k \geq 1$  such that  $f^{(k)}(t_0) \neq 0$  and  $q$  the smallest  $k > p$  such that  $(f^{(p)}(t_0), f^{(q)}(t_0))$  are linearly independant.

Then, in the reference frame centered around  $f(t_0)$  and with basis vectors  $(f^{(p)}(t_0), f^{(q)}(t_0))$ , we have

$$f(t) \approx \begin{pmatrix} \lambda(t) \\ \mu(t) \end{pmatrix} \text{ where } \lambda(t) = \frac{(t - t_0)^p}{p!} \text{ and } \mu(t) = \frac{(t - t_0)^q}{q!}. \quad [12]$$

TODO : Add figure

**Remark :** In practice,  $\lambda(t) \gg \mu(t)$ , thus the influence of  $f^{(q)}$  is vastly exaggerated in the figure.

### Metric properties of curves

#### Length of a rectifiable curve

In this section  $\|\cdot\|$  is implicitly assumed to be the euclidean norm  $\|\cdot\|_2$ .

Let's first define the length of a curve in the most geometrical way possible.

## Length of a rectifiable curve

Definition 2.1

Let  $\gamma : [a, b] \rightarrow \mathbb{R}^d$  be  $\mathcal{C}^k$  with  $k \geq 1$ , and  $\mathcal{S}$  be the set of uniform subdivisions of  $[a, b]$ . For  $s = (t_0, t_1, \dots, t_n) \in \mathcal{S}$ , we define

$$l(s) = \sum_{i=0}^{n-1} \|\gamma(t_{i+1}) - \gamma(t_i)\|. \quad [13]$$

If  $\{l(s), s \in \mathcal{S}\}$  is bounded, we say  $\gamma$  is **rectifiable** and define the length  $L(\gamma)$  of  $\gamma$  as

$$L(\gamma) = \sup_{s \in \mathcal{S}} l(s) \quad [14]$$

This leads to the natural integral characterisation that follows.

## Integral expression of the length of a rectifiable curve

Theorem 2.1

Let  $\gamma : [a, b] \rightarrow \mathbb{R}^d$  be a rectifiable curve. We have :

$$L(\gamma) = \int_a^b \|\gamma'(t)\| dt \quad [15]$$

## Arc-length

## Arc-length of a rectifiable curve

Definition 2.2

Let

## Notion of regular curve

## Regular Curve

Definition 2.3

A curve  $\gamma : I \subset \mathbb{R} \rightarrow \mathbb{R}^d$  of class  $\mathcal{C}^k$  is said to be regular at  $t$  if  $\gamma'(t) \neq 0$ .

## Remarks:

- A  $\mathcal{C}^1$  regular curve admits a tangent at every point
- We sometimes call the set  $\tilde{C} = \gamma(I) \subset \mathbb{R}^d$  the “geometric curve” or the support of the curve.
- For  $\mathcal{C}^1$  regular curves,  $I$  is connected/compact  $\Rightarrow \tilde{C}$  is connected/compact

**A non regular smooth curve**

Example 2.1

Consider the following  $\mathcal{C}^0$  curve :

$$t \rightarrow \begin{cases} \left(t, t^{\frac{3}{2}}\right) & \text{if } t \geq 0 \\ \left(|t| - |t|^{\frac{3}{2}}\right) & \text{else} \end{cases} \quad [16]$$




Figure 1: The geometric curve

This geometric curve can be represented by the map  $g : t \rightarrow (t^2, t^3)$  which is  $\mathcal{C}^\infty$ . The curve is not regular in 0 because  $g'(0) = 0$ .

**Reparametrization**

Proposition 2.1

Let  $\gamma : I \subset \mathbb{R} \rightarrow \mathbb{R}^d$  and  $\varphi : I \rightarrow J$  a  $\mathcal{C}^k$  diffeomorphism (i.e  $\varphi$  bijective,  $\varphi$  of class  $\mathcal{C}^k$ ,  $\varphi'(t) \neq 0$ , thus  $\varphi^{-1}$  is of class  $\mathcal{C}^k$ ).

Then  $\gamma \circ \varphi^{-1} : J \rightarrow \mathbb{R}^d$  is called an admissible reparametrization of  $\gamma$  and  $\varphi$  or  $\varphi^{-1}$  is called an admissible change of variable.

**Remark:**  $\varphi(t) = t^3$  is not an admissible change of variable, because  $\varphi'(0) = 0$ , even tho  $\varphi$  is bijective and  $\mathcal{C}^k$ .

We sometimes also see the following alternative definition of geometric curve, which states that a geometric curve does not depend on the parametrization.

**Geometric Curve**

Definition 2.4

A “geometric curve” is an equivalence class for the relation

$$\gamma : I \rightarrow \mathbb{R}^d \sim g : J \rightarrow \mathbb{R}^d \text{ if } \exists \varphi : J \rightarrow I \text{ a } \mathcal{C}^k\text{-diffeomorphism s.t } \gamma = g \circ \varphi^{-1}. \quad [17]$$

Proposition 2.2

If  $\gamma : I \rightarrow \mathbb{R}^d$  is  $\mathcal{C}^k$  regular and  $\varphi : J \rightarrow I$  is an admissible change of variable, then  $\gamma \circ \varphi$  is  $\mathcal{C}^k$  regular.

**Proof:**  $(\gamma \circ \varphi)'(t) = \gamma'(\varphi(t)) \cdot \varphi'(t) \neq 0$

**Remark :** Non-admissible reparametrization exist and can change the regularity of the curve, but not the admissible ones.

**Plane curves**

We will now study the curvature of plane and space curves separately, starting with plane curves.

**Serret-Fresnet frame**

Definition 2.5

Let  $\gamma : I \rightarrow \mathbb{R}^2$  a  $\mathcal{C}^1$  regular curve. The frame  $(f(s), T(s), N(s))$  is called the Serret-Fresnet frame at  $t$ .

**Remark :** When the curve is parametrized with arc-length,  $T(s) = \gamma'(s)$ , thus  $N(s) = \text{Rot}_{\frac{\pi}{2}}(\gamma'(s))$ .

### Curvature

#### Curvature for plane curves

Definition 2.6

Let  $\gamma : I \rightarrow \mathbb{R}^2$  a  $\mathcal{C}^2$  regular curve parametrized by arc length. The curvature of  $\gamma$  at  $\gamma(s)$  is defined by

$$\kappa_s(s) = \langle \gamma''(s), N(s) \rangle = \pm \|\gamma''(s)\| \quad [18]$$

**Remark :**

- We define curvature to be positive when  $\gamma''(s) = N(s)$  and negative when  $\gamma''(s) = -N(s)$ . This can be interpreted by convexity  $\Leftrightarrow$  positive curvature for graphs of  $\mathbb{R} \rightarrow \mathbb{R}$  functions.
- For any non arc-length parametrization,  $\gamma''(t)$  is not (in general) orthogonal to  $\gamma'(t)$  and the above formula doesn't hold. It still holds if  $\|\gamma'(t)\|$  is constant as  $T(t) = \frac{\gamma'(t)}{\|\gamma'(t)\|}$ . We can however derive a general formula for any parametrization.

#### Planar curvature for general parametrizations

Proposition 2.3

Let  $\gamma : I \rightarrow \mathbb{R}^2$  a  $\mathcal{C}^2$  regular curve. We have

$$\kappa_t(t) = \frac{\det(\gamma'(t), \gamma''(t))}{\|\gamma'(t)\|^3} \quad [19]$$

**Proof:** Let  $\tilde{\gamma} = \gamma \circ \varphi^{-1}$  be the arc-length reparametrization of  $\gamma$ . We denote  $t = \varphi^{-1}(s)$ . Then,

$$\tilde{\gamma}'(s) = (\gamma \circ \varphi^{-1})'(s) = \gamma'(\varphi^{-1}(s)) \cdot (\varphi^{-1})'(s) = \frac{\gamma'(t)}{\|\gamma'(t)\|}. \quad [20]$$

Hence

$$\begin{aligned} \tilde{\gamma}''(s) &= \frac{\gamma''(t)}{\|\gamma'(t)\|^2} + \gamma'(t) \frac{d}{dt} \left( \frac{1}{\|\gamma'(t)\|} \right) \\ \Rightarrow \det(\tilde{\gamma}'(s), \tilde{\gamma}''(s)) &= \frac{\det(\gamma'(t), \gamma''(t))}{\|\gamma'(s)\|^3}. \end{aligned} \quad [21]$$

Indeed,  $\gamma'(t) \frac{d}{dt} \left( \frac{1}{\|\gamma'(t)\|} \right)$  is colinear to  $\gamma'(t)$  and thus vanishes in the det. Furthermore, since  $\kappa(s) = \langle \tilde{\gamma}''(s), N(s) \rangle$  and  $\tilde{\gamma}''(s)$  is colinear to  $N(s)$ , we have  $\det(\tilde{\gamma}'(s), \tilde{\gamma}''(s)) = \det(T(s), \kappa_s(s)N(s))$ . Thus, since  $(T(s), N(s))$  is a direct basis,  $\det(T(s), N(s)) = 1$ , hence  $\det(T(s), \kappa_s(s)N(s)) = \kappa_s(s)$ . Finally,

$$\kappa_s(s) = \kappa_t(t) = \det(\tilde{\gamma}'(s), \tilde{\gamma}''(s)) = \frac{\det(\gamma'(t), \gamma''(t))}{\|\gamma'(s)\|^3}. \quad [22]$$

#### Serret-Fresnet formula

Proposition 2.4

Let  $\gamma : I \rightarrow \mathbb{R}^2$  a  $\mathcal{C}^2$  regular curve parametrized by arc length. We have

$$\begin{cases} T'(s) = \kappa(s)N(s) \\ N'(s) = -\kappa(s)T(s) \end{cases} \quad [23]$$

**Proof:**

- TODO
- We have  $\langle N(s), T(s) \rangle = 0$ . Differentating this equation yields

$$\langle N'(s), T(s) \rangle + \underbrace{\langle N(s), T'(s) \rangle}_{\kappa(s)} = 0. \quad [24]$$

Moreover,

$$\langle N(s), N(s) \rangle = 1 \Rightarrow \langle N'(s), N(s) \rangle = 0. \quad [25]$$

Thus, in the direct basis  $(T(s), N(s))$ , we have  $N'(s) = \begin{pmatrix} -\kappa(s) \\ 0 \end{pmatrix}$ , i.e  $N'(s) = -\kappa(s)T(s)$ .

### Osculating circle and center of curvature

#### Center and radius of curvature

Definition 2.7

Let  $\gamma : I \rightarrow \mathbb{R}^2$  a  $\mathcal{C}^2$  regular curve. For all  $t$  where  $\kappa(t) \neq 0$ , we define the center of curvature  $c(t)$  as

$$c(t) = f(t) + \frac{1}{\kappa(t)}N(t). \quad [26]$$

The quantity  $R(t) = \frac{1}{\|\kappa(t)\|}$  is called the radius of curvature.

#### Remark :

- The circle of center  $c(t)$  and radius  $R(t)$  is called the osculating circle to  $\gamma$  at  $\gamma(t)$ .
- The osculating circle is placed towards the (local) interior of the curve : this justifies the sign convention in the curvature definition.

#### Osculating circle approximation

Proposition 2.5

The osculating circle to  $\gamma$  at  $\gamma(t)$  is tangent to  $\mathcal{C}(\gamma)$  at  $\gamma(t)$  and approaches  $\gamma$  at order 2 around it.

**Proof:** Admitted.

### Total curvature

#### Total curvature

Proposition 2.6

Let  $\gamma : [a, b] \rightarrow \mathbb{R}^2$  a  $\mathcal{C}^2$ -regular curve parametrized by arc-length. We have

$$\int_a^b k(s)ds = \theta(b, a) (\text{:= total angle variation of } \gamma) \quad [27]$$

This quantity is called total curvature.

#### Proof:

We denote  $f(s) = \begin{pmatrix} x(s) \\ y(s) \end{pmatrix}$  and  $\theta(s) = (O_x, T(s))$  the angle between the tangent vector to  $f$  and the  $x$ -axis. We define  $\theta$  to be differentiable thus continuous, thus defined up to  $2\pi k$ . Then,

$$T(s) = \begin{pmatrix} \cos \theta(s) \\ \sin \theta(s) \end{pmatrix} \quad \text{and} \quad N(s) = \begin{pmatrix} -\sin \theta(s) \\ \cos \theta(s) \end{pmatrix}. \quad [28]$$

Hence,

$$T'(s) = \theta'(s) \begin{pmatrix} -\sin \theta(s) \\ \cos \theta(s) \end{pmatrix} \quad \text{and} \quad T'(s) = k(s)N(s). \quad [29]$$

This yields:

$$\theta'(s) = k(s) \quad [30]$$

And the conclusion trivially follows.

### Space curves

In this section, we let  $\gamma : I \subset \mathbb{R} \rightarrow \mathbb{R}^3$  be a  $\mathcal{C}^k$  regular space curve ( $k$  will be specified).

#### Principal normal vector and curvature

#### Space curvature

Definition 2.8

If  $\gamma$ ,  $\mathcal{C}^2$  regular, is parametrized by arc-length, the curvature  $\kappa$  is defined by:

$$\kappa(s) = \|\gamma''(s)\|. \quad [31]$$

**Remark :** Space curvature is unsigned, due to the fact there is no unique normal vector to the curve, but a plane.



## Space curvature for general parametrizations

Proposition 2.7

For any regular parametrization of  $\gamma$ , we have

$$\kappa(t) = \frac{\|\gamma'(t) \wedge \gamma''(t)\|}{\|\gamma'(t)\|^3}. \quad [32]$$

**Proof:** We denote  $\tilde{\gamma} = \gamma \circ \varphi^{-1}$  the arc-length parametrization of  $\gamma$ , with  $t = \varphi^{-1}(s)$ . We have

$$\begin{aligned} \langle \tilde{\gamma}'(s), \tilde{\gamma}'(s) \rangle &= 1 \implies 2\langle \tilde{\gamma}''(s), \tilde{\gamma}'(s) \rangle = 0 \\ \implies \tilde{\gamma}''(s) &\perp \tilde{\gamma}'(s) \quad \text{and} \quad \|\tilde{\gamma}'(s)\| = 1 \\ \implies \kappa(s) &= \|\tilde{\gamma}''(s)\| = \|\tilde{\gamma}'''(s) \wedge \tilde{\gamma}'(s)\|. \end{aligned} \quad [33]$$

We already derived in 2D (same proof works) that

$$\tilde{\gamma}''(s) = \frac{\gamma''(t)}{\|\gamma''(t)\|} + \gamma'(t) \cdot \frac{d}{ds} \left( \frac{1}{\|\gamma'(t)\|} \right). \quad [34]$$

Hence,

$$\kappa_s(s) = \left\| \frac{\gamma''(t)}{\|\gamma''(t)\|^2} + \gamma'(t) \cdot \lambda(t) \wedge \frac{\gamma'(t)}{\|\gamma'(t)\|} \right\| = \frac{\|\gamma''(t) \wedge \gamma'(t)\|}{\|\gamma'(t)\|^3} := k_t(t). \quad [35]$$

## Principal normal vector

Definition 2.9

If  $\gamma$  is  $\mathcal{C}^2$  regular and parametrized by arc-length, when  $\kappa(s) \neq 0$ , i.e  $\gamma''(s) \neq 0$ , we define the principal normal as

$$N(s) = \frac{\gamma''(s)}{\|\gamma''(s)\|} = \left( \frac{1}{\kappa(s)} \frac{d}{ds} T(s) \right) \quad [36]$$

Where:

- $T(s) = \gamma'(s)$  is a tangent vector
- $\gamma$  is said to be biregular if  $\forall s, \gamma''(s) \neq 0$ .

**Remark :**

- The term **principal** comes from the fact that when you perform a Taylor expansion of  $\gamma$  around  $s$ , the curve lies (at order 2) in the  $(T(s), N(s))$  plane, since  $N(s)$  is defined through  $\gamma''(s) \perp T(s) = \gamma'(s)$ .
- We can also express the principal normal using a non-arc length parametrization.
- The principal normal and the planar normal vector are the same up to the sign. Thus, the principal normal is not continuous in general. Indeed, it can change orientation instantly on inflexion points.

## Osculating plane/sphere, radius and center of curvature, evolute

Definition 2.10

- The osculating plane at point  $\gamma(t)$  is the plane spanned by  $T(t)$  and  $N(t)$ .
- The radius of curvature is  $R(t) = \frac{1}{\kappa(t)}$ .
- The center of curvature is  $c(t) = \gamma(t) + R(t)N(t)$ .
- The evolute is the set of centers of curvatures.
- The osculating sphere at point  $\gamma(t)$  is  $\mathcal{S}(C(t), R(t))$ .
- The osculating circle is the osculating sphere's cut through the osculating plane.

## Binormal vector and Serret-Fresnet frame

Definition 2.11

If  $\gamma$  is  $\mathcal{C}^2$  regular and arc-length parametrized and  $s$  is a biregular point of  $\gamma$ , then we define

- $B(s) = T(s) \wedge N(s)$  the binormal vector at  $\gamma(s)$ .
- $\gamma(s), T(s), N(s), B(s)$  the Serret-Fresnet frame.

Another interesting property of  $\gamma$  is encoded by the variation of the osculating plane, that we call torsion. This can be defined using the derivative of the binormal vector, which is orthogonal to the osculating plane. This requires  $\gamma$  to be  $\mathcal{C}^3$  regular.

## Proposition 2.8

If  $\gamma$  is  $\mathcal{C}^3$  regular and arc-length parametrized, then  $B'(s)$  is colinear to  $N(s)$ .

**Proof:**

**First method :**  $\|B(s)\|^2 = \langle B(s), B(s) \rangle = 1 \implies B'(s) \perp B(s)$ . Hence, since  $B(s) \perp N(s)$ ,  $B'(s)$  is colinear to  $N(s)$ .

**Second method :** Differentiating  $B(s)$  using the product rule yields:

$$\begin{aligned} B(s) &= T(s) \wedge N(s) \implies B'(s) = \underbrace{T'(s) \wedge N(s)}_0 + T(s) \wedge N'(s) = T(s) \wedge N'(s) \\ &\implies B'(s) \perp T(s) \\ &\implies B'(s) \text{ is colinear to } N(s). \end{aligned} \quad [37]$$

## Torsion

## Definition 2.12

For  $\gamma$  a  $\mathcal{C}^3$  regular space curve parametrized by arc-length, we define torsion of  $\gamma$  in the point  $\gamma(s)$  for all biregular points  $s$  as

$$\tau(s) = -\langle B'(s), N(s) \rangle = \pm \|B'(s)\| \quad [38]$$

**Remark :**

- Torsion measures the speed of rotation of the binormal vector at the given point. It's sign informs on the direction of rotation : the negative sign in the definition is a matter of convention. We thus define torsion to be negative when in the direction of the normal vector, and negative in the opposite case.
- As  $N(s)$  is colinear to  $B'(s)$ , this is equivalent to  $B'(s) = -\tau(s)N(s)$ .
- As  $N(s) \perp B(s)$ , this is also equivalent to  $\langle N'(s), B(s) \rangle$ .

## Torsion formula for arc-length parametrized curves

## Proposition 2.9

For  $\gamma$  a  $\mathcal{C}^3$  regular space curve parametrized by arc-length, we have

$$\tau(s) = \frac{\det(\gamma'(s), \gamma''(s), \gamma'''(s))}{\|\gamma''(s)\|^2}. \quad [39]$$

**Proof:**

$$\begin{aligned} \tau(s) &= -\langle N(s), B'(s) \rangle \\ &= \text{TODO} \end{aligned} \quad [40]$$

Thus  $\tau(s) = \det(T(s), N(s), N'(s))$ . However,

$$\begin{aligned} \cdot T(s) &= \gamma'(s) \\ \cdot N(s) &= R(s)\gamma''(s) \\ \cdot N'(s) &= R'(s)\gamma''(s) + R(s)\gamma'''(s) \end{aligned} \quad [41]$$

Hence,

$$\begin{aligned} \tau(s) &= \det(\gamma'(s), R(s)\gamma''(s), R'(s)\gamma''(s) + R(s)\gamma'''(s)) \\ &= R(s)^2 \det(\gamma'(s), \gamma''(s), \gamma'''(s)). \end{aligned} \quad [42]$$

The conclusion trivially follows.

**Torsion formula for general biregular parametrization**

Proposition 2.10

For  $\gamma$  a  $\mathcal{C}^3$  regular space curve parametrized by  $t$ , we have

$$\tau(t) = \frac{\det(\gamma'(t), \gamma''(t), \gamma'''(t))}{\|\gamma'(t) \wedge \gamma''(t)\|^2} \quad [43]$$

**Proof:** Admitted.**Geometric interpretation of vanishing torsion**

Proposition 2.11

Let  $\gamma$  be a  $\mathcal{C}^3$  biregular space curve. We have :

$$\gamma \text{ is planar} \Leftrightarrow \tau \equiv 0. \quad [44]$$

**Proof:**

$$\bullet \quad \gamma \text{ is planar} \implies \gamma', \gamma'' \text{ and } \gamma''' \text{ are coplanar} \implies \tau \equiv 0. \quad [45]$$

• Suppose  $\tau \equiv 0$ , i.e  $B(s) \equiv B_0$  (parametrizing  $\gamma$  by arc-length). Then  $\forall s$ ,

$$(\langle \gamma(s), B_0 \rangle)' = \langle \gamma'(s), B_0 \rangle = \langle T(s), B_0 \rangle = 0. \quad [46]$$

This tells us that  $\langle \gamma(s), B_0 \rangle$  is constant, thus  $\gamma(s)$  lies in a plane orthogonal to  $B_0$ .

**Serret-Fresnet formula**

Proposition 2.12

Let  $\gamma$  be a arc-length  $\mathcal{C}^3$  parametrized space curve. We have

- $T'(s) = \kappa(s)N(s)$
- $B'(s) = -\tau(s)N(s)$
- $N'(s) = -\kappa(s)T(s) + \tau(s)B(s)$ .

**Proof:** First two points were already proven earlier.

•  $(\|N(s)\|) = 1 \Rightarrow N' \perp N$ . Since  $(T(s), B(s))$  is thus a direct basis, we have  $N'(s) = \lambda(s)T(s) + \mu(s)B(s)$ . We have

$$\lambda(s) = \langle N'(s), T(s) \rangle = -\langle N(s), T'(s) \rangle = -\langle N(s), \kappa(s)N(s) \rangle = -\kappa(s). \quad [47]$$

And

$$\mu(s) = \langle N'(s), B(s) \rangle = -\langle N(s), B'(s) \rangle = -\langle N(s), -\tau(s)N(s) \rangle = \tau(s). \quad [48]$$

**Fundamental theorem of local theory of curves**

Theorem 2.2

Let  $\kappa : [a, b] \mapsto \mathbb{R}$  and  $\tau : [a, b] \mapsto \mathbb{R}$  be  $\mathcal{C}^1$  functions, and such that  $\kappa$  is always strictly positive.

Then, there exists a unique (up to a rigid motion)  $\mathcal{C}^3$  arc-length parametrized curve  $\gamma : [a, b] \mapsto \mathbb{R}^3$  whose curvature is  $\kappa$  and torsion  $\tau$ .

**Proof:** Admitted. Relies on Cauchy-Lipschitz.**Chapter 3: Differential geometry of surfaces**

Representations of surfaces include :

- Parametrization :  $f : U \subset \mathbb{R}^2 \mapsto \mathbb{R}^3$ 
  - Special case : the graph of a map  $\varphi : \mathbb{R}^2 \mapsto \mathbb{R}$  is  $f(x, y, z) = (x, y, \varphi(x, y))$ .
- Implicit : An inverse image of a map  $g : \mathbb{R}^3 \mapsto \mathbb{R}$ .
  - Example : Let  $g(x, y, z) = x^2 + y^2 + z^2 - R^2$ . Then  $g^{-1}(\{0\})$  is the sphere of center  $0_{\mathbb{R}^3}$  and radius  $R$ .
- Drawings !

These representations are locally equivalent when surfaces are “regular” (existence of tangent spaces).

In the following section,  $U$  and  $V$  are taken to be open sets of  $\mathbb{R}^2$ .


## Generalities

### Parametrized surface

Definition 3.1

A  $\mathcal{C}^k$  parametrized surface is a  $\mathcal{C}^k$  map  $f : U \subset \mathbb{R}^2 \mapsto \mathbb{R}^3$ .

We are interested in the geometric surface  $S = f(U)$ . However, one such geometric surface can admit many different parametrizations. They are all the same up to a diffeomorphism. More precisely, with  $\varphi$  a  $\mathcal{C}^k$  diffeomorphism, we have:



### Admissible change of variable

Definition 3.2

We say  $\varphi : U \mapsto V$  is an admissible change of variable for a  $\mathcal{C}^k$  parametrized surface if  $\varphi$  is a  $\mathcal{C}^k$  diffeomorphism. The induced reparametrization is  $g = f \circ \varphi^{-1}$ . Indeed,

$$g(V) = f(\varphi^{-1}(V)) = f(U). \quad [49]$$

We are interested in **geometric** properties of curves, thus that do not depend on reparametrization by an admissible change of variable.

## Tangent spaces

### Curve on surface

Definition 3.3

Let  $f : U \mapsto \mathbb{R}^3$  be a  $\mathcal{C}^1$  surface and  $\gamma : I \subset \mathbb{R} \mapsto U$  a  $\mathcal{C}^1$  curve. Then,  $f \circ \gamma$  is a curve on  $S = f(U)$ .

**Examples:** Let  $I_1, I_2 \subset \mathbb{R}$  such that  $I_1 \times I_2 \subset U$ . For any point  $(u_0, v_0) \in I_1 \times I_2$ , we define the **coordinate curves** on  $S$  as  $\gamma_{u_0} : I_1 \mapsto S$  and  $\gamma_{v_0} : I_2 \mapsto S$ :

$$\gamma_{u_0} : v \mapsto \gamma(u_0, v) \quad \text{and} \quad \gamma_{v_0} : u \mapsto \gamma(u, v_0). \quad [50]$$

Remark that  $\gamma'_{u_0}(v_0) = \frac{\partial f}{\partial v}(u_0, v_0)$  and  $\gamma'_{v_0}(u_0) = \frac{\partial f}{\partial u}(u_0, v_0)$ .

### Tangent space

Definition 3.4

The tangent space to  $S = f(U)$  at  $m_0 = f(u_0, v_0)$  is the affine space passing through  $m_0$  and spanned by  $\left(\frac{\partial f}{\partial u}(u_0, v_0), \frac{\partial f}{\partial v}(u_0, v_0)\right)$ . We denote it  $T_{m_0}S$ .

**Remark :** We will often abusively refer to  $T_{m_0}S$  as a vector space, omitting the translation from 0 to  $m_0$ .

Proposition 3.1

$T_{m_0}S$  is (as a vector space) the set of derivatives of parametrized curves on  $S$  at 0 where  $C(0) = m_0$ .

**Proof:**

Let  $C = f \circ \gamma$  be a curve on  $S$ . Then  $C'(t) = D_f(\gamma(t)) \cdot \gamma'(t)$ . We set  $\gamma(0) = (u_0, v_0)$  and  $\gamma'(0) = (\lambda, \mu)$ . Then,

$$C'(0) = D_f(u_0, v_0) \cdot (\lambda, \mu) = \frac{\partial f}{\partial u}(u_0, v_0)\lambda + \frac{\partial f}{\partial v}(u_0, v_0)\mu \in T_{m_0}S. \quad [51]$$

For the reciprocal inclusion, let  $X \in T_{m_0}S$ . We know

$$\exists(\lambda, \mu) \in \mathbb{R}^2, X = \frac{\partial f}{\partial u}(u_0, v_0)\lambda + \frac{\partial f}{\partial v}(u_0, v_0)\mu. \quad [52]$$

We thus pick a curve  $\gamma : I \subset \mathbb{R} \mapsto U$  such that  $\gamma(0) = (u_0, v_0)$  and  $\gamma'(0) = (\lambda, \mu)$ . We denote  $C = f \circ \gamma$ .

Then  $X = D_f(u_0, v_0) \cdot (\lambda, \mu) = C'(0)$ .

**Remarks :**


- We can express the tangent space (as a vector space) as  $T_{m_0}S = \text{Im}(D_f(u_0, v_0))$ .
- A parametrized surface  $f$  is regular at  $m_0$  if  $\text{Rank}(D_f(u_0, v_0)) = 2$ , i.e the tangent space is a **plane**.
- A parametrized surface  $f$  is said to be regular if it is regular at every point.
- A geometric surface  $S$  is said to be regular if there exists a regular parametrization of  $S$ .

**Invariance of regularity of surfaces by reparametrization**

Proposition 3.2

Let  $f : U \mapsto \mathbb{R}^3$  be a  $\mathcal{C}^1$  regular parametrized surface and  $g : V \mapsto \mathbb{R}^3$  a reparametrization of  $f$ .  
Then,  $g$  is regular and the tangent spaces coincide.

**Proof:**



We have  $f = g \circ \varphi$ . Thus,

$$D_f(u_0, v_0) = D_g(\varphi(u_0, v_0)) \circ D_\varphi(u_0, v_0) \quad \text{where} \quad D_\varphi(u_0, v_0) : \mathbb{R}^2 \mapsto \mathbb{R}^2 \text{ is an isomorphism.} \quad [53]$$

Hence,

$$\text{Im}(D_f(u_0, v_0)) = \text{Im}(D_g(u_0, v_0)). \quad [54]$$

It trivially follows that  $g$  is regular if and only if  $f$  is.

## Metric properties

### First fundamental form

**First fundamental form**

Definition 3.5

Let  $f : U \subset \mathbb{R}^2 \mapsto \mathbb{R}^3$  be a  $\mathcal{C}^1$ -regular surface. The first fundamental form is given by :

$$\begin{aligned} i_{m_0} : T_{m_0}S \times T_{m_0}S &\mapsto \mathbb{R} \\ (x, y) &\longrightarrow \langle x, y \rangle \end{aligned} \quad [55]$$

**Remark:** - Since  $f$  is regular, the first fundamental form is canonically identified to a 2D dot product on  $T_{m_0}S$  via change of basis. This justifies focusing on the expression of the first fundamental form in this basis.

- The first fundamental form is a dot product, i.e a positive definite symmetric bilinear form. Our notations will thus be based on the matrix expression of bilinear form

**Matrix of the first fundamental form in the tangent space basis**

Proposition 3.3

We can express the first fundamental form in the basis  $\left(\frac{\partial f}{\partial u}(u_0, v_0), \frac{\partial f}{\partial v}(u_0, v_0)\right) = (U, V)$  of  $T_{m_0}S$  by the following matrix:

$$I_{m_0} = \begin{pmatrix} E_{m_0} & F_{m_0} \\ F_{m_0} & G_{m_0} \end{pmatrix} \quad [56]$$

Where  $E_{m_0} = \langle U, U \rangle$ ,  $F_{m_0} = \langle U, V \rangle$  and  $G_{m_0} = \langle V, V \rangle$ .

**Proof:** Let  $X = \begin{pmatrix} x_u \\ x_v \end{pmatrix}$  and  $Y = \begin{pmatrix} y_u \\ y_v \end{pmatrix}$  be two tangent vectors expressed in the given basis. We have

$$\begin{aligned} i_{m_0}(X, Y) &= \langle x_u U + x_v V, y_u U + y_v V \rangle \\ &= x_u y_u \langle U, U \rangle + (x_u y_v + x_v y_u) \langle U, V \rangle + x_v y_v \langle V, V \rangle \\ &= X^T I_{m_0} Y. \end{aligned} \quad [57]$$

**Remark :**

- From a differential geometry point of view,  $I_{m_0}$  as a matrix amounts to a pull back of the dot product on the parameter space.
- We usually like to represent bilinear forms in orthonormal bases, beware that it is not the case in general for the tangent space basis.

**Local area multiplicative factor expressed by the fundamental form**

Proposition 3.4

We have

$$\left\| \frac{\partial f}{\partial u}(u_0, v_0) \wedge \frac{\partial f}{\partial v}(u_0, v_0) \right\| = \sqrt{\det(I_{m_0})} = \sqrt{E_{m_0} G_{m_0} - F_{m_0}^2}. \quad [58]$$

**Proof:** Let  $u, v \in \mathbb{R}^3$ . We have

$$\|u \wedge v\|^2 = \|u\|^2 \|v\|^2 \sin^2(u, v) \quad \text{and} \quad |\langle u, v \rangle|^2 = \|u\|^2 \|v\|^2 \cos^2(u, v). \quad [59]$$

Thus,  $\|u \wedge v\|^2 = \|u\|^2 \|v\|^2 - \langle u, v \rangle^2$ .

Conclusion trivially follows using  $(u, v) = \left(\frac{\partial f}{\partial u}(u_0, v_0), \frac{\partial f}{\partial v}(u_0, v_0)\right)$ .

**Remark :** Since  $I_{m_0}$  is a positive bilinear form,  $\det(I_{m_0}) > 0$ .

**Length of a curve on a surface****Length of a curve on a surface**

Proposition 3.5

Let  $\gamma : [a, b] \mapsto U$  defined by  $\gamma(t) = \begin{pmatrix} u(t) \\ v(t) \end{pmatrix}$  and  $f : U \mapsto \mathbb{R}^3$ . Denote  $C_\gamma = f \circ \gamma$  and  $l(C_\gamma)$  the length of  $C$ . We have

$$l(C_\gamma) = \int_a^b \sqrt{I_{(f \circ \gamma)(t)}(\gamma'(t), \gamma''(t))} dt \quad [60]$$

**Proof:** We have

$$l(C_\gamma) = l(f \circ \gamma) = \int_a^b \|(f \circ \gamma)'(t)\| dt. \quad [61]$$

Also,

$$\begin{aligned}
\|(f \circ \gamma)'(t)\|^2 &= \|D_f(\gamma(t)) \circ D_\gamma(t)\|^2 \\
&= \left\| \frac{\partial f}{\partial u}(u(t), v(t)) \cdot u'(t) + \frac{\partial f}{\partial v}(u(t), v(t)) \cdot v'(t) \right\|^2 \\
&= u'(t)^2 \underbrace{\left\| \frac{\partial f}{\partial u}(u(t), v(t)) \right\|^2}_E + 2 \cdot u'(t)v'(t) \underbrace{\left\langle \frac{\partial f}{\partial u}(u(t), v(t)), \frac{\partial f}{\partial v}(u(t), v(t)) \right\rangle}_F + v'(t)^2 \underbrace{\left\| \frac{\partial f}{\partial v}(u(t), v(t)) \right\|^2}_G \\
&= I_{\gamma(t)}(\gamma'(t), \gamma''(t)).
\end{aligned} \tag{62}$$

This yields the wanted formula.

#### Area of a surface

#### Proposition 3.6

Let  $f : U \mapsto \mathbb{R}^3$  be a  $\mathcal{C}^1$  regular surface. The area of the surface  $S$  is the following (independent of parametrization) quantity :

$$A(S) = \int_U \sqrt{\det(I_f(u, v))} du dv. \tag{63}$$

**Proof:** Admitted, although it is just an application of [Proposition 3.4](#) to the integral definition of the area of a surface. See [here](#) for details..

## Curvature

### Differentiable map between surfaces

We want to give a meaning to the differential of a map  $\varphi$  between surfaces. More precisely, consider the following diagram :

$$\begin{array}{ccc}
S_1 \subset \mathbb{R}^3 & \xrightarrow{\varphi} & S_2 \subset \mathbb{R}^3 \\
\uparrow f & & \uparrow g \\
U \subset \mathbb{R}^2 & & V \subset \mathbb{R}^2
\end{array}$$

Our goal is to build a linear map that locally represents  $\varphi$ .

### Differentiable map between surfaces

#### Definition 3.6

We say that  $\varphi : S_1 \mapsto S_2$  is differentiable if  $f \circ \varphi \circ \varphi^{-1}$  is differentiable.

It's differential  $D_\varphi(m_0) : T_{m_0}S_1 \mapsto T_{m_1}S_2$  is then defined for every  $X \in T_{m_0}S_1$  as

$$D_\varphi(m_0) := D(\varphi \circ f)(u_0, v_0)(h, k) \in T_{m_0}S_2 \text{ where } (h, k) \in \mathbb{R}^2 \text{ is a vector s.t } X = D_f(u_0, v_0) \cdot (h, k) \tag{64}$$

### Gauss map and second fundamental form

Let  $f : U \mapsto \mathbb{R}^3$  be a  $\mathcal{C}^2$  regular surface and  $m_0 = f(u_0, v_0) \in S$ . The vector

$$N(u_0, v_0) = \frac{\frac{\partial f}{\partial u}(u_0, v_0) \wedge \frac{\partial f}{\partial v}(u_0, v_0)}{\left\| \frac{\partial f}{\partial u}(u_0, v_0) \wedge \frac{\partial f}{\partial v}(u_0, v_0) \right\|} \tag{65}$$

is a unit vector orthogonal to  $T_{m_0}S$ .

#### Proposition 3.7

The line  $\text{Span}(N_{f(u_0, v_0)})$  does not depend on the parametrization of  $S = f(U)$ .

**Proof:** Let  $g : V \mapsto \mathbb{R}^3$  be a reparametrization of  $S$ ,  $f = g \circ \varphi$ . A few lines of algebra yield:

$$N_f(u_0, v_0) = N_g(\varphi(u_0, v_0)) \cdot \frac{\det(D_\varphi(u_0, v_0))}{|\det(D_\varphi(u_0, v_0))|}. \quad [66]$$

Thus  $N_{f(u_0, v_0)}$  is defined up to the sign, hence the span is invariant. Note that this sign is constant since  $\varphi$  is a diffeomorphism.

#### Orientable surface

Definition 3.7

We say that a surface  $S$  is orientable if there exists a continuous unit normal vector on  $S$ .

**Remark:**

- Orientability is a global notion.
- An orientable surface thus has a defined inside and outside based on this normal vector.
- The Moebius strip is non orientable.

#### Gauss map

Definition 3.8

Let  $S$  be a  $\mathcal{C}^2$ -regular orientable surface of  $\mathbb{R}^3$  and  $\mathbb{S}^2$  the unit sphere of  $\mathbb{R}^3$ . We define the continuous map :

$$\begin{aligned} K : S &\longrightarrow \mathbb{S}^2 \\ m_0 &\longrightarrow K(m_0) \end{aligned} \quad [67]$$

Where  $K(m_0)$  is the unit normal vector to  $f$  in  $m_0$ .  $K$  is called the Gauss map.

Proposition 3.8

The map  $D_K(m_0) : T_{m_0}S \longrightarrow T_{K(m_0)}\mathbb{S}^2$  is an endomorphism.

**Remark :** As done previously, we identify the tangent spaces to vector spaces here.

**Proof:**