3.2 The Derivative as a Function


DEFINITION The **derivative** of the function $f(x)$ with respect to the variable x is the function f' whose value at x is

$$f'(x) = \lim_{h \to 0} \frac{f(x + h) - f(x)}{h},$$

provided the limit exists.

Alternative Formula for the Derivative

$$f'(x) = \lim_{z \to x} \frac{f(z) - f(x)}{z - x}.$$



Calculating Derivatives from the Definition

EXAMPLE 1 Differentiate $f(x) = \dfrac{x}{x-1}$.

Solution We use the definition of derivative, which requires us to calculate $f(x+h)$ and then subtract $f(x)$ to obtain the numerator in the difference quotient. We have

$$f(x) = \frac{x}{x-1} \quad \text{and} \quad f(x+h) = \frac{(x+h)}{(x+h)-1}, \text{ so}$$

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} \qquad \text{Definition}$$

$$= \lim_{h \to 0} \frac{\dfrac{x+h}{x+h-1} - \dfrac{x}{x-1}}{h}$$

$$= \lim_{h \to 0} \frac{1}{h} \cdot \frac{(x+h)(x-1) - x(x+h-1)}{(x+h-1)(x-1)} \qquad \frac{a}{b} - \frac{c}{d} = \frac{ad-cb}{bd}$$

$$= \lim_{h \to 0} \frac{1}{h} \cdot \frac{-h}{(x+h-1)(x-1)} \qquad \text{Simplify.}$$

$$= \lim_{h \to 0} \frac{-1}{(x+h-1)(x-1)} = \frac{-1}{(x-1)^2}. \qquad \text{Cancel } h \neq 0.$$

EXAMPLE 2

(a) Find the derivative of $f(x) = \sqrt{x}$ for $x > 0$.

(b) Find the tangent line to the curve $y = \sqrt{x}$ at $x = 4$.

Solution

(a) We use the alternative formula to calculate f':

$$f'(x) = \lim_{z \to x} \frac{f(z) - f(x)}{z - x}$$

$$= \lim_{z \to x} \frac{\sqrt{z} - \sqrt{x}}{z - x}$$

$$= \lim_{z \to x} \frac{\sqrt{z} - \sqrt{x}}{(\sqrt{z} - \sqrt{x})(\sqrt{z} + \sqrt{x})}$$

$$= \lim_{z \to x} \frac{1}{\sqrt{z} + \sqrt{x}} = \frac{1}{2\sqrt{x}}.$$

(b) The slope of the curve at $x = 4$ is


$$f'(4) = \frac{1}{2\sqrt{4}} = \frac{1}{4}.$$

The tangent is the line through the point (4, 2) with slope 1/4

The tangent is the line through the point (4, 2) with slope 1/4

$$y = 2 + \frac{1}{4}(x - 4)$$

$$y = \frac{1}{4}x + 1.$$



Notations

There are many ways to denote the derivative of a function $y = f(x)$, where the independent variable is x and the dependent variable is y. Some common alternative notations for the derivative are

$$f'(x) = y' = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx} f(x) = D(f)(x) = D_x f(x).$$

To indicate the value of a derivative at a specified number $x = a$, we use the notation

$$f'(a) = \left.\frac{dy}{dx}\right|_{x=a} = \left.\frac{df}{dx}\right|_{x=a} = \left.\frac{d}{dx} f(x)\right|_{x=a}.$$

For instance, in Example 2

$$f'(4) = \left.\frac{d}{dx}\sqrt{x}\right|_{x=4} = \left.\frac{1}{2\sqrt{x}}\right|_{x=4} = \frac{1}{2\sqrt{4}} = \frac{1}{4}.$$


Differentiable on an Interval; One-Sided Derivatives

A function $y = f(x)$ is **differentiable on an open interval** (finite or infinite) if it has a derivative at each point of the interval. It is **differentiable on a closed interval** $[a, b]$ if it is differentiable on the interior (a, b) and if the limits

$$\lim_{h \to 0^+} \frac{f(a + h) - f(a)}{h} \qquad \textbf{Right-hand derivative at } a$$

$$\lim_{h \to 0^-} \frac{f(b + h) - f(b)}{h} \qquad \textbf{Left-hand derivative at } b$$

exist at the endpoints



Remark

a function has a derivative at a point if and only if it has left-hand and right-hand derivatives there, and these one-sided derivatives are equal.

EXAMPLE 4 Show that the function $y = |x|$ is differentiable on $(-\infty, 0)$ and $(0, \infty)$ but has no derivative at $x = 0$.

$$\text{Right-hand derivative of } |x| \text{ at zero} = \lim_{h \to 0^+} \frac{|0 + h| - |0|}{h} = \lim_{h \to 0^+} \frac{|h|}{h}$$

$$= \lim_{h \to 0^+} \frac{h}{h} \qquad |h| = h \text{ when } h > 0$$

$$= \lim_{h \to 0^+} 1 = 1$$

$$\text{Left-hand derivative of } |x| \text{ at zero} = \lim_{h \to 0^-} \frac{|0 + h| - |0|}{h} = \lim_{h \to 0^-} \frac{|h|}{h}$$

$$= \lim_{h \to 0^-} \frac{-h}{h} \qquad |h| = -h \text{ when } h < 0$$

$$= \lim_{h \to 0^-} -1 = -1. \quad \blacksquare$$


EXAMPLE 5 In Example 2 we found that for $x > 0$,

$$\frac{d}{dx}\sqrt{x} = \frac{1}{2\sqrt{x}}.$$


We apply the definition to examine if the derivative exists at $x = 0$:

$$\lim_{h \to 0^+} \frac{\sqrt{0 + h} - \sqrt{0}}{h} = \lim_{h \to 0^+} \frac{1}{\sqrt{h}} = \infty.$$


When Does a Function *Not* Have a Derivative at a Point?




1. a *corner*, where the one-sided derivatives differ.



2. a *cusp*, where the slope of PQ approaches ∞ from one side and $-\infty$ from the other.



3. a *vertical tangent*, where the slope of PQ approaches ∞ from both sides or approaches $-\infty$ from both sides (here, $-\infty$).



4. a *discontinuity* (two examples shown).

Differentiable Functions Are Continuous

A function is continuous at every point where it has a derivative.

THEOREM 1—Differentiability Implies Continuity If f has a derivative at $x = c$, then f is continuous at $x = c$.

Remark

Theorem 1 says that if a function has a discontinuity at a point (for instance, a jump discontinuity), then it cannot be differentiable there. The greatest integer function $y = \lfloor x \rfloor$ fails to be differentiable at every integer $x = n$ (Example 4, Section 2.5).

The converse of Theorem 1 is false. A function need not have a derivative at a point where it is continuous, as we saw in Example 4.

Exercises 3.2

In Exercises 17–18, differentiate the functions. Then find an equation of the tangent line at the indicated point on the graph of the function.

17. $y = f(x) = \dfrac{8}{\sqrt{x-2}}, \quad (x, y) = (6, 4)$

$$f'(x) = \lim_{h\to 0}\frac{f(x+h)-f(x)}{h} = \lim_{h\to 0}\frac{\dfrac{8}{\sqrt{x+h-2}} - \dfrac{8}{\sqrt{x-2}}}{h}$$

$$= \lim_{h\to 0}\frac{\dfrac{8\sqrt{x-2} - 8\sqrt{x+h-2}}{\sqrt{x-2}\sqrt{x+h-2}}}{h} = 8\lim_{h\to 0}\frac{\sqrt{x-2}-\sqrt{x+h-2}}{h\left[\sqrt{x-2}\sqrt{x+h-2}\right]}$$

$$= 8\lim_{h\to 0}\frac{\sqrt{x-2}-\sqrt{x+h-2}}{h\left[\sqrt{x-2}\sqrt{x+h-2}\right]}\left(\frac{\sqrt{x-2}+\sqrt{x+h-2}}{\sqrt{x-2}+\sqrt{x+h-2}}\right)$$

$$= 8\lim_{h\to 0}\frac{x-2-(x+h-2)}{h\left[\sqrt{x-2}\sqrt{x+h-2}\right]\left(\sqrt{x-2}+\sqrt{x+h-2}\right)}$$

$$= 8\lim_{h\to 0}\frac{-h}{h\left[\sqrt{x-2}\sqrt{x+h-2}\right]\left(\sqrt{x-2}+\sqrt{x+h-2}\right)}$$

$$= 8\lim_{h\to 0}\frac{-1}{\left[\sqrt{x-2}\sqrt{x+h-2}\right]\left(\sqrt{x-2}+\sqrt{x+h-2}\right)}$$

$$\text{Slope} = f'(6) = \frac{-4}{(6-2)^{\frac{3}{2}}} = \frac{-4}{(4)^{\frac{3}{2}}} = -\frac{1}{2}$$

The equation of the tangent line is


$$\frac{y-4}{x-6} = -\frac{1}{2}$$

$$y = -\frac{1}{2}(x-6)+4 \quad \Rightarrow y = -\frac{1}{2}x + 7$$

One-Sided Derivatives

Compute the right-hand and left-hand derivatives as limits to show that the functions in Exercises 37–40 are not differentiable at the point P.

37.



$$\text{Right hand derivative} = \lim_{h\to 0^+}\frac{f(0+h)-f(0)}{h} = \lim_{h\to 0^+}\frac{f(h)-f(0)}{h} = \lim_{h\to 0^+}\frac{h-0}{h} = 1$$

$$\text{Left hand derivative} = \lim_{h\to 0^-}\frac{f(0+h)-f(0)}{h} = \lim_{h\to 0^-}\frac{f(h)-f(0)}{h} = \lim_{h\to 0^+}\frac{h^2-0}{h} = \lim_{h\to 0^+} h = 0$$

This implies that $f'(0)$ does not exist.