

Polyhedra (2)

Consider the following half-spaces in \mathbb{R}^2 :

$$-x_1 + 3x_2 \leq 12$$

$$x_1 + 2x_2 \leq 12$$

$$3x_1 - 2x_2 \leq 8$$

$$x_1 \geq 0$$

$$x_2 \geq 0$$

Questions

1. Draw a plot of their supporting hyper-planes.
2. Draw the polyhedron given by the intersection of the half-spaces. Is it a polytope?
3. What happens if we add constraint $x_1 \geq 5$ to the polyhedron? What if we add constraint $x_1 \leq 5$?

Solution

Given $a_{11}, a_{12}, a_{21}, a_{22}, b_1, b_2 \in \mathbb{R}$, if $(a_{11} a_{22} - a_{12} a_{21}) \neq 0$, we have:

$$\begin{cases} a_{11} x_1 + a_{12} x_2 = b_1 \\ a_{21} x_1 + a_{22} x_2 = b_2 \end{cases} \Rightarrow (x_1, x_2) = \left(\frac{b_1 a_{22} - a_{12} b_2}{a_{11} a_{22} - a_{12} a_{21}}, \frac{a_{11} b_2 - b_1 a_{21}}{a_{11} a_{22} - a_{12} a_{21}} \right)$$

1. The supporting hyper-planes associated with the given half-spaces are:

$$-x_1 + 3x_2 = 12$$


$$x_1 + 2x_2 = 12$$

$$3x_1 - 2x_2 = 8$$

$$x_1 = 0$$

$$x_2 = 0$$

Their plot is shown below.




We compute the intersection points:

$$\begin{cases} -x_1 + 3x_2 = 12 \\ x_1 + 2x_2 = 12 \end{cases} \Rightarrow (x_1, x_2) = \left(\frac{12 \cdot 2 - 3 \cdot 12}{(-1) \cdot 2 - 3 \cdot 1}, \frac{(-1) \cdot 12 - 12 \cdot 1}{(-1) \cdot 2 - 3 \cdot 1} \right) = \left(\frac{12}{5}, \frac{24}{5} \right)$$

$$\begin{cases} x_1 + 2x_2 = 12 \\ 3x_1 - 2x_2 = 8 \end{cases} \Rightarrow (x_1, x_2) = \left(\frac{12 \cdot (-2) - 2 \cdot 8}{1 \cdot (-2) - 2 \cdot 3}, \frac{1 \cdot 8 - 12 \cdot 3}{1 \cdot (-2) - 2 \cdot 3} \right) = \left(5, \frac{7}{2} \right)$$

2. The polyhedron, given by the intersection of the given half-spaces, is:



Since the polyhedron is bounded, it is also a polytope.

3. If we add constraint $x_1 \geq 5$ the polyhedron reduces to the point $\tilde{\mathbf{x}} = (\tilde{x}_1, \tilde{x}_2) = (5, \frac{7}{2})$. If we add constraint $x_1 \leq 5$ the polyhedron does not change since for any $\tilde{\mathbf{x}} \in P$ we have that $\tilde{x}_1 \leq 5$.