Flexible sexual appetite - Analysis Report

Malika Ihle

November 9, 2017

descriptive numbers

dead females during training

```
DeadFemales
##
##
      RedAverse RedPreference
##
AliveFemales
##
##
      RedAverse RedPreference
chisq.test(rbind(DeadFemales,AliveFemales))
  Pearson's Chi-squared test with Yates' continuity correction
##
## data: rbind(DeadFemales, AliveFemales)
## X-squared = 1.5161, df = 1, p-value = 0.2182
dead males during tests
Males tests excluded out of 107
##
## BlackMaleDied
                    FemaleDied FemaleStarved
                                               RedMaleDied
chisq.test(rbind(c(10,3),c(97,104)))
## Pearson's Chi-squared test with Yates' continuity correction
## data: rbind(c(10, 3), c(97, 104))
## X-squared = 2.9483, df = 1, p-value = 0.08597
```

duration to male consumption

```
number valid tests: 80
```

number test male consumed during video:15

percentage of test where consumption within first day: 27.5

Delay to consumption all valid tests:

```
## Min. 1st Qu. Median Mean 3rd Qu. Max. NA's
## 0.000 0.000 2.000 4.339 5.042 31.000 3
```

Bug test

```
##
## Call:
## glm(formula = AttackBugYN ~ Trt + Fcondition, family = "binomial",
       data = MY_TABLE_BugTest)
##
##
## Deviance Residuals:
                     Median
      Min
                1Q
                                  3Q
                                          Max
## -1.3822 -1.0170
                     0.9951
                              1.0342
                                       1.4017
##
## Coefficients:
                   Estimate Std. Error z value Pr(>|z|)
                    -0.4422
                                0.2748 -1.609
                                                 0.1076
## (Intercept)
                    0.8293
                                0.3704
## TrtRedPreference
                                         2.239
                                                 0.0252 *
## Fcondition
                    -8.7534
                               31.1049 -0.281
                                                 0.7784
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## (Dispersion parameter for binomial family taken to be 1)
##
      Null deviance: 173.28 on 124 degrees of freedom
## Residual deviance: 168.15 on 122 degrees of freedom
## AIC: 174.15
## Number of Fisher Scoring iterations: 4
```

Termite test

equality of motivation to feed

```
##
## Welch Two Sample t-test
##
## data: LogLatencyAttackTermiteRedPreference and LogLatencyAttackTermiteRedAverse
## t = 0.26756, df = 110.91, p-value = 0.7895
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## -0.5228022    0.6860234
## sample estimates:
## mean of x mean of y
## 2.092866    2.011255
```

Male test

```
##
## glm(formula = CannibalizedRedYN ~ Trt + DeltaMsize + DeltaMcondition,
##
       family = "binomial", data = MY_TABLE_MaleTestValid)
##
## Deviance Residuals:
##
      Min
              1Q
                    Median
                                  3Q
                                          Max
## -1.0056 -0.9148 -0.7826
                             1.3988
                                       1.7301
## Coefficients:
                   Estimate Std. Error z value Pr(>|z|)
##
## (Intercept)
                    -0.6140
                                0.3312 -1.854
                                                 0.0637 .
## TrtRedPreference -0.4199
                                0.5024 -0.836
                                                 0.4033
                    -2.5473
                                6.0305 -0.422
                                                 0.6727
## DeltaMsize
## DeltaMcondition
                    10.7190
                               97.0272
                                        0.110
                                                 0.9120
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## (Dispersion parameter for binomial family taken to be 1)
##
      Null deviance: 99.374 on 79 degrees of freedom
## Residual deviance: 98.333 on 76 degrees of freedom
```

```
## AIC: 106.33
##
## Number of Fisher Scoring iterations: 4
```

equality of male motivation to court

videos not watched yet

FID repeatability

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
    Approximation) [glmerMod]
## Family: binomial (logit)
## Formula: AttackRedYN ~ Trt + (1 | FID)
##
     Data: MY_TABLE_Step
##
##
       AIC
               BIC logLik deviance df.resid
##
     468.6
             480.1 -231.3
                             462.6
##
## Scaled residuals:
     Min 1Q Median
                             ЗQ
##
## -0.9812 -0.8297 -0.6957 1.0606 1.3677
##
## Random effects:
## Groups Name
                     Variance Std.Dev.
## FID (Intercept) 0.1698 0.4121
## Number of obs: 343, groups: FID, 125
## Fixed effects:
                  Estimate Std. Error z value Pr(>|z|)
## (Intercept)
                  ## TrtRedPreference 0.4318
                             0.2428 1.778 0.07535 .
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Correlation of Fixed Effects:
## TrtRdPrfrnc -0.753
## Bootstrap Progress:
##
## Repeatability estimation using the glmm method and logit link
##
## Repeatability for FID
## -----
## Link-scale approximation:
## R = 0.039
## SE = 0.037
## CI = [0, 0.129]
## P = 0.178 [LRT]
       NA [Permutation]
```

```
##
## Original-scale approximation:
## R = 0.039
## SE = 0.036
## CI = [0, 0.124]
## P = 0.178 [LRT]
      NA [Permutation]
##
## Repeatability for Fixed
## Link-scale approximation:
## R = 0.011
## SE = 0.012
## CI = [0, 0.047]
## P = NA [LRT]
##
       NA [Permutation]
##
## Original-scale approximation:
## R = 0.011
## SE = 0.012
## CI = [0, 0.044]
## P = NA [LRT]
##
       NA [Permutation]
```

Odds ratio

Bug

```
## OR 2.5 % 97.5 %
## 2.291711 1.117010 4.792772
```


Termite

```
## OR 2.5 % 97.5 %
## 1.4800000 0.7399342 3.0057548
```

Male

```
## OR 2.5 % 97.5 %
## 0.6571182 0.2400022 1.7463719
```

Termite graph against 50/50



Male graph against 50/50

