

# THE AMAZING

$$P(A/B) = \frac{P(B/A) P(A)}{P(B)}$$

## Probability and Statistics (Monsoon 2022)

### Lecture-24

# Thomas Bayes

Pawan Kumar  
IIIT, Hyderabad  
November 16, 2022

$P(B)$

$P(A)$



# ① Bayesian Inference

## Motivating Example

Prior and Posterior  
Maximum A Priori Estimation  
Minimum Mean Squared Error

# Table of Contents

- ① Bayesian Inference
  - Motivating Example
  - Prior and Posterior
  - Maximum Apriori Estimation
  - Minimum Mean Squared Error

## Recall: Statistical Inference...

## Recall: Statistical Inference...

Statistical Inference: Compare frequentist and Bayesian

General setup for a statistical inference problem:

## Recall: Statistical Inference...

Statistical Inference: Compare frequentist and Bayesian

General setup for a statistical inference problem: There is an unknown quantity that we would like to estimate. We get some data.

## Recall: Statistical Inference...

### Statistical Inference: Compare frequentist and Bayesian

**General setup for a statistical inference problem:** There is an unknown quantity that we would like to estimate. We get some data. From the data, we estimate the desired quantity.

#### Frequentist Approach

In that approach, the unknown quantity  $\theta$  is assumed to be a **fixed (non-random)** quantity that is to be estimated by the observed data.

## Recall: Statistical Inference...

### Statistical Inference: Compare frequentist and Bayesian

**General setup for a statistical inference problem:** There is an unknown quantity that we would like to estimate. We get some data. From the data, we estimate the desired quantity.

#### Frequentist Approach

In that approach, the unknown quantity  $\theta$  is assumed to be a **fixed (non-random)** quantity that is to be estimated by the observed data.

#### Bayesian Approach

In the Bayesian framework, we treat the unknown quantity,  $\Theta$ , as a random variable. More specifically, we assume that we have some **initial guess** about the distribution of  $\Theta$ . This distribution is called the **prior distribution**. After observing some data, we update the distribution of  $\Theta$  (based on the observed data).



# Table of Contents

- ① Bayesian Inference
  - Motivating Example
  - Prior and Posterior
  - Maximum Apriori Estimation
  - Minimum Mean Squared Error

## Motivating Example...

## Motivating Example...

Example (Motivating Example)

## Motivating Example...

### Example (Motivating Example)

Suppose that you would like to estimate the portion of voters in your town that plan to vote for Party A in an upcoming election.

## Motivating Example...

### Example (Motivating Example)

Suppose that you would like to estimate the portion of voters in your town that plan to vote for Party A in an upcoming election. To do so, you take a random sample of size  $n$  from the likely voters in the town.

## Motivating Example...

### Example (Motivating Example)

Suppose that you would like to estimate the portion of voters in your town that plan to vote for Party A in an upcoming election. To do so, you take a random sample of size  $n$  from the likely voters in the town. Since you have a limited amount of time and resources, your sample is relatively small.

## Motivating Example...

### Example (Motivating Example)

Suppose that you would like to estimate the portion of voters in your town that plan to vote for Party A in an upcoming election. To do so, you take a random sample of size  $n$  from the likely voters in the town. Since you have a limited amount of time and resources, your sample is relatively small. Specifically, suppose that  $n = 20$ . After doing your sampling, you find out that 6 people in your sample say they will vote for Party A.

## Some Observations...



## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

- In fact, in absence of any other data, that seems to be a reasonable estimate. However, you might feel that  $n = 20$  is too small.

## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

- In fact, in absence of any other data, that seems to be a reasonable estimate. However, you might feel that  $n = 20$  is too small.
- Thus, your guess is that the error in your estimation might be too high.

## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

- In fact, in absence of any other data, that seems to be a reasonable estimate. However, you might feel that  $n = 20$  is too small.
- Thus, your guess is that the error in your estimation might be too high.
- While thinking about this problem, you remember that the data from the previous election is available to you.

## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

- In fact, in absence of any other data, that seems to be a reasonable estimate. However, you might feel that  $n = 20$  is too small.
- Thus, your guess is that the error in your estimation might be too high.
- While thinking about this problem, you remember that the data from the previous election is available to you.
- You look at that data and find out that, in the previous election, 40% of the people in your town voted for Party A.

## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

- In fact, in absence of any other data, that seems to be a reasonable estimate. However, you might feel that  $n = 20$  is too small.
- Thus, your guess is that the error in your estimation might be too high.
- While thinking about this problem, you remember that the data from the previous election is available to you.
- You look at that data and find out that, in the previous election, 40% of the people in your town voted for Party A.
- How can you use this data to possibly improve your estimate of  $\theta$ ?
- Although the portion of votes for Party A changes from one election to another, the change is not usually very drastic.

## Some Observations...

- Let  $\theta$  be the true portion of voters in your town who plan to vote for Party A. You might want to estimate  $\theta$  as

$$\hat{\theta} = 6/20 = 0.3$$

- In fact, in absence of any other data, that seems to be a reasonable estimate. However, you might feel that  $n = 20$  is too small.
- Thus, your guess is that the error in your estimation might be too high.
- While thinking about this problem, you remember that the data from the previous election is available to you.
- You look at that data and find out that, in the previous election, 40% of the people in your town voted for Party A.
- How can you use this data to possibly improve your estimate of  $\theta$ ?
- Although the portion of votes for Party A changes from one election to another, the change is not usually very drastic.
- Therefore, given that in the previous election 40% of the voters voted for Party A, you might want to model the portion of votes for Party A in the next election as a random variable  $\Theta$  with a probability density function,  $f_{\Theta}(\theta)$ , that is mostly concentrated around  $\theta = 0.4$ .

## Some Observations...



## Some Observations...

- For example, you might want to choose the density such that

$$E[\Theta] = 0.4$$

## Some Observations...

- For example, you might want to choose the density such that

$$E[\Theta] = 0.4$$

- That is, before taking your random sample of size  $n = 20$ , this is your guess about the distribution of  $\Theta$ .

## Some Observations...

- For example, you might want to choose the density such that

$$E[\Theta] = 0.4$$

- That is, before taking your random sample of size  $n = 20$ , this is your guess about the distribution of  $\Theta$ .
- Therefore, you initially have the prior distribution  $f_{\Theta}(\theta)$ . Then you collect some data, shown by  $D$ .

## Some Observations...

- For example, you might want to choose the density such that

$$E[\Theta] = 0.4$$

- That is, before taking your random sample of size  $n = 20$ , this is your guess about the distribution of  $\Theta$ .
- Therefore, you initially have the prior distribution  $f_{\Theta}(\theta)$ . Then you collect some data, shown by  $D$ .
- More specifically, here your data is a random sample of size  $n=20$  voters, 6 of whom are voting for Party A.

## Some Observations...

- For example, you might want to choose the density such that

$$E[\Theta] = 0.4$$

- That is, before taking your random sample of size  $n = 20$ , this is your guess about the distribution of  $\Theta$ .
- Therefore, you initially have the prior distribution  $f_{\Theta}(\theta)$ . Then you collect some data, shown by  $D$ .
- More specifically, here your data is a random sample of size  $n=20$  voters, 6 of whom are voting for Party A.
- you can then proceed to find an updated distribution for  $\Theta$ , called the posterior distribution, using Bayes' rule:

$$f_{\Theta|D}(\theta|D) = \frac{P(D|\theta)f_{\Theta}(\theta)}{P(D)}. \quad (1)$$

- We can now use the posterior density,  $f_{\Theta|D}(\theta|D)$  to further draw inferences about  $\Theta$

## Bayesian Inference...

## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$

## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$
- 2 The **unknown** variable is modelled as a random variable  $X$ , with **prior distribution**  $f_X(x)$ , if  $X$  is continuous,  $P_X(x)$ , if  $X$  is discrete



## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$
- 2 The **unknown** variable is modelled as a random variable  $X$ , with **prior distribution**  $f_X(x)$ , if  $X$  is continuous,  $P_X(x)$ , if  $X$  is discrete
- 3 After observing the value of the random variable  $Y$ , we find the **posterior distribution** of  $X$ .

## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$
- 2 The **unknown** variable is modelled as a random variable  $X$ , with **prior distribution**  $f_X(x)$ , if  $X$  is continuous,  $P_X(x)$ , if  $X$  is discrete
- 3 After observing the value of the random variable  $Y$ , we find the **posterior distribution** of  $X$ . This is the conditional PDF (or PMF) of  $X$  given  $Y = y$ ,  $f_{X|Y}(x|y)$  or  $P_{X|Y}(x|y)$

## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$
- 2 The **unknown** variable is modelled as a random variable  $X$ , with **prior distribution**  $f_X(x)$ , if  $X$  is continuous,  $P_X(x)$ , if  $X$  is discrete
- 3 After observing the value of the random variable  $Y$ , we find the **posterior distribution** of  $X$ . This is the conditional PDF (or PMF) of  $X$  given  $Y = y$ ,  $f_{X|Y}(x|y)$  or  $P_{X|Y}(x|y)$
- 4 The **posterior distribution** is usually found using Bayes' formula.

## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$
- 2 The **unknown** variable is modelled as a random variable  $X$ , with **prior distribution**  $f_X(x)$ , if  $X$  is continuous,  $P_X(x)$ , if  $X$  is discrete
- 3 After observing the value of the random variable  $Y$ , we find the **posterior distribution** of  $X$ . This is the conditional PDF (or PMF) of  $X$  given  $Y = y$ ,  $f_{X|Y}(x|y)$  or  $P_{X|Y}(x|y)$
- 4 The **posterior distribution** is usually found using Bayes' formula. Using the posterior distribution, we can then find point or interval estimates of  $X$

## Bayesian Inference...

### Bayesian Inference: main ideas

- 1 The goal is to draw inferences about an unknown variable  $X$  by observing a related random variable  $Y$
- 2 The **unknown** variable is modelled as a random variable  $X$ , with **prior distribution**  $f_X(x)$ , if  $X$  is continuous,  $P_X(x)$ , if  $X$  is discrete
- 3 After observing the value of the random variable  $Y$ , we find the **posterior distribution** of  $X$ . This is the conditional PDF (or PMF) of  $X$  given  $Y = y$ ,  $f_{X|Y}(x|y)$  or  $P_{X|Y}(x|y)$
- 4 The **posterior distribution** is usually found using Bayes' formula. Using the posterior distribution, we can then find point or interval estimates of  $X$
- 5 Note that in the above setting,  $X$  or  $Y$  (or possibly both) could be random vectors

### Example

Solved example Let  $X \sim N(0, 1)$ . Suppose that we know

$$Y \mid X = x \sim N(x, 1).$$

Show that the posterior density of  $X$  given  $Y = y$ ,  $f_{X|Y}(x \mid y)$  is given by

$$X \mid Y = y \sim N\left(\frac{y}{2}, \frac{1}{2}\right).$$

# Table of Contents

## ① Bayesian Inference

Motivating Example

Prior and Posterior

Maximum Apriori Estimation

Minimum Mean Squared Error

## Prior and Posterior...



## Prior and Posterior...

### Prior and Posterior

- 1 Using our notation for PMF and CDF, we have

## Prior and Posterior...

### Prior and Posterior

- 1 Using our notation for PMF and CDF, we have

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)P_X(x)}{P_Y(y)}$$

## Prior and Posterior...

### Prior and Posterior

- 1 Using our notation for PMF and CDF, we have

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)P_X(x)}{P_Y(y)}$$

- 2 If  $X$  is continuous RV and  $Y$  is discrete,

## Prior and Posterior...

### Prior and Posterior

- 1 Using our notation for PMF and CDF, we have

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)P_X(x)}{P_Y(y)}$$

- 2 If  $X$  is continuous RV and  $Y$  is discrete,

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)f_X(x)}{P_Y(y)}$$

## Prior and Posterior...

### Prior and Posterior

- 1 Using our notation for PMF and CDF, we have

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)P_X(x)}{P_Y(y)}$$

- 2 If  $X$  is continuous RV and  $Y$  is discrete,

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)f_X(x)}{P_Y(y)}$$

- 3 To find the denominator  $P_Y(y)$  or  $f_Y(y)$ , we often use the law of total probability

## Prior and Posterior...

### Prior and Posterior

- 1 Using our notation for PMF and CDF, we have

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)P_X(x)}{P_Y(y)}$$

- 2 If  $X$  is continuous RV and  $Y$  is discrete,

$$P_{X|Y}(x|y) = \frac{P_{Y|X}(y|x)f_X(x)}{P_Y(y)}$$

- 3 To find the denominator  $P_Y(y)$  or  $f_Y(y)$ , we often use the law of total probability
- 4 Here  $f_{X|Y}(x|y)$  is called posterior distribution

## Solved Example...

## Solved Example...

### Example

Let  $X \sim \text{Uniform}(0, 1)$ .



## Solved Example...

### Example

Let  $X \sim \text{Uniform}(0, 1)$ . Suppose that we know

## Solved Example...

### Example

Let  $X \sim \text{Uniform}(0, 1)$ . Suppose that we know

$$Y \mid X = x \sim \text{Geometric}(x).$$

## Solved Example...

### Example

Let  $X \sim \text{Uniform}(0, 1)$ . Suppose that we know

$$Y \mid X = x \sim \text{Geometric}(x).$$

Find the posterior density of  $X$  given  $Y = 2$ ,  $f_{X|Y}(x|2)$

# Table of Contents

## ① Bayesian Inference

Motivating Example

Prior and Posterior

Maximum Apriori Estimation

Minimum Mean Squared Error

## Maximum Apriori Estimation (MAP)...

## Maximum Apriori Estimation (MAP)...

### Definiton of MAP

The **posterior distribution**,  $f_{X|Y}(x|y)$  (or  $P_{X|Y}(x|y)$ ), contains **all** the knowledge about the unknown quantity  $X$ .

## Maximum Apriori Estimation (MAP)...

### Definiton of MAP

The **posterior distribution**,  $f_{X|Y}(x|y)$  (or  $P_{X|Y}(x|y)$ ), contains **all** the knowledge about the unknown quantity  $X$ . Therefore, we can use the **posterior distribution** to find **point or interval estimates** of  $X$ .

## Maximum Apriori Estimation (MAP)...

### Definiton of MAP

The **posterior distribution**,  $f_{X|Y}(x|y)$  (or  $P_{X|Y}(x|y)$ ), contains **all** the knowledge about the unknown quantity  $X$ . Therefore, we can use the **posterior distribution** to find **point or interval estimates** of  $X$ . One way to obtain a point estimate is to choose the value of  $x$  that **maximizes** the **posterior PDF (or PMF)**.



## Maximum Apriori Estimation (MAP)...

### Definiton of MAP

The **posterior distribution**,  $f_{X|Y}(x|y)$  (or  $P_{X|Y}(x|y)$ ), contains **all** the knowledge about the unknown quantity  $X$ . Therefore, we can use the **posterior distribution** to find **point or interval estimates** of  $X$ . One way to obtain a point estimate is to choose the value of  $x$  that **maximizes** the **posterior PDF (or PMF)**. This is called the **maximum a posteriori (MAP) estimation**.

## Maximum Apriori Estimation (MAP)...

### Definiton of MAP

The **posterior distribution**,  $f_{X|Y}(x|y)$  (or  $P_{X|Y}(x|y)$ ), contains **all** the knowledge about the unknown quantity  $X$ . Therefore, we can use the **posterior distribution** to find **point or interval estimates** of  $X$ . One way to obtain a point estimate is to choose the value of  $x$  that **maximizes** the **posterior PDF (or PMF)**. This is called the **maximum a posteriori (MAP) estimation**.




Figure: Here  $\hat{x}_{MAP}$  is the value of  $X$  for which the posterior  $f_{X|Y}(x|y)$  is maximized

## Finding the MAP Estimate...

## Finding the MAP Estimate...

### MAP Estimate

We note that  $f_Y(y)$  does not depend on the value of  $x$ .

## Finding the MAP Estimate...

### MAP Estimate

We note that  $f_Y(y)$  does not depend on the value of  $x$ . Hence, to find the MAP estimate of  $X$  given that we have observed  $Y = y$ ,

## Finding the MAP Estimate...

### MAP Estimate

We note that  $f_Y(y)$  does not depend on the value of  $x$ . Hence, to find the MAP estimate of  $X$  given that we have observed  $Y = y$ , we find the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x).$$

## Finding the MAP Estimate...

### MAP Estimate

We note that  $f_Y(y)$  does not depend on the value of  $x$ . Hence, to find the MAP estimate of  $X$  given that we have observed  $Y = y$ , we find the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x).$$

Whenever,  $X$  or  $Y$  is discrete,

## Finding the MAP Estimate...

### MAP Estimate

We note that  $f_Y(y)$  does not depend on the value of  $x$ . Hence, to find the MAP estimate of  $X$  given that we have observed  $Y = y$ , we find the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x).$$

Whenever,  $X$  or  $Y$  is discrete, we replace PDF by its PMF.



## Example of MAP Estimate...

## Example of MAP Estimate...

Example (Example of MAP Estimate)

Let  $X$  be a continuous random variable with the following PDF:

## Example of MAP Estimate...

### Example (Example of MAP Estimate)

Let  $X$  be a continuous random variable with the following PDF:

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Example of MAP Estimate...

### Example (Example of MAP Estimate)

Let  $X$  be a continuous random variable with the following PDF:

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, let

## Example of MAP Estimate...

### Example (Example of MAP Estimate)

Let  $X$  be a continuous random variable with the following PDF:

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, let

$$Y|X = x \sim \text{Geometric}(x).$$

## Example of MAP Estimate...

### Example (Example of MAP Estimate)

Let  $X$  be a continuous random variable with the following PDF:

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, let

$$Y|X = x \sim \text{Geometric}(x).$$

Find the **MAP estimate** of  $X$  given  $Y = 3$ .

## Comparison of MAP to ML Estimator...

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value.



## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The maximum likelihood (ML) estimate of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The maximum likelihood (ML) estimate of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The ML estimate is shown by  $\hat{x}_{ML}$ .

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The maximum likelihood (ML) estimate of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The ML estimate is shown by  $\hat{x}_{ML}$ .

- 2 On the other hand,

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The **maximum likelihood (ML) estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The **ML estimate** is shown by  $\hat{x}_{ML}$ .

- 2 On the other hand, the **MAP estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x)$$

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The **maximum likelihood (ML) estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The **ML estimate** is shown by  $\hat{x}_{ML}$ .

- 2 On the other hand, the **MAP estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x)$$

- 3 The two expressions are somewhat similar.

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The **maximum likelihood (ML) estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The **ML estimate** is shown by  $\hat{x}_{ML}$ .

- 2 On the other hand, the **MAP estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x)$$

- 3 The two expressions are somewhat similar. The MAP has one extra term  $f_X(x)$

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The **maximum likelihood (ML) estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The **ML estimate** is shown by  $\hat{x}_{ML}$ .

- 2 On the other hand, the **MAP estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x)$$

- 3 The two expressions are somewhat similar. The MAP has one extra term  $f_X(x)$
- 4 If  $X$  is uniformly distributed over a finite interval,

## Comparison of MAP to ML Estimator...

### Comparison of ML and MAP

- 1 Let  $Y = y$  be observed value. The **maximum likelihood (ML) estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x).$$

The **ML estimate** is shown by  $\hat{x}_{ML}$ .

- 2 On the other hand, the **MAP estimate** of  $X$  is the value of  $x$  that maximizes

$$f_{Y|X}(y|x)f_X(x)$$

- 3 The two expressions are somewhat similar. The MAP has one extra term  $f_X(x)$
- 4 If  $X$  is uniformly distributed over a finite interval, then **ML and MAP estimate** is same



## Solved Example...

## Solved Example...

### Example

Suppose that the signal  $X \sim N(0, \sigma_X^2)$  is transmitted over a communication channel.

## Solved Example...

### Example

Suppose that the signal  $X \sim N(0, \sigma_X^2)$  is transmitted over a communication channel. Assume that the received signal is given by

$$Y = X + W,$$

where  $W \sim N(0, \sigma_W^2)$  is independent of  $X$ .

## Solved Example...

### Example

Suppose that the signal  $X \sim N(0, \sigma_X^2)$  is transmitted over a communication channel. Assume that the received signal is given by

$$Y = X + W,$$

where  $W \sim N(0, \sigma_W^2)$  is independent of  $X$ .

- 1 Find the ML estimate of  $X$ , given  $Y = y$  is observed

## Solved Example...

### Example

Suppose that the signal  $X \sim N(0, \sigma_X^2)$  is transmitted over a communication channel. Assume that the received signal is given by

$$Y = X + W,$$

where  $W \sim N(0, \sigma_W^2)$  is independent of  $X$ .

- 1 Find the ML estimate of  $X$ , given  $Y = y$  is observed
- 2 Find the MAP estimate of  $X$ , given  $Y = y$  is observed

# Table of Contents

## ① Bayesian Inference

Motivating Example

Prior and Posterior

Maximum Apriori Estimation

Minimum Mean Squared Error

## Minimum Mean Squared Error (MMSE) Estimation...

## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .



## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .
- 2 To find a point estimate of  $X$ , we can just choose a summary statistic of the posterior such as its mean, median, or mode

## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .
- 2 To find a point estimate of  $X$ , we can just choose a summary statistic of the posterior such as its mean, median, or mode
- 3 If we choose the mode (the value of  $x$  that maximizes  $f_{X|Y}(x|y)$ ), we obtain the MAP estimate of  $X$

## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .
- 2 To find a point estimate of  $X$ , we can just choose a summary statistic of the posterior such as its mean, median, or mode
- 3 If we choose the mode (the value of  $x$  that maximizes  $f_{X|Y}(x|y)$ ), we obtain the MAP estimate of  $X$
- 4 Another possibility would be to choose the posterior mean, i.e.,  $\hat{x} = E[X|Y = y]$

## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .
- 2 To find a point estimate of  $X$ , we can just choose a summary statistic of the posterior such as its mean, median, or mode
- 3 If we choose the mode (the value of  $x$  that maximizes  $f_{X|Y}(x|y)$ ), we obtain the MAP estimate of  $X$
- 4 Another possibility would be to choose the posterior mean, i.e.,  $\hat{x} = E[X|Y = y]$

The **minimum mean squared error** (MMSE) estimate of the random variable  $X$ ,

## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .
- 2 To find a point estimate of  $X$ , we can just choose a summary statistic of the posterior such as its mean, median, or mode
- 3 If we choose the mode (the value of  $x$  that maximizes  $f_{X|Y}(x|y)$ ), we obtain the MAP estimate of  $X$
- 4 Another possibility would be to choose the posterior mean, i.e.,  $\hat{x} = E[X|Y = y]$

The **minimum mean squared error** (MMSE) estimate of the random variable  $X$ , given that we have observed  $Y = y$ , is given by

## Minimum Mean Squared Error (MMSE) Estimation...

### MMSE

- 1 The posterior distribution,  $f_{X|Y}(x|y)$ , contains all the knowledge that we have about the unknown quantity  $X$ .
- 2 To find a point estimate of  $X$ , we can just choose a summary statistic of the posterior such as its mean, median, or mode
- 3 If we choose the mode (the value of  $x$  that maximizes  $f_{X|Y}(x|y)$ ), we obtain the MAP estimate of  $X$
- 4 Another possibility would be to choose the posterior mean, i.e.,  $\hat{x} = E[X|Y = y]$

The **minimum mean squared error** (MMSE) estimate of the random variable  $X$ , given that we have observed  $Y = y$ , is given by

$$\hat{x}_M = E[X|Y = y].$$

## Example of MMSE Computation...

## Example of MMSE Computation...

### Example

Let  $X$  be a continuous random variable with the following PDF



## Example of MMSE Computation...

### Example

Let  $X$  be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Example of MMSE Computation...

### Example

Let  $X$  be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, we are given that

## Example of MMSE Computation...

### Example

Let  $X$  be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, we are given that

$$f_{Y|X}(y|x) = \begin{cases} 2xy - x + 1 & \text{if } 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Example of MMSE Computation...

### Example

Let  $X$  be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, we are given that

$$f_{Y|X}(y|x) = \begin{cases} 2xy - x + 1 & \text{if } 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Find the **MMSE estimate** of  $X$ , given  $Y = y$  is observed.

## Example of MMSE Computation...

### Example

Let  $X$  be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, we are given that

$$f_{Y|X}(y|x) = \begin{cases} 2xy - x + 1 & \text{if } 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Find the **MMSE estimate** of  $X$ , given  $Y = y$  is observed.

## Mean Squared Error (MSE) ...

## Mean Squared Error (MSE) ...

### MSE

Let  $\hat{X} = g(Y)$  be an estimator of the random variable  $X$ , given that we have observed the random variable  $Y$ .

## Mean Squared Error (MSE) ...

### MSE

Let  $\hat{X} = g(Y)$  be an estimator of the random variable  $X$ , given that we have observed the random variable  $Y$ . The **mean squared error (MSE)** of this estimator is defined as



## Mean Squared Error (MSE) ...

### MSE

Let  $\hat{X} = g(Y)$  be an estimator of the random variable  $X$ , given that we have observed the random variable  $Y$ . The **mean squared error (MSE)** of this estimator is defined as

$$E[(X - \hat{X})^2] = E[(X - g(Y))^2]$$

## Mean Squared Error (MSE) ...

### MSE

Let  $\hat{X} = g(Y)$  be an estimator of the random variable  $X$ , given that we have observed the random variable  $Y$ . The **mean squared error (MSE)** of this estimator is defined as

$$E[(X - \hat{X})^2] = E[(X - g(Y))^2]$$

The **MMSE estimator** of  $X$ ,

## Mean Squared Error (MSE) ...

### MSE

Let  $\hat{X} = g(Y)$  be an estimator of the random variable  $X$ , given that we have observed the random variable  $Y$ . The **mean squared error (MSE)** of this estimator is defined as

$$E[(X - \hat{X})^2] = E[(X - g(Y))^2]$$

The **MMSE estimator** of  $X$ ,

$$\hat{X}_M = E[X|Y],$$

## Mean Squared Error (MSE) ...

### MSE

Let  $\hat{X} = g(Y)$  be an estimator of the random variable  $X$ , given that we have observed the random variable  $Y$ . The **mean squared error (MSE)** of this estimator is defined as

$$E[(X - \hat{X})^2] = E[(X - g(Y))^2]$$

The **MMSE estimator** of  $X$ ,

$$\hat{X}_M = E[X|Y],$$

has the **lowest MSE** among **all** possible estimators.