

November 11, 2022

## Tutorial 8 - Probability and Statistics

---

### Problem 1

The following data is from a random sample: 5, 1, 3, 3, 8.  
Compute the sample mean and sample standard deviation.

### Problem 2

Let  $X_1, X_2, X_3$  be a random sample of size  $n = 3$  from a distribution with the geometric probability mass function:

$$f(x) = \left(\frac{3}{4}\right) \left(\frac{1}{4}\right)^{x-1}$$

for  $x = 1, 2, 3, \dots$ . What is  $P(\max X_i \leq 2)$ ?

### Problem 3

If  $X_i$  is a Bernoulli random variable with parameter  $p$ , then:

$$\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$$

is the maximum likelihood estimator (MLE) of  $p$ . Is the MLE of  $p$  an unbiased estimator of  $p$ ?

### Problem 4

If  $X_i$  are normally distributed random variables with mean  $\mu$  and variance  $\sigma^2$ , then:

$$\hat{\mu} = \frac{\sum X_i}{n} = \bar{X} \text{ and } \hat{\sigma}^2 = \frac{\sum (X_i - \bar{X})^2}{n}$$

are the maximum likelihood estimators of  $\mu$  and  $\sigma^2$ , respectively. Are the MLEs unbiased for their respective parameters?

## Tutorial 8 - Probability and Statistics

---

### Problem 5

Let  $Y_1 < Y_2 < Y_3 < Y_4 < Y_5 < Y_6$  be the order statistics associated with  $n = 6$  independent observations each from the distribution with probability density function:

$$f(x) = \frac{1}{2}x \quad ; \text{ for } 0 < x < 2$$

What is the probability that the next-to-largest order statistic, that is,  $Y_5$ , is less than 1?

### Problem 6

**Example 2:** Suppose  $X_1, X_2, \dots, X_n$  are i.i.d. random variables with density function  $f(x|\sigma) = \frac{1}{2\sigma} \exp\left(-\frac{|x|}{\sigma}\right)$ , please find the maximum likelihood estimate of  $\sigma$ .

### Problem 7

**Example 3. Light bulbs**

Suppose that the lifetime of *Badger* brand light bulbs is modeled by an exponential distribution with (unknown) parameter  $\lambda$ . We test 5 bulbs and find they have lifetimes of 2, 3, 1, 3, and 4 years, respectively. What is the MLE for  $\lambda$ ?

### Problem 8

Suppose our data  $x_1, \dots, x_n$  are independently drawn from a uniform distribution  $U(a, b)$ . Find the MLE estimate for  $a$  and  $b$ .

## Tutorial 8 - Probability and Statistics

---

### Problem 9

**Example 7. Hardy-Weinberg.** Suppose that a particular gene occurs as one of two alleles ( $A$  and  $a$ ), where allele  $A$  has frequency  $\theta$  in the population. That is, a random copy of the gene is  $A$  with probability  $\theta$  and  $a$  with probability  $1 - \theta$ . Since a diploid genotype consists of two genes, the probability of each genotype is given by:

genotype	AA	Aa	aa
probability	$\theta^2$	$2\theta(1 - \theta)$	$(1 - \theta)^2$

Suppose we test a random sample of people and find that  $k_1$  are  $AA$ ,  $k_2$  are  $Aa$ , and  $k_3$  are  $aa$ . Find the MLE of  $\theta$ .

### Problem 10

**Example 2.2.2 (Weibull with known  $\alpha$ )**  $\{Y_i\}$  are iid random variables, which follow a Weibull distribution, which has the density

$$\frac{\alpha y^{\alpha-1}}{\theta^\alpha} \exp(-(y/\theta)^\alpha) \quad \theta, \alpha > 0.$$

Suppose that  $\alpha$  is known, but  $\theta$  is unknown. Our aim is to find the MLE of  $\theta$ .

## Tutorial 8 - Probability and Statistics

### Problem 11

Let  $X_1, X_2, X_3, \dots, X_n$  be a random sample from the following distribution:

$$f_X(x) = \begin{cases} \theta \left(x - \frac{1}{2}\right) + 1 & \text{for } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

where  $\theta \in [-2, 2]$  is an unknown parameter. We define the estimator  $\hat{\Theta}_n$  as

$$\hat{\Theta}_n = 12\bar{X} - 6$$

to estimate  $\theta$ .

- (a) Is  $\hat{\Theta}_n$  an unbiased estimator of  $\theta$ ?
- (b) Is  $\hat{\Theta}_n$  a consistent estimator of  $\theta$ ?
- (c) Find the mean squared error (MSE) of  $\hat{\Theta}_n$ .

### Problem 12

Let  $X$  be one observation from a  $N(0, \sigma^2)$  distribution.

- (a) Find an unbiased estimator of  $\sigma^2$ .
- (b) Find the log likelihood,  $\log(L(x; \sigma^2))$ , using

$$f_X(x; \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{x^2}{2\sigma^2}\right\}$$

as the PDF.

- (c) Find the Maximum Likelihood Estimate (MLE) for the standard deviation  $\sigma$ ,  $\hat{\sigma}_{ML}$ .

## Tutorial 8 - Probability and Statistics

---

### Problem 13

15. Let  $X_1, X_2, X_3, X_4, X_5$  be a random sample from a  $N(\mu, 1)$  distribution, where  $\mu$  is unknown. Suppose that we have observed the following values

5.45, 4.23, 7.22, 6.94, 5.98

We would like to decide between

$$H_0: \mu = \mu_0 = 5,$$

$$H_1: \mu \neq 5.$$

- (a) Define a test statistic to test the hypotheses and draw a conclusion assuming  $\alpha = 0.05$ .
- (b) Find a 95% confidence interval around  $\bar{X}$ . Is  $\mu_0$  included in the interval? How does the exclusion of  $\mu_0$  in the interval relate to the hypotheses we are testing?

## Tutorial 8 - Probability and Statistics

---

### Problem 14

Let  $X_1, X_2, \dots, X_{150}$  be a random sample from an unknown distribution. After observing this sample, the sample mean and the sample variance are calculated to be as follows:

$$\bar{X} = 52.28, \quad S^2 = 30.9$$

Design a level 0.05 test to choose between

$$H_0: \mu = 50,$$

$$H_1: \mu > 50.$$

Do you accept or reject  $H_0$ ?

### Problem 15

Let  $X_1, X_2, \dots, X_{121}$  be a random sample from an unknown distribution. After observing this sample, the sample mean and the sample variance are calculated to be as follows:

$$\bar{X} = 29.25, \quad S^2 = 20.7$$

Design a test to decide between

$$H_0: \mu = 30,$$

$$H_1: \mu < 30,$$

and calculate the  $P$ -value for the observed data.

## Tutorial 8 - Probability and Statistics

---

### Problem 16

*Example 2.9* Let  $X_1, X_2, \dots, X_n$  be a random sample from  $f(x, \alpha, \beta) = \beta e^{-\beta(x-\alpha)}$ ;  $\alpha \leq x < \infty$  and  $\beta > 0$ .  
Find MLE's of  $\alpha, \beta$ .

### Problem 17

*Example 2.10* Let  $X_1, X_2, \dots, X_n$  be a random sample from  $f(x, \alpha, \beta) = \begin{cases} \frac{1}{\beta-\alpha}, & \alpha \leq x \leq \beta \\ 0, & \text{Otherwise} \end{cases}$   
(a) Show that the MLE of  $(\alpha, \beta)$  is  $(\text{Min } X_i, \text{Max } X_i)$ .  
(b) Also find the estimators of  $\alpha$  and  $\beta$  by the method of moments.

### Problem 18

*Example 5.2* Let  $X_1, X_2, \dots, X_n$  be a random sample from  $N(\mu, \sigma^2)$ . Find  $(1 - \alpha)$  level confidence interval for  $\sigma^2$  when (i)  $\mu$  is known and (ii)  $\mu$  is unknown.

### Problem 19

*Example 5.3* Let  $X_1, X_2, \dots, X_n$  be a random sample from density function  $f(x|\theta) = \left(\frac{1}{\theta}\right), 0 < x < \theta$ . Find  $100(1 - \alpha)\%$  confidence interval of  $\theta$ .

November 11, 2022

## Tutorial 8 - Probability and Statistics

---

### Problem 20

There are two candidates in a presidential election: Candidate A and Candidate B. Let  $\theta$  be the portion of people who plan to vote for Candidate A. Our goal is to find a confidence interval for  $\theta$ . Specifically, we choose a random sample (with replacement) of  $n$  voters and ask them if they plan to vote for Candidate A. Our goal is to estimate the  $\theta$  such that the margin of error is 3 percentage points. Assume a 95% confidence level. That is, we would like to choose  $n$  such that

$$P\left(\bar{X} - 0.03 \leq \theta \leq \bar{X} + 0.03\right) \geq 0.95,$$

where  $\bar{X}$  is the portion of people in our random sample that say they plan to vote for Candidate A. How large does  $n$  need to be?