

MA 6.101
Probability and Statistics

Tejas Bodas

Assistant Professor, IIIT Hyderabad

Agenda for the next two lectures

- ▶ Intro to Stochastic Simulation
 - ▶ We will generate samples from discrete or continuous r.v's using samples from uniform distribution.
- ▶ Limit theorems for Convergence of random variables
 - ▶ Sure convergence
 - ▶ Almost sure convergence & SLLN
 - ▶ Convergence in probability
 - ▶ Convergence in r^{th} mean
 - ▶ Weak Convergence or Convergence in distribution & CLT


Generate samples using uniform distribution

Our aim: Obtain samples from a discrete random variable

- ▶ Suppose you have access to samples from a uniform random variable U over support $[0, 1]$.

- ▶

```
import numpy as np
import matplotlib.pyplot as plt
uni_samples = np.random.uniform(0, 1, 5000)
plt.hist(uni_samples, bins = 10, density = True)
plt.show()
```



- ▶ *uni_samples* is a vector of 5000 realizations of uniform random variable U .
- ▶ You can also see it as a realization of $U_1, U_2, \dots, U_{5000}$ i.i.d uniform variables.


How to simulate a dice using these samples?

- ▶ Can you use these 5000 samples and convert them into outcomes of a dice ?

```
t=0
dice_samples=np.zeros(5000)
for u in uni_samples:
    if u < 1/6:
        dice_sample = 1
    if 1/6 < u < 2/6:
        dice_sample = 2
    if 2/6 < u < 3/6:
        dice_sample = 3
    if 3/6 < u < 4/6:
        dice_sample = 4
    if 4/6 < u < 5/6:
        dice_sample = 5
    if 5/6 < u < 6/6:
        dice_sample = 6
    dice_samples[t] = dice_sample
    t = t+1
plt.hist(dice_samples, bins = 6, density = True)
```

▶ [0.02, 0.8, 0.6, 0.03]

▶ [1, 5, 4, 1]



Our aim: Obtain samples from a discrete random variable

- ▶ Consider a discrete random variable X with support set $\{x_0, x_1, \dots\}$ and pmf $p_X(x_j) = p_j$ for $j = 0, 1, \dots$ such that $\sum_j p_j = 1$.
- ▶ Cardinality of the support set of X could be finite or infinite.
- ▶ Our aim: Create i.i.d. samples of r.v. X using i.i.d. random samples of U .
- ▶ We shall now formally see the **inverse transform method** to do this.

The inverse transform method

- ▶ **Aim:** We wish to create i.i.d. samples of a discrete r.v. X with $p_X(x_j) = p_j$ using i.i.d. samples of a uniform r.v. U over $[0, 1]$.
- ▶ Let $u \in [0, 1]$ be a realization of r.v. U . Then the corresponding sample of X is generated as follows

$$X = \begin{cases} x_0 & \text{if } u < p_0 \\ x_1 & \text{if } p_0 \leq u < p_0 + p_1 \\ x_2 & \text{if } p_0 + p_1 \leq u < p_0 + p_1 + p_2 \\ \vdots & \\ x_j & \text{if } \sum_{i=0}^{j-1} p_i \leq u < \sum_{i=0}^j p_i \\ \vdots & \end{cases}$$

- ▶ Why is this method correct?

The inverse transform method

- ▶ A sample of X is generated using the sample of U as follows

$$X = x_j \quad \text{if} \quad \sum_{i=0}^{j-1} p_i \leq U < \sum_{i=0}^j p_i$$

- ▶ Why the name “inverse transform method”?
 - ▶ Recall that $\{x_0, x_1, x_2, \dots\}$ is the support set of X and without loss of generality (WLOG), suppose $x_0 < x_1 < x_2 < \dots$
 - ▶ Let $F_X(x) := \mathbb{P}[X \leq x]$ denote the cdf of X and thus we have

$$F_X(x_j) = \sum_{i=0}^j p_i$$

- ▶ This implies that

$$X = x_j \quad \text{if} \quad F_X(x_{j-1}) \leq U < F_X(x_j) \quad (\text{implying } p_X(x_j) = p_j)$$

- ▶ After generating a random number U , we determine the value of X by finding the interval $[F_X(x_{j-1}), F_X(x_j))$ in which u lies.
- ▶ We are thus finding the inverse of $F_X(U)$!

How to generate samples of a continuous random variable

(Using samples of a continuous uniform variable over $[0, 1]$)

Our aim: Obtain samples from a continuous random variable

- ▶ Suppose you have access to samples from a uniform random variable U over support $[0, 1]$. (We will not study how to generate such samples.)
- ▶ Consider a continuous random variable X with support set \mathcal{X} and let $F_X(x)$ denotes its cdf.
- ▶ Support set of X could be arbitrary.
- ▶ Our aim: Create i.i.d. samples of r.v. X using i.i.d. random samples of U .
- ▶ We shall again see the **inverse transform method** to do this.

Sampling from continuous random variables

Lemma

Let U be uniform random variable over $[0, 1]$. Consider any continuous cdf $F(\cdot)$. Consider a random variable X defined as follows

$$X := F^{-1}(U)$$

Then the cdf of X is $F(\cdot)$.

Proof:

- ▶ Let $F_X(x)$ be the cdf of X , i.e., $F_X(x) := \mathbb{P}[X \leq x]$. Then

$$\begin{aligned} F_X(x) &= \mathbb{P}[F^{-1}(U) \leq x] \\ &= \mathbb{P}[U \leq F(x)] \\ &= F(x) \end{aligned}$$

Sampling from continuous random variables

Lemma

Let U be uniform random variable over $[0, 1]$. Consider any continuous cdf $F(\cdot)$. Consider a random variable X defined as follows

$$X := F^{-1}(U)$$

Then the cdf of X is $F(\cdot)$.

- ▶ Using this lemma, how to generate samples of a continuous random variable X using samples of uniform random variable U ?
- ▶ **Answer:** Draw $u \sim U$ and obtain $F^{-1}(u)$. This is a sample from X .
- ▶ Do you observe anything “special” about this lemma?

Application in data analysis

- ▶ Lemma: $X = F^{-1}(U)$ has distribution $F(\cdot)$.
- ▶ What will be cdf of a random variable $Y = F(X)$? **Uniform!**
- ▶ A consequence of this lemma is that $F(X)$ is a uniform distribution.
- ▶ This property is known as “probability integral transform or universality of uniform”.
- ▶ This property is used to test whether a set of observations can be modelled as arising from a specified distribution $G(\cdot)$ or not.
 - ▶ Given set of data samples s_1, s_2, \dots, s_n , plot $G(s_i)$ for different samples.
 - ▶ If these points are spread uniformly over the interval $[0, 1]$ then it indicates that the samples are indeed from $G(\cdot)$.

Stochastic Simulation

- ▶ This was a brief introduction to this area of stochastic simulation.
- ▶ Refer the book Simulation by Sheldon Ross!
- ▶ Some popular techniques in simulation are:
 - ▶ The inverse transform method
 - ▶ Accept-Reject method (rejection sampling)
 - ▶ Importance sampling
 - ▶ Markov Chain Monte Carlo (MCMC) methods
 - ▶ Hasting-Metropolis algorithm
 - ▶ Gibbs sampling
 - ▶ Slice sampling