

THE AMAZING

$$P(A/B) = \frac{P(B/A) P(A)}{P(B)}$$

Probability and Statistics (Monsoon 2022)

Lecture-24

Thomas Bayes

Pawan Kumar
IIIT, Hyderabad
November 17, 2022

$P(B)$

$P(A)$



① Statistical Inference
Interval Estimation and Confidence Level

Hypothesis Testing for the Mean
P-Values

Table of Contents

- ① Statistical Inference
 - Interval Estimation and Confidence Level
 - Hypothesis Testing for the Mean
 - P-Values

Table of Contents

- ① Statistical Inference
 - Interval Estimation and Confidence Level
 - Hypothesis Testing for the Mean
 - P-Values

Trade-off between α and β

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .
 - if we want to decrease the probability of type I error (α), then the probability of type II error (β) increases, and vice versa.

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .
 - if we want to decrease the probability of type I error (α), then the probability of type II error (β) increases, and vice versa.
- When we look at our analysis in previous Example, we found

$$\begin{aligned}\alpha &= 1 - \Phi(3c), \\ \beta &= \Phi(3(c - 1)).\end{aligned}$$

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .
 - if we want to decrease the probability of type I error (α), then the probability of type II error (β) increases, and vice versa.
- When we look at our analysis in previous Example, we found

$$\begin{aligned}\alpha &= 1 - \Phi(3c), \\ \beta &= \Phi(3(c - 1)).\end{aligned}$$

- Since Φ is an increasing function, we have

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .
 - if we want to decrease the probability of type I error (α), then the probability of type II error (β) increases, and vice versa.
- When we look at our analysis in previous Example, we found

$$\begin{aligned}\alpha &= 1 - \Phi(3c), \\ \beta &= \Phi(3(c - 1)).\end{aligned}$$

- Since Φ is an increasing function, we have
 - 1 If we make c larger, α becomes smaller, and β becomes larger.

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .
 - if we want to decrease the probability of type I error (α), then the probability of type II error (β) increases, and vice versa.
- When we look at our analysis in previous Example, we found

$$\alpha = 1 - \Phi(3c),$$
$$\beta = \Phi(3(c - 1)).$$

- Since Φ is an increasing function, we have
 - 1 If we make c larger, α becomes smaller, and β becomes larger.
 - 2 On the other hand, if we make c smaller, α becomes larger, and β becomes smaller.

Trade-off between α and β

- Since α and β indicate error probabilities, we would ideally like both of them to be small.
- There is in fact a trade-off between α and β .
 - if we want to decrease the probability of type I error (α), then the probability of type II error (β) increases, and vice versa.
- When we look at our analysis in previous Example, we found

$$\alpha = 1 - \Phi(3c),$$
$$\beta = \Phi(3(c - 1)).$$

- Since Φ is an increasing function, we have
 - 1 If we make c larger, α becomes smaller, and β becomes larger.
 - 2 On the other hand, if we make c smaller, α becomes larger, and β becomes smaller.




Table of Contents

① Statistical Inference

Interval Estimation and Confidence Level

Hypothesis Testing for the Mean

P-Values

Hypothesis Testing for Mean




Figure: Hypothesis Testing for Mean.

Definition (Two sided hypothesis test)

- Consider a random sample X_1, X_2, \dots, X_n from a distribution.
- Our goal is to make inference about the mean of the distribution μ .
- Two sided hypothesis test: Decide between the following hypotheses:
 - $H_0 : \mu = \mu_0$
 - $H_1 : \mu \neq \mu_0$

Here, null hypothesis is simple hypothesis, alternate hypothesis is **two-sided** ($\mu < \mu_0, \mu > \mu_0$)

Definition (Two sided hypothesis test)

- Consider a random sample X_1, X_2, \dots, X_n from a distribution.
- Our goal is to make inference about the mean of the distribution μ .
- Two sided hypothesis test: Decide between the following hypotheses:
 - $H_0 : \mu = \mu_0$
 - $H_1 : \mu \neq \mu_0$

Here, null hypothesis is simple hypothesis, alternate hypothesis is **two-sided** ($\mu < \mu_0, \mu > \mu_0$)

Definition (One-Sided Hypothesis)

- The second and the third cases are **one-sided tests**. More specifically

$$H_0 : \mu \leq \mu_0, \quad H_1 : \mu > \mu_0.$$

- Here, both H_0 and H_1 are one-sided, so we call this test a **one-sided test**.
- The third case is similar

$$H_0 : \mu \geq \mu_0, \quad H_1 : \mu < \mu_0.$$

Statistics and Test Statistics

- In all cases, we use sample mean as statistics.

Statistics and Test Statistics

- In all cases, we use sample mean as statistics.

$$\bar{X} = \frac{X_1 + X_2 + \cdots + X_n}{n}$$

Statistics and Test Statistics

- In all cases, we use sample mean as statistics.

$$\bar{X} = \frac{X_1 + X_2 + \cdots + X_n}{n}$$

- Furthermore, if we know the variance $\text{Var}(X_i) = \sigma^2$, define test statistics

$$W(X_1, X_2, \cdots, X_n) = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Statistics and Test Statistics

- In all cases, we use sample mean as statistics.

$$\bar{X} = \frac{X_1 + X_2 + \cdots + X_n}{n}$$

- Furthermore, if we know the variance $\text{Var}(X_i) = \sigma^2$, define test statistics

$$W(X_1, X_2, \cdots, X_n) = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

- If we don't know the variance, then use sample variance

$$W(X_1, X_2, \cdots, X_n) = \frac{\bar{X} - \mu_0}{S/\sqrt{n}},$$

where S is sample standard deviation

$$S = \sqrt{\frac{1}{n-1} \sum_{k=1}^n (X_k - \bar{X})^2} = \sqrt{\frac{1}{n-1} \left(\sum_{k=1}^n X_k^2 - n\bar{X}^2 \right)}. \quad (1)$$

Plan for two-sided tests for the mean

Plan for two-sided tests for the mean

- Given a random sample X_1, X_2, \dots, X_n from a distribution. Let $\mu = E[X_i]$. Our goal is to decide between

$$H_0 : \mu = \mu_0, \quad H_1 : \mu \neq \mu_0.$$

Plan for two-sided tests for the mean

- Given a random sample X_1, X_2, \dots, X_n from a distribution. Let $\mu = E[X_i]$. Our goal is to decide between

$$H_0 : \mu = \mu_0, \quad H_1 : \mu \neq \mu_0.$$

- If H_0 is true, we expect \bar{X} to be close to μ_0 , and so we expect $W(X_1, X_2, \dots, X_n)$ to be close to 0.

Plan for two-sided tests for the mean

- Given a random sample X_1, X_2, \dots, X_n from a distribution. Let $\mu = E[X_i]$. Our goal is to decide between

$$H_0 : \mu = \mu_0, \quad H_1 : \mu \neq \mu_0.$$

- If H_0 is true, we expect \bar{X} to be close to μ_0 , and so we expect $W(X_1, X_2, \dots, X_n)$ to be close to 0.
- Therefore, we can suggest the following test. Choose a threshold, and call it c .

Plan for two-sided tests for the mean

- Given a random sample X_1, X_2, \dots, X_n from a distribution. Let $\mu = E[X_i]$. Our goal is to decide between

$$H_0 : \mu = \mu_0, \quad H_1 : \mu \neq \mu_0.$$

- If H_0 is true, we expect \bar{X} to be close to μ_0 , and so we expect $W(X_1, X_2, \dots, X_n)$ to be close to 0.
- Therefore, we can suggest the following test. Choose a threshold, and call it c .
- If $|W| \leq c$, accept H_0 , and if $|W| > c$, accept H_1 . How do we choose c ? If α is the required significance level, we must have

$$\begin{aligned} P(\text{type I error}) &= P(\text{Reject } H_0 \mid H_0) \\ &= P(|W| > c \mid H_0) \leq \alpha. \end{aligned}$$

Plan for two-sided tests for the mean

- Given a random sample X_1, X_2, \dots, X_n from a distribution. Let $\mu = E[X_i]$. Our goal is to decide between

$$H_0 : \mu = \mu_0, \quad H_1 : \mu \neq \mu_0.$$

- If H_0 is true, we expect \bar{X} to be close to μ_0 , and so we expect $W(X_1, X_2, \dots, X_n)$ to be close to 0.
- Therefore, we can suggest the following test. Choose a threshold, and call it c .
- If $|W| \leq c$, accept H_0 , and if $|W| > c$, accept H_1 . How do we choose c ? If α is the required significance level, we must have

$$\begin{aligned} P(\text{type I error}) &= P(\text{Reject } H_0 \mid H_0) \\ &= P(|W| > c \mid H_0) \leq \alpha. \end{aligned}$$

- Thus, we can choose c such that $P(|W| > c \mid H_0) = \alpha$.

Example (level α hypothesis test for mean)

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ distribution, where μ is unknown but σ is known. Design a level α test to choose between

$$H_0 = \mu = \mu_0, \quad H_1 : \mu \neq \mu_0.$$

https://www.probabilitycourse.com/chapter8/8_4_3_hypothesis_testing_for_mean.php

Hypothesis Testing is Related to Confidence Interval

Hypothesis Testing is Related to Confidence Interval

- From previous example, the acceptance region for H_0 is

$$\left| \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \right| \leq z_{\frac{\alpha}{2}}.$$

Hypothesis Testing is Related to Confidence Interval

- From previous example, the acceptance region for H_0 is

$$\left| \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \right| \leq z_{\frac{\alpha}{2}}.$$

- Rewriting the above equation as

$$\mu_0 \in \left[\bar{X} - z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}, \bar{X} + z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \right].$$

Hypothesis Testing is Related to Confidence Interval

- From previous example, the acceptance region for H_0 is

$$\left| \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \right| \leq z_{\frac{\alpha}{2}}.$$

- Rewriting the above equation as

$$\mu_0 \in \left[\bar{X} - z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}, \bar{X} + z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \right].$$

- Recall that It is the $(1 - \alpha)100\%$ confidence interval for μ_0 .

Hypothesis Testing is Related to Confidence Interval

- From previous example, the acceptance region for H_0 is

$$\left| \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \right| \leq z_{\frac{\alpha}{2}}.$$

- Rewriting the above equation as

$$\mu_0 \in \left[\bar{X} - z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}, \bar{X} + z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \right].$$

- Recall that It is the $(1 - \alpha)100\%$ confidence interval for μ_0 .
- Relationship between confidence interval problems and hypothesis testing problems.

Example

For the above example, find β , the probability of type II error, as a function of μ .

https://www.probabilitycourse.com/chapter8/8_4_3_hypothesis_testing_for_mean.php

Example

For the above example, find β , the probability of type II error, as a function of μ .

https://www.probabilitycourse.com/chapter8/8_4_3_hypothesis_testing_for_mean.php

Example

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ distribution, where μ and σ are unknown. Design a level α test to choose between

$$H_0 : \mu = \mu_0 \quad H_1 : \mu \neq \mu_0.$$

https://www.probabilitycourse.com/chapter8/8_4_3_hypothesis_testing_for_mean.php#example8_24

Example

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ distribution, where μ and σ are unknown. Design a level α test to choose between

$$H_0 : \mu \leq \mu_0, \quad H_1 : \mu > \mu_0.$$

https://www.probabilitycourse.com/chapter8/8_4_3_hypothesis_testing_for_mean.php#example8_24

- The above can be extended to any other distribution. Let

$$W(X_1, X_2, \dots, X_n) = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}, \quad (2)$$

- If $W \leq c$, accept H_0 , otherwise accept H_1 . To choose c

$$\begin{aligned} P(\text{type I error}) &= P(\text{Reject } H_0 \mid H_0) \\ &= P(W > c \mid \mu \leq \mu_0) \\ &\leq P(W > c \mid \mu = \mu_0). \end{aligned}$$

- Last inequality is because we assume worst-case scenario.

Table of Contents

① Statistical Inference

Interval Estimation and Confidence Level

Hypothesis Testing for the Mean

P-Values

P-Values

- We only reported an "accept" or a "reject" decision as the conclusion of a hypothesis test.

P-Values

- We only reported an "accept" or a "reject" decision as the conclusion of a hypothesis test.
- We can provide more information using what we call P -values.

P-Values

- We only reported an "accept" or a "reject" decision as the conclusion of a hypothesis test.
- We can provide more information using what we call P -values.
- suppose we end up rejecting H_0 at at significance level $\alpha = 0.05$.

P-Values

- We only reported an "accept" or a "reject" decision as the conclusion of a hypothesis test.
- We can provide more information using what we call P -values.
- suppose we end up rejecting H_0 at at significance level $\alpha = 0.05$.
- Then we could ask: "How about if we require significance level $\alpha = 0.01$?" Can we still reject H_0 ?

P-Values

- We only reported an "accept" or a "reject" decision as the conclusion of a hypothesis test.
- We can provide more information using what we call P -values.
- suppose we end up rejecting H_0 at at significance level $\alpha = 0.05$.
- Then we could ask: "How about if we require significance level $\alpha = 0.01$?" Can we still reject H_0 ?

Definition (P -value)

P -value is the lowest significance level α that results in rejecting the null hypothesis.

- If the P -value is small, it means that the observed data is very unlikely to have occurred under H_0 , so we are more confident in rejecting the null hypothesis.

Example

You have a coin and you would like to check whether it is fair or biased.

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$.

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$. Suppose that you need to choose between the following hypotheses:

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$. Suppose that you need to choose between the following hypotheses:

- H_0 (the null hypothesis): The coin is fair, i.e., $\theta = \theta_0 = 1/2$.
- H_1 (the alternative hypothesis): The coin is not fair, i.e., $\theta > 1/2$.

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$. Suppose that you need to choose between the following hypotheses:

- H_0 (the null hypothesis): The coin is fair, i.e., $\theta = \theta_0 = 1/2$.
- H_1 (the alternative hypothesis): The coin is not fair, i.e., $\theta > 1/2$.

We toss the coin 100 times and observe 60 heads.

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$. Suppose that you need to choose between the following hypotheses:

- H_0 (the null hypothesis): The coin is fair, i.e., $\theta = \theta_0 = 1/2$.
- H_1 (the alternative hypothesis): The coin is not fair, i.e., $\theta > 1/2$.

We toss the coin 100 times and observe 60 heads.

- 1 Can we reject H_0 at significance level $\alpha = 0.05$?

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$. Suppose that you need to choose between the following hypotheses:

- H_0 (the null hypothesis): The coin is fair, i.e., $\theta = \theta_0 = 1/2$.
- H_1 (the alternative hypothesis): The coin is not fair, i.e., $\theta > 1/2$.

We toss the coin 100 times and observe 60 heads.

- 1 Can we reject H_0 at significance level $\alpha = 0.05$?
- 2 Can we reject H_0 at significance level $\alpha = 0.01$?

Example

You have a coin and you would like to check whether it is fair or biased. More specifically, let θ be the probability of heads, $\theta = P(H)$. Suppose that you need to choose between the following hypotheses:

- H_0 (the null hypothesis): The coin is fair, i.e., $\theta = \theta_0 = 1/2$.
- H_1 (the alternative hypothesis): The coin is not fair, i.e., $\theta > 1/2$.

We toss the coin 100 times and observe 60 heads.

- 1 Can we reject H_0 at significance level $\alpha = 0.05$?
- 2 Can we reject H_0 at significance level $\alpha = 0.01$?
- 3 What is the P -value?

https://www.probabilitycourse.com/chapter8/8_4_4_p_vals.php

Computing P values

Definition

Consider a hypothesis test for choosing between H_0 and H_1 . Let W be the test statistic, and w_1 be the observed value of W

Computing P values

Definition

Consider a hypothesis test for choosing between H_0 and H_1 . Let W be the test statistic, and w_1 be the observed value of W

- Assume H_0 is true.

Computing P values

Definition

Consider a hypothesis test for choosing between H_0 and H_1 . Let W be the test statistic, and w_1 be the observed value of W

- Assume H_0 is true.
- The P -value is $P(\text{type I error})$ when the test threshold c is chosen to be $c = w_1$.
- For the above example, we can consider

$$W = \frac{X - 50}{5},$$

which is approximately $N(0, 1)$ under H_0 . The observed value of W is

$$w_1 = \frac{60 - 50}{5} = 2.$$

- Thus,

$$\begin{aligned} P - \text{value} &= P(\text{type I error when } c = 2) \\ &= P(W > 2) = 1 - \Phi(2) = 0.023 \end{aligned}$$

Likelihood Ratio Test for Simple Hypotheses

Let $X_1, X_2, X_3, \dots, X_n$ be a random sample from a distribution with a parameter θ . Suppose that we have observed $X_1 = x_1, X_2 = x_2, \dots, X_n = x_n$. To decide between two simple hypotheses

$$H_0 : \theta = \theta_0$$

$$H_1 : \theta = \theta_1$$

We define

$$\lambda(x_1, x_2, \dots, x_n) = \frac{L(x_1, x_2, \dots, x_n; \theta_0)}{L(x_1, x_2, \dots, x_n; \theta_1)}.$$

To perform a likelihood ratio test (LRT), we choose a constant c . We reject H_0 if $\lambda < c$ and accept it if $\lambda \geq c$. The value of c can be chosen based on the desired α .

Solved Example

Example

Here, we look again at the radar problem. More specifically, we observe the random variable X :

$$X = \theta + W,$$

Solved Example

Example

Here, we look again at the radar problem. More specifically, we observe the random variable X :

$$X = \theta + W,$$

where $W \sim N(0, \sigma^2 = \frac{1}{9})$. We need to decide between

Solved Example

Example

Here, we look again at the radar problem. More specifically, we observe the random variable X :

$$X = \theta + W,$$

where $W \sim N(0, \sigma^2 = \frac{1}{9})$. We need to decide between

$$H_0 : \theta = \theta_0 = 0,$$

$$H_1 : \theta = \theta_1 = 1.$$

Solved Example

Example

Here, we look again at the radar problem. More specifically, we observe the random variable X :

$$X = \theta + W,$$

where $W \sim N(0, \sigma^2 = \frac{1}{9})$. We need to decide between

$$H_0 : \theta = \theta_0 = 0,$$

$$H_1 : \theta = \theta_1 = 1.$$

Let $X = x$. Design a level 0.05 test ($\alpha = 0.05$) to decide between H_0 and H_1 .

https://www.probabilitycourse.com/chapter8/8_4_5_likelihood_ratio_tests.php