

## HEART DISEASE IN AYURVEDA III : A HISTORICAL PERSPECTIVE

D. SURESH KUMAR\* & Y.S. PRABHAKAR

### ABSTRACT

Data on all aspects of diagnosis and treatment of heart disease was collected from Charaka Samhita, Sushruta Samhita, Ashtangasangraha, Ashtangahrdaya, Chakradatta, Sharangadhara Samhita and Bhaishajyaratnavali which cover a period of nearly two thousand years. This communication reviews the historical aspects of the subject. Chronology and frequency distribution of the 229 single drugs used in the preparation of the 146 formulations identified from the survey are determined. The results are discussed and special mention is made of the areas of future research.

### Introduction

The ancient Indian medical system of Ayurveda is based on a set of unique concepts. Basing on the six schools of Indian philosophy, Ayurveda maintains that the states of health and disease are influenced by three humours namely vāta, pitta and kapha collectively known as tridōṣa. Because of the wholistic approach to solve problems of health and disease, Ayurveda is able to offer guidelines for the diagnosis and treatment of many diseases for which Western medicine is yet to develop effective medicines devoid of adverse side-effects<sup>1</sup>. As a part of our studies

on the scientific evaluation of ayurvedic medical literature<sup>2-4</sup>, we undertook an extensive search of seven Sanskrit medical texts for cardiological information. Thirteen single drug remedies and 133 multi-component formulations indicated in heart disease were identified<sup>5,6</sup>. This communication treats the subject from the point of view of history of medicine. Attempt has also been made to determine the chronology and frequency-distribution of 229 single drugs used in the preparation of these formulations.

Agniveśa, the author of Caraka

\*Dept. of Physiology and Ethnobiology Unit, International Institute of Ayurveda,  
Ramanathapuram, Coimbatore-641 045.

Saṁhitā flourished in the eighth century BC. The Saṁhitā of Suśruta is also believed to be equally time-honoured. Aṣṭāṅgahṛdaya and Aṣṭāṅgasaṅgraha are attributed to Vāgbhaṭa (600 AD), Cakradatta to Cakrapāṇidatta (1100 AD), Śāringadhara Saṁhitā to Śāringadhara (1300-1400 AD) and Bhaiṣajyaratnāvalī to Govindadāsa (1800 AD)<sup>7-10</sup>. While Caraka Saṁhitā, Suśruta Saṁhitā, Aṣṭāṅgasaṅgraha, and Aṣṭāṅgahṛdaya are treatises on the theory and practice of Ayurveda, Cakradatta and Bhaiṣajyaratnāvalī deal with the treatment of diseases only. Same is the case with Śāringadhara Saṁhitā which is the only authoritative text on ayurvedic pharmacy.

#### Materials and Methods

The seven texts were carefully examined in their entirety for information on all aspects of heart disease. Key words like hṛdroga, hṛdpīḍa, hṛdārti, hṛdgada (heart disease), hṛcchūla, hṛdruk (piercing pain in precordial region), hṛdgraha (seizing pain in precordial region) etc. were used for locating the relevant portions of the texts. After analysis and categorisation of the drug formulae, frequency-distribution of each constituent was determined. Ingredients of all formulations including their variants were subjected to study.

#### Results and Discussion

All the seven authors unanimously agree that heart disease is of five

varieties viz., vātaja (vāta-dominant), pittaja (pitta-dominant), kaphaja (kapha-dominant), sannipātaja (due to combinations of the tridoṣa) and krimija (due to "worms"). However, the presentation of medical information shows interesting patterns in the seven texts. Caraka mainly presents the data on heart, disease in the trimarmīya chapter dealing with the three vital areas of the head, the heart and the pelvis<sup>11</sup>. Suśruta Saṁhitā describes heart disease in the hṛdroga pratiṣedha chapter<sup>12</sup>. Aṣṭāṅgasaṅgraha and Aṣṭāṅgahṛdaya include heart disease in a chapter entitled chardihṛdrogatṛṣṇācikitsā (treatment of emesis, heart disease and thirst)<sup>13,14</sup>. Notwithstanding, Cakradatta and Bhaisajyaratnāvalī contain chapters solely devoted to the treatment of the disease<sup>15, 16</sup>.

#### Causes of heart disease

According to Ayurveda, heart and the alimentary tract are the central points where channels conducting the vital energy (prāṇa) converge. Heart and ten vessels are said to be the conduits for rasa, one of the seven dhātu (tissue elements) which constitute the human body<sup>17</sup>. Vitiation of these channels results in symptoms characteristic of diseases collectively known as hṛdroga (heart disease)<sup>18,19</sup>.

Caraka and Suśruta state that symptoms of heart disease are manifested due to excessive indul-

gence in incompatible food, suppression of natural urges, physical and psychological trauma, fear and anxiety<sup>11, 12</sup>. Caraka further states in the chapter on treatment of rheumatic diseases (*vātavyādhicikitsā*) that vitiation of *vāta* in the alimentary tract, if untreated often leads to disorders of the heart<sup>20</sup>. Vāgbhaṭa on the other hand says that the same factors which cause *gulma* are responsible for causing heart disease<sup>21, 14</sup>. Mādhabavaka (800 AD), the author of the celebrated text *Mādhabanidāna* also gives similar opinion on the causative factors of the disease<sup>25</sup>.

The influence of nutritional factors on the pathogenesis of heart disease is well-studied in Western medicine. It is documented that excessive consumption of fats and proteins eventually leads to obesity, occlusion and hardening of blood vessels and finally to heart disease<sup>26</sup>. Alcohol, a *tikṣṇa* substance according to Ayurveda is proved to cause alcoholic myopathy in addition to its varied effects on the intermediary metabolism<sup>27, 28</sup>.

### Symptoms

Caraka considers paleness of body, fainting, fever, cough, hiccough, dyspepsia, thirst, nausea and anorexia as general symptoms of cardiac disorders. According to him *vātaja* heart disease is characterised by a feeling of emptiness and dryness of

heart, pain in precordial area and fainting<sup>11</sup>. However Suśruta considers intense pain in precordial region to be an ominous symptom of the *vātaja* variety<sup>12</sup>. Vāgbhaṭa also gives emphasis to excruciating pain in precordial area. Nevertheless, he opines that a wise physician should consider intolerance to sound, insomnia, fainting and depressed moods also as typical symptoms<sup>20</sup>.

Burning sensation all over the body, thirst, fainting and the like are considered to be symptoms of pittaja heart disease<sup>11, 12, 20</sup>. However, Caraka and Vāgbhaṭa give importance to fever and discolourisation of body also.

Caraka, Suśruta and Vāgbhaṭa unanimously agree that discomfort in precordial region and increased salivation are the heralding signs of kaphaja heart disease<sup>11, 12, 20</sup>. Interestingly, anorexia, dyspepsia, hypersomnia, heaviness in head, fever, cough etc. are also mentioned.

A combination of symptoms of the three major groups is said to be observed in the *sannipātaja* variety. Though the Sanskrit word *krimi* literally denotes a worm<sup>29</sup>, descriptions available in ayurvedic literature suggest that the word also connotes a subtle pathogen<sup>30, 31</sup>. Thus *krimi hṛdroga* may be the ayurvedic counterpart of infective endocarditis described in Western medicine<sup>32</sup>.

### Conditioning of patients prior to therapy

Pre-conditioning of patients prior to drug treatment was not a subject of much discussion in ayurvedic texts anterior to Cakradatta. Starvation, purgation, induced emesis and smearing of pastes of appropriate drugs are usually recommended by Cakrapāṇi and Govindadāsa. In the case of krimi hṛdroga, the patient is fed with goghṛta and meat for three days followed by purgation<sup>15, 16</sup>.

Govindadasa gives clear instructions on dietetics of heart patients. Grains of *Oryza sativa* L. (with red bran and endosperm), seeds of *Phaseolus mungo* L., fruits of *Mangifera indica* L., *Momordica charantia* L., *Musa paradisiaca* L., *Punica granatum* L., and *Trichosanthes dioica* Roxb., fresh buttermilk and meat of herbivores inhabiting grasslands or forests are some of the prescribed foods. Food possessing qualities similar to those of the humour (s) in question are strictly prohibited<sup>16</sup>.

Ayurvedic practitioners belonging to the traditional school of thought are of the opinion that for achieving excellence in clinical practice a student of the system should study carefully the axioms detailed in the section on fundamental principles (*sūstrasthāna*) of every classical text and delve into their deeper meanings. Caraka advises that even in the

absence of the name of the disease manifested in an individual, detailed information on the humoral imbalance is sufficient to choose the correct line of treatment<sup>33</sup>. A practitioner is therefore expected to extrapolate the axioms in the *sūstrasthāna* and discover their usefulness in diagnosis and therapeutics. This may explain the paucity of information in the four classical texts on pre-treatment and dietetics of heart patients.

### History of medicinal formulations

Details of the 13 single drug remedies and 133 multicomponent formulations identified in the present survey are presented elsewhere<sup>5, 6</sup>. The 146 medicinal formulations involve 203 plants distributed in 69 families, 6 substances of animal origin, 5 inorganic compounds, 7 metals and 8 salts or ashes. As far as the multicomponent formulations are concerned, Caraka Saṃhitā contains 53, Suśruta Saṃhitā 8, Aṣṭāṅgahṛdaya 40, Aṣṭāṅgasaṅgraha 47, Cakradatta 49, Śāṅgadhara Saṃhitā 17 and Bhaiṣajyaratnāvalī 36 formulae. Interestingly enough, Cakradatta is found to contain the maximum number of single drug formulae (9), many of them innovative, followed by Bhaiṣajyaratnāvalī (8). Suśruta Saṃhitā is singular by the absence of any single drug formulae for heart disease.

Frequency distribution of the 229

constituent drugs of the 146 formulations is presented in Table 1, which reveals interesting information. For example, *Zingiber officinale* (91, 68.42%) and *Piper longum* (91, 68.42%) are the two drugs having maximum incidence in recipes. These are followed by *Piper nigrum* (61, 45.86%), *Plumbago zeylanica* (58, 43.60%), *Terminalia chebula* (57, 42.85%), *Emblica officinalis* (49, 36.84%) and *Terminalia bellirica* (35, 26.31%).

*Piper longum*, *Piper nigrum* and *Zingiber officinale* form the famous trikaṭu group of drugs. It is now clear beyond doubt that one of the functions of the trikaṭu drugs is to increase the bioavailability of other drugs<sup>34-36</sup>. This is astonishingly in agreement with Caraka's statement made nearly two thousand years ago that *Piper longum* intensifies the action of drugs to which it is added<sup>37</sup>. Stimulation of digestive processes and dissolution of toxic substances produced due to impaired digestion (āma) are also the other possible modes of action of the trikaṭu drugs. This is more striking especially in the wake of the ayurvedic axiom that pathology of the srotas (minute channels) is the fundamental cause of heart disease<sup>17</sup>. The important position of *Zingiber officinale* in ayurvedic pharmacopoeia is highlighted by the fact that a proverb extolling its medicinal value is still in vogue in Kannada, Malayalam, Tamil and Telugu languages<sup>38</sup>.

*Emblica officinalis*, *Terminalia bellirica* and *Terminalia chebula* make the time-honoured ayurvedic rejuvenator Triphala, a minor variant of which is recommended by Suśruta for heart disease<sup>5</sup>. According to ayurvedic literature the three ingredients have a variety of actions, the most important one being the clearing of srotas<sup>39</sup>. Pharmacological information on these plants indicate that they have choleric, cholagogic, hypolipidemic, anti-necrotic, hypotensive and coronary vasodilatory properties<sup>40-44</sup>. *Plumbago zeylanica* is carminative and antimicrobial<sup>39</sup>. Modern studies show that it normalises intestinal flora<sup>45</sup>.

*Curcuma aromatica* and *Croton tiglium* are two of the minor ingredients which seem to have been accepted from other systems notably the Tamil, Greco-Arabic and Chinese medicines. Similarly many drugs of inorganic and animal origin made their appearance in cardioactive formulations of post-Vāgbhaṭa period obviously due to interaction with Tamil medicine. This is evidenced by Govindadāsa's treatment of the subject. Enrichment of the ayurvedic formulary with the introduction of such drugs during the medieval period points towards the dynamic state of Ayurveda as suggested by Meulenbeld<sup>9</sup>.

Taking into consideration the volume of information on heart

disease, it can be said with certainty that the earlier authors of Ayurveda were familiar with the disease and the formulations useful in its treatment. For example, though described in widely-separated sections like the sūtrasthāna, vimānasthāna (section on specialised groupings of factors), and cikitsāsthāna (section on therapeutics), the Caraka Saṃhitā contains the maximum information on aetiology, nosology and therapeutics of heart disease. However, by the 11 century AD there seems to have emerged the need to deal with the disease in a more organised manner. This explains the highly systematised presentation of cardiological literature by Cakrapāṇidatta and the introduction of *Inula racemosa*, *Moringa oleifera*, *Ricinus communis*, *Terminalia arjuna* etc. as specific single drug remedies for heart disease. Continued wars and political instability which affected medieval Indian society must have necessitated such efforts.

### Conclusions

This survey shows that unanimity exists among the various authors of ayurvedic classics on the aetiology, nosology and therapeutics of heart disease. Careful examination of their opinions shows that obstruction of the srotas (srotorodha) is the fundamental cause of heart disease. The emphasis laid by Ayurveda on the need to regulate digestive physiology sounds rational on the basis of the

argument that all diseases spring from impaired digestion. Interestingly enough, one of the synonyms of the word "disease" is āmaya, which means that which originates from āma<sup>46</sup>. The wide-spectrum nature of the drug formulations identified from this survey is also explainable in this way.

Evidences from Western medical literature<sup>47</sup> suggest that the four major types of heart disease described in ayurvedic literature may be groups of known cardiac ailments and syndromes, krimija hṛdroga being comparable to infective endocarditis. In the backdrop of ayurvedic theory it is also possible to speculate that kaphaja hṛdroga in course of time may get transformed into the vātaja variety. However, well-controlled comparative clinical studies are essential to clarify these points. It is needless to state that in order to do justice to the subject, patients may be examined separately according to the diagnostic procedures of both the systems, while keeping treatment protocol exclusively on ayurvedic lines. It is also essential to pay due attention to the several variables which if overlooked, will certainly increase the "noise" in the clinical data<sup>48</sup>. Western diagnostic parameters can be used for evaluating the success of treatment and comparing the nosological details.

During the course of this survey

it was observed that in addition to the specific formulations prescribed in the sections on therapeutics of heart disease, almost every other chapter of a text contained some useful data or formula which included heart disease among its many indications. For example, Suśruta states in the chapter on the aetiology of pramēha (urinary disorders of polyuric nature) that heart disease can be its secondary affliction<sup>49</sup>. Moreover, in the chapters on the classification of vidradhi (abscess) Suśruta and Vāgbhaṭa describe hṛdayavidradhi<sup>50, 51</sup>. It will be rewarding to ascertain whether the disease has any semblance with the various tumours of the heart described in Western medicine<sup>52</sup>. It is therefore evident that a text needs to be surveyed in its entirety for gathering information on a specific point.

Ayurveda has always remained dynamic in the course of its peregrination through the ages. Accordingly, many new plants introduced into the country through various routes were well-incorporated

into the system without sacrificing its tenets. Bhāvamiśra's monumental work bears testimony to this<sup>53</sup>. This was achieved obviously by understanding the rasa (taste), guṇa (quality), vīrya (potency), vipāka (post-transformational taste) and prabhāva (specific action) of these drugs. Such a feat can be achieved in modern times as well so as to enrich the ayurvedic pharmacopoeia. The technological advances made by Western medicine and biomedical sciences can become handy tools for achieving this goal. This obviates the importance of studies on the utility of the tridoṣa doctrine in the diagnosis and treatment of diseases. Preliminary attempts show that this is not an insurmountable task<sup>54</sup>. Though a wealth of information is available in Ayurveda on heart disease, cardiology is a highly neglected area in contemporary ayurvedic practice. However, the situation can be vastly improved by adopting a multi-disciplinary approach to the problem. It is hoped that this study serves as a harbinger of the much-needed change.

### Acknowledgement

The authors gratefully acknowledge the financial assistance received from the AVR Educational Foundation of Ayurveda, Coimbatore (DSK) and the Ministry of Environment, Forests and Wildlife, Government of India (YSP).

TABLE 1

FREQUENCY-DISTRIBUTION OF THE INGREDIENTS OF THE  
CARDIOACTIVE FORMULATIONS

Ingredient	Sanskrit name	Official part	Incidence in recipes (preference of authors)*
(1)	(2)	(3)	(4)
<b>I Vegetable Drugs</b>			
<i>Acanthaceae</i>			
1) <i>Adhatoda vasica</i> Nees.	Vāsā	root	7 (S1, CS5, A2, C3, Sa1, B2)
2) <i>Andrographis paniculata</i> (Burm. f.) Wall. ex Nees.	Kirātatkta	root	5 (CS1, A5)
3) <i>Barleria prionitis</i> L.	Sairīyā	root	1 (A1)
<i>Amaranthaceae</i>			
4) <i>Achyranthes aspera</i> L.	Apāmārga	root	1 (S1, CS1, A1)
5) <i>Aerva lanata</i> (L.) Juss.	Bhadra	root	1 (A1)
6) <i>Cyathula prostrata</i> (L.) Bl.	Kṣudrāpamārga	root entire	2 (C1, B2)
<i>Anacardiaceae</i>			
7) <i>Buchanania lanzan</i> Spreng.	Priyāla	seed bark	2 (A1, C1)
8) <i>Rhus succedanea</i> L.	Karkaṭaśringi	gall	8 (CS3, A4, C2, B1)
9) <i>Semecarpus anacardium</i> L. F.	Bhallātaka	seed	5 (CS2, A4, C2)
<i>Apocynaceae</i>			
10) <i>Alstonia scholaris</i> (L.) R. Br.	Saptaparṇa	bark	2 (S1, CS1, A1, C1, Sa1)

(1)	(2)	(3)	(4)
11) <i>Holarrhena antidysenterica</i> (Roth.) Wall. ex DC.	Kuṭaja	bark,seed entire	14 (S2, CS4, A5, C8, Sa3, B1)
12) <i>Ichnocarpus frutescens</i> (L.) R. Br.	Kṛṣṇasārīvā	root	2 (S1, CS2, C1, Sa1)
<i>Araceae</i>			
13) <i>Acorus calamus</i> L.	Vacā	rhizome	31 (S3, CS13, A14, C13, Sa4, B13)
14) <i>Alocasia indica</i> Schott.	Mānaka	tuber	1 (B1)
15) <i>Syndapsus officinalis</i> Schott.	Gajapippalī	fruit	15 (S1, CS3, A7, C6 Sa1, B4)
<i>Asclepiadaceae</i>			
16) <i>Calotropis gigantea</i> (L.) R. Br. ex Ait.	Arka	root, latex entire	4 (CS1, A2, C1, B2)
17) <i>Hemidesmus indicus</i> Br.	Sārīvā	root	2 (S1, CS2, C1, Sa1)
18) <i>Holostemma annulare</i> (Roxb.) K. Schum.	Jīvantī	root	9 (CS6, A8, C5, B2)
<i>Berberidaceae</i>			
19) <i>Berberis aristata</i> DC.	Dāruharidrā	root-bark	15 (S1, CS3, A5, C6, Sa3, B2)
<i>Bignoniaceae</i>			
20) <i>Bignonia colais</i> (Buch. Ham. ex Dillw.) Mabberley.	Pāṭalā	root	10 (S1, CS5, A5, C7, Sa1, B2)
21) <i>Oroxylum indicum</i> (L.) Vent.	Śyonāka	root	10 (S1, CS5, A4, C7, Sa1, B2)
<i>Boraginaceae</i>			
22) <i>Heliotropium indicum</i> L.	Hastiśunda	entire	1 (B1)

(1)	(2)	(3)	(4)
23) <i>Trichodesma indicum</i> R. Br.	Avākapuṣpi	root	1 (A1)
<i>Burseraceae</i>			
24) <i>Commiphora mukul</i> (Hook. ex Stocks.) Engl.	Guggulu		gum-resin 3 (A1, C3)
<i>Caesalpiniaceae</i>			
25) <i>Cassia absus</i> L.	Prapouṇḍarīka	seed	1 (CS1, A1)
26) <i>Cassia fistula</i> L.	Āragvadha	bark	1 (S1, CS1, C1, Sa1)
27) <i>Tamarindus indicus</i> L.	Amlikā	fruit, root-bark	9 (S1, CS2, A4. C4, Sa5, B5)
<i>Capparidaceae</i>			
28) <i>Crataev religiosa</i> Forst.	Varuṇa	entire	1 (A1, B1)
<i>Combretaceae</i>			
29) <i>Terminalia arjuna</i> (Roxb.) Wight & Arn.	Arjuna	bark	17 (CS1, C6, B15)
30) <i>Terminalia bellirica</i> (Gaertn.) Roxb.	Vibhītaka	fruit	35 (S1, CS12, A12, C13, Sa8, B8)
31) <i>Terminalia chebula</i> (Gaertn.) Retz.	Harītakī	fruit	57 (S7, CS22, A23, C22, Sa11, B23)
<i>Compositae</i>			
32) <i>Centratherum anthelmin-</i> <i>ticum</i> (Willd.) Kuntze.	Somarājī	seed	1 (B1)
33) <i>Inula racemosa</i> Hook. f.	Puṣkaramūla	root	33 (S3, CS14, A14, C17, Sa5, B16)
34) <i>Saussurea lappa</i> Clark.	Kuṣṭha	root	13 (CS5, A6, C7, Sa1, B5)
35) <i>Sphaeranthus indicus</i> L.	Mahāśrāvaṇī	root	11 (S1, CS4, A5, C7, Sa3, B6)

(1)	(2)	(3)	(4)
36) <i>Vernonia cinerea</i> (L.) Less.	Dāṇḍotpala	root	1 (B1)
<i>Convulvulaceae</i>			
37) <i>Argyreia speciosa</i> Sweet	Vṛddhadāru	root	2 (C2, B1)
38) <i>Ipomoea paniculata</i> R. Br.	Vidārī	root	4 (CS3, A1, C1)
39) <i>Ipomoea pes-tigridis</i> L.	Vyāghranakhi	root	1 (A1)
40) <i>Operculina turpethum</i> (L.) Manso	Tṛiṣṭ	entire	11 (CS4, A5, C4, Sa3, B6)
<i>Cruciferae</i>			
41) <i>Brassica juncea</i> (L.) Czern. & Coss.	Āsuri	seed	1 (C1)
42) <i>Raphanus sativus</i> L.	Mūlaka	root	4 (CS2, A1, Sa1, B1)
<i>Cucurbitaceae</i>			
43) <i>Benincasa cerifera</i> Savi	Kūśmāṇḍa	fruit	3 (C3)
44) <i>Citrullus colocynthis</i> Schrad.	Indravārunī	root	12 (S2, CS6, A4, C4, Sa3, B4)
45) <i>Luffa acutangula</i> Linn. Roxb.	Kōṣātakī	root, entire	2 (A1, B1)
46) <i>Luffa aegyptica</i> Mill.	Dāhmārgava	entire	1 (B1)
47) <i>Solena heterophylla</i> Lour.	Amlavetasa	root	14 (S1, CS4, A4, C6, Sa7, B5)
48) <i>Trichosanthes bracteata</i> (Lamk.) Voight.	Mahākala	root	2 (CS1, A1)
49) <i>Trichosanthes cucumerina</i> L.	Paṭola	leaf, stem, root	14 (S1, CS2, A9, C5, Sa2)
<i>Cyperaceae</i>			
50) <i>Cyperus rotundus</i> L.	Mustā	tuber	31 (CS11, A13, C10, Sa6, B5)
51) <i>Scirpus kynsoor</i> Roxb.	Kāśeruka	tuber	3 (CS3)

(1)	(2)	(3)	(4)
<i>Ebenaceae</i>			
52) <i>Diospyros peregrina</i> (Gaertn.) Gurke	Tinduka	fruit	1 (CS1)
<i>Euphorbiaceae</i>			
53) <i>Baliospermum montanum</i> (Willd.) Muell. & Arg.	Nāgadantī	root	9 (S1, CS3, A4, C3, Sa3, B5)
54) <i>Croton tiglium</i> L.	Jayapāla	seed	1 (B1)
55) <i>Embelica officinalis</i> Gaertn.	Āmalakī	fruit	49 (S4, CS22, A19, C19, Sa8, B10)
56) <i>Euphorbia neriiifolia</i> L.	Sudhā	root, stem, entire	3 (CS1, A1, C1, B2)
57) <i>Euphorbia tirucalli</i> L.	Tṛkantaka	latex	1 (CS1, A1, C1, B1)
58) <i>Homonoia riparia</i> Lour.	Pāṣāṇabheda	root	1 (C1, Sa1, B1)
59) <i>Phyllanthus niruri</i> L.	Tāma'lakī	entire	8 (CS6, A6, C1, B1,)
60) <i>Ricinus communis</i> L. white variety	Eraṇḍla	root	3 (C2, Sa2, B3)
61) <i>Tragia involucrata</i> L.	Vṛścikāli	root	15 (S1, CS5, A4, C6, Sa2, B3)
<i>Ficoidaceae</i>			
62) <i>Mollugo cerviana</i> Ser.	Parpaṭaka	entire	7 (S1, CS2, A5, C1, Sa2)
<i>Gramineae</i>			
63) <i>Cymbopogon schoenanthus</i> (L.) Spreng.	Kattrīṇa	entire	2 (CS2, A1, B1)
64) <i>Desmostachya bipinnata</i> Stapf.	Darbha	root	1 (CS1, A1, C1, B1)
65) <i>Hordeum vulgare</i> L.	Yava	grain	6 (S3, CS3, A4, C2, B2)

(1)	(2)	(3)	(4)
66) <i>Saccharum arundinaceum</i> Retz.	Muñja	root	1 (B1)
67) <i>Triticum aestivum</i> L.	Godhūma	grain	1 (CS1, A1, C1, B1)
68) <i>Vetiveria zizanioides</i> (L.) Nash.	Uśira	root	10 (S1, CS4, A6, C3, Sa3, B1)
<i>Guttiferae</i>			
69) <i>Calophyllum apetalum</i> Willd.	Tejovatī	seed	3 (CS1, A3, C2)
70) <i>Garcinia indica</i> Choisy.	Vṛksāmla	fruit	4 (A4)
71) <i>Mesua ferrea</i> L.	Nāgakesara	flower	13 (CS5, A4, C5, Sa5, B2)
<i>Haemodoraceae</i>			
72) <i>Sansevieria roxburghiana</i> Schult.	Mūrvā	root	6 (S1, CS3, A2, C1, Sa2, B1)
<i>Labiatae</i>			
73) <i>Coleus vettiveroides</i> Jacob	Hrivera	root	5 (CS2, C1, Sa2, B1)
74) <i>Ocimum sanctum</i> L.	Tulasi	root	2 (CS1, A1)
<i>Lauraceae</i>			
75) <i>Cinnamomum camphora</i> (L.) Nees & Eberm.	Karpūra	entire	1 (Sa1)
76) <i>Cinnamomum tamala</i> Nees & Eberm.	Patra	leaf	20 (CS6, A8, C8, Sa5, B2)
77) <i>Cinnamomum verum</i> J.S.Presl.	Tvak	bark	26 (CS10, A13, C10, Sa7, B2)
78) <i>Litsea polyantha</i> Juss.	Meda	bark	11 (CS6, A5, C3, B3)

(1)	(2)	(3)	(4)
<i>Lecythidaceae</i>			
79) <i>Barringtonia acutangula</i> (L.) Gaertn.	Nicula	root, bark	2 (A1, B1)
<i>Liliaceae</i>			
80) <i>Allium sativum</i> L.	Lašuna	bulb	1 (CS1, A2, C1, B1)
81) <i>Asparagus racemosus</i> Willd.	Šatāvarī	root	11 (S1, CS6, A3, C5, Sa2, B2)
82) <i>Fritillaria roylei</i> Hook.	Kṣirakākōli	bulb	7 (CS3, A2, C2, B2,)
83) <i>Lilium polyphyllum</i> D. Don	Kākōli	bulb	8 (CS3, A3, C2, Sa1, B2)
84) <i>Polygonatum cirrhifolium</i> Royle.	Mahāmeda	bulb	7 (CS2, A3, C2, B3)
<i>Lythraceae</i>			
85) <i>Woodfordia fruticosa</i> Kurz.	Dhātakī	flower	3 (CS1, A1, Sa1, B1)
<i>Malvaceae</i>			
86) <i>Sida cordifolia</i> L.	Balā	root	14 (S1, CS7, A8, C6, Sa1, B5)
87) <i>Sida rhombifolia</i> L.	Atibalā	root	5 (CS3, A1, C4, B3)
<i>Marantaceae</i>			
88) <i>Maranta arundinacea</i> L.	Tugākṣīrī	starch	11 (CS3, A2, C3, Sa2, B3)
<i>Meliaceae</i>			
89) <i>Aglaia roxburghiana</i> Miq.	Priyañgu	flower	2 (CS2)
90) <i>Aphanamixis polystachya</i> (Wall.) Parker	Rohitaka	bark	3 (CS1, A1, C1, Sa1, B1)
91) <i>Azadirachta indica</i> A. Juss	Nimba.	bark, seed	14 (S1, CS4, A8, C4, Sa2, B3)

(1)	(2)	(3)	(4)
92) <i>Chukrasia tabularis</i> A. Juss. Aguru		heartwood 1 (A1)	
<i>Menispermaceae</i>			
93) <i>Cissampelos pareira</i> L.	Pāṭhā	root	21 (S3, CS10, A7, C7, Sa3, B4)
94) <i>Tinospora cordifolia</i> (Willd.) Miers. ex Hook. f. & Thomas.	Guḍūcī	stem	13 (S1, CS6, A3, S8, Sa2)
<i>Mimosaceae</i>			
95) <i>Acacia catechu</i> Willd.	Khadira	heartwood 2 (CS1, A1, Sa1)	
96) <i>Acacia concinna</i> DC.	Sātala	entire	2 (CS1, A1, C1,
97) <i>Acacia leucophloea</i> Willd.	Arimeda	bark	1 (C1)
98) <i>Acacia polyantha</i> Willd.	Somavalka	bark	1 (A1)
99) <i>Albizia procera</i> (Roxb.) Benth.	Kaṭabhi	bark	2 (CS1, Sa1)
<i>Moraceae</i>			
100) <i>Ficus benghalensis</i> L.	Vaṭa	bark	1 (CS1, A1)
101) <i>Ficus racemosa</i> L.	Udumbara	bark	1 (CS1, A1)
102) <i>Ficus religiosa</i> L.	Āśvattha	bark	1 (CS1, A1)
<i>Moringaceae</i>			
103) <i>Moringa oleifera</i> Lam.	Śigru	bark, fruit, seed	6 (CS2, A4, C2, Sa2, B1)
<i>Myristicaceae</i>			
104) <i>Myristica fragrans</i> Houtt.	Jātiphala	aril	2 (Sa)
<i>Myrsinaceae</i>			
105) <i>Embelia ribes</i> Burm. f.	Vidāṅga	seed	29 (S1, CS9, A7, C10, Sa7, B9)

(1)	(2)	(3)	(4)
<i>Myrtaceae</i>			
106) <i>Syzygium aromaticum</i> (L.) Merr. & L.M. Perry	Lavaṅga	flowerbud 3 (Sa3)	
<i>Nyctaginaceae</i>			
107) <i>Boerhaavia diffusa</i> L.	Rakta Punarnavā	root, entire 9	(CS4, A3, C2, B3)
<i>Nymphaeaceae</i>			
108) <i>Neelumbo nucifera</i> Gaertn.	Padma	fruit-torus 4	(CS2, A2, A1, C1, Sa1, B1)
109) <i>Nymphaea stellata</i> Willd.	Nilotpala	root-stock 5	(CS5, A1, Sa1, B1)
<i>Onagraceae</i>			
110) <i>Trapa bispinosa</i> Roxb.	Śrīngātaka	fruit	1 (CS1, A1)
<i>Orchidaceae</i>			
111) <i>Habenaria intermedia</i> D. Don.	Rddhi Vṛddhi	entire	8 (CS5, A6, C5, B4)
112) <i>Microstylus muscifera</i> Ridl.	Jivaka	entire	10 (CS5, A5, C2, Sa1, B1)
113) <i>Microstylus wallichii</i> Lindl.	Rśabhaka	entire	7 (CS5, A5, C3, Sa1, B2)
<i>Oxalidaceae</i>			
114) <i>Oxalis corniculata</i> L.	Cāṅgeri	entire	1 (CS1)
<i>Palmae</i>			
115) <i>Areca catechu</i> L.	Kramuka	root, fruit 3	(CS3, A1)
116) <i>Borassus flabellifer</i> L.	Tāla	root	1 (B1)
117) <i>Cocos nucifera</i> L.	Nālikera	seed	1 (C1, B1)
118) <i>Phoenix dactylifera</i> L.	Kharjūra	fruit	2 (CS1, A2)
119) <i>Phoenix pusilla</i> Gaertn.	Sallaki	root	3 (CS2, C1, B1)

(1)	(2)	(3)	(4)
<i>Papilionaceae</i>			
120) <i>Butea monosperma</i> (Lam.) Kuntze	Palāśa	bark, seed, entire	6 (CS4, A4, C2, B2)
121) <i>Cajanus indicus</i> Spreng.	Āḍhakī	seed	1 (B1)
122) <i>Clitoria ternatea</i> L.	Śāṅkhapuṣṭī	root	1 (S1, CS1, A1, C1)
123) <i>Desmodium gangeticum</i> (L.) DC.	Śāliparnī	root	21 (S1, CS10, A10, C10, Sa2, B4)
124) <i>Dolichos biflorus</i> L.	Kulattha	seed	3 (CS1, A1, Sa1, B1)
125) <i>Erythrina variegata</i> L. var. <i>orientalis</i> L.	Mandāra	bark, entire	2 (CS1, B1)
126) <i>Glycyrrhiza glabra</i> L.	Yaṣṭīmadhu	root	18 (S1, CS9, A10, C7, Sa2, B4)
127) <i>Indigofera tinctoria</i> L.	Nīlī	root	1 (B1)
128) <i>Lens culinaris</i> Medic.	Masūra	seed	1 (A1)
129) <i>Mucuna pruriens</i> Hook.	Ātmaguptā	root, seed	6 (S1, CS6, A4, C2, B1)
130) <i>Pongamia pinnata</i> (L.) Pierre	Karañja	bark	2 (A1, C1)
131) <i>Pseuderarthria viscosa</i> (L.) W & A	Pr̄śniparnī	root	16 (S1, CS8, A7, C9 Sa2, B5)
132) <i>Psoralea corylifolia</i> L.	Bākuci	seed	1 (Sa1)
133) <i>Pterocarpus marsupium</i> Roxb.	Asana	heartwood	2 (A2)
134) <i>Pterocarpus santalinus</i> L.	Raktacandana	heartwood	2 (A1, C1)
135) <i>Tephrosia purpurea</i> (L.) Pers.	Śarapuṇkhā	root	1 (B1)
136) <i>Teramnus labialis</i> Spreng.	Māśaparnī	root	1 (CS1)

(1)	(2)	(3)	(4)
<i>Pinaceae</i>			
137) <i>Abies webbiana</i> Lindl.	Tālīsa	leaf	8 (CS2, A3, C7, Sa3, B1)
138) <i>Cedrus deodara</i> Hook. f.	Devadāru	heartwood	21 (CS8, A7, C4, Sa4, B4)
139) <i>Pinus longifolia</i> Roxb. <i>Piperaceae</i>	Sarala	heartwood	1 (CS1, A1)
140) <i>Piper chaba</i> Hunter	Cavya	root	30 (S1, CS10, A7, C10, Sa4, B9)
141) <i>Piper cubeba</i> Linn. f.	Kakkola	fruit	2 (Sa2)
142) <i>Piper longum</i> L.	Pippalī	fruit	91 (S6, CS37 A39, C38, Sa14, B25)
143) <i>Piper nigrum</i> L.	Marica	fruit	61 (S4, CS21, A23, C26, Sa10, B17)
144) <i>Piper wallichii</i> Hand.-Mazz. <i>Plumbaginaceae</i>	Rēnuka	fruit	2 (CS1, B1)
145) <i>Plumbago zeylanica</i> L. <i>Punicaceae</i>	Citraka	root, entire	58 (S3, CS23, A21, C20, Sa9, B20)
146) <i>Punica granatum</i> L. <i>Ranunculaceae</i>	Dāḍima	fruit, seed	18 (S1, CS8, A7, C7, Sa7, B9)
147) <i>Aconitum chasmanthum</i> Stapf. ex Holmes	Vatsanābha	root	1 (A1)
148) <i>Aconitum heterophyllum</i> Wall.	Ativisā	root	12 (S1, CS5, A6, C5, Sa2, B1)
149) <i>Nigella sativa</i> L.	Sthūlajīraka	seed	4 (CS1, A1, Sa2)

(1)	(2)	(3)	(4)
<i>Rhamnaceae</i>			
150) <i>Zizyphus jujuba</i> L.	Badari	fruit, seed	8 (CS4, A4, C2, Sa4, B1)
<i>Rosaceae</i>			
151) <i>Prunus cerasus</i> L.	Elāvāluka	fruit	6 (S1, CS3, A2, C1)
152) <i>Prunus pugettii</i> Roxb.	Padmaka	bark	4 (S1, CS2, A2, C2, Sa2)
<i>Rubiaceae</i>			
153) <i>Gardenia lucida</i> Roxb.	Hīngupatrī	leaf	1 (Sa1)
154) <i>Randia dumetorum</i> Lamk.	Madanaphala	fruit	1 (A1)
155) <i>Rubia cordifolia</i> L.	Mañjiṣṭhā	stem	7 (CS3, A3, C2, Sa1, B1)
156) <i>Spermacoce hispida</i> L.	Vasuka	entire	1 (B1)
<i>Rutaceae</i>			
157) <i>Aegle marmelos</i> (L.) Corr.	Bilva	root, unripe	19 (S1, CS8, A8, C11, Sa2, B5)
		fruit, entire	
158) <i>Citrus decumana</i> L.	Mātulunga	root, fruit	7 (CS1, C4, B3)
159) <i>Citrus medica</i> L. var. <i>Limonum</i>	Bijapūraka	root, fruit	4 (C1, Sa1, B2)
160) <i>Feronia limonia</i> (L.) Swingle	Kapiththa	fruit, bark	3 (S1, CS2, A1)
161) <i>Zanthoxylum alatum</i> Roxb.	Tumburu	seed	8 (A2, C3, Sa1, B2)
<i>Santalaceae</i>			
162) <i>Santalum album</i> L.	Candana	heartwood	8 (S1, CS3, A4, C2, Sa3, B1)

(1)	(2)	(3)	(4)
<i>Sapotaceae</i>			
163) <i>Madhuca longifolia</i> (Koenig) Macbride	Madhūka	flower,	3 (A3, B1)
<i>Scrophulariaceae</i>			
164) <i>Bacopa monniera</i> (L.) Wettst.	Brāhmī	entire	9 (S1, CS4, A6, C2, Sa2, B1)
165) <i>Limnophila gratiissima</i> Bl.	Āmrāgandhīka	entire	1 (CS1)
166) <i>Picrorhiza kurroa</i> Royle ex Benth.	Kāturohīnī	root	19 (S1, CS7, A9, C10, Sa2, B1)
<i>Solanaceae</i>			
167) <i>Hyoscyamus niger</i> L.	Pārasikayavānī	leaf	14 (CS2, A4, C8, Sa5, B7)
168) <i>Solanum ferox</i> L.	Bṛhatī	root,entire	16 (S1, CS6, A6, C10, Sa2, B6)
169) <i>Solanum indicum</i> L.	Bṛhatī	root, fruit	25 (S1, CS11, A10, C12, Sa3, B6)
170) <i>Solanum melongena</i> L.	Vṛntāka	fruit	1 (C1)
171) <i>Solanum nigrum</i> L.	Kākamācī	root	1 (A1, B1)
172) <i>Solanum torvum</i> Swartz.	Vṛntāka	fruit	1 (A1)
173) <i>Solanum violaceum</i> Ortega	Tikta	root	5 (S1, CS2, C4, Sa2)
174) <i>Solanum xanthocarpum</i> Sch. & Wendl.	Vyāghrī	root	4 (A2, C1, Sa1)
175) <i>Withania somnifera</i> Dunal	Aśvagandhā	root	1 (A1, C1, Sa1)
<i>Symplocaceae</i>			
176) <i>Symplocos racemosa</i> Roxb.	Lodhra	bark	4 (S1, CS3, A1)

(1)	(2)	(3)	(4)
<i>Thymelaceae</i>			
177) <i>Aquilaria agallocha</i> Roxb. <i>Ulmaceae</i>	Kṛṣṇāgaru	heartwood 1 (Sa1)	
178) <i>Holoptelia integrifolia</i> (Roxb.) Planch. <i>Umbelliferae</i>	Pūtikarañja	bark, entire 2 (CS1, A1, C1)	
179) <i>Carum carvi</i> L.	Kṛṣṇajīraka	fruit	13 (CS2, A3, C5, Sa4, B8)
180) <i>Coriandrum sativum</i> L.	Kustumburu	seed	19 (S2, CS6, A9, C11, Sa8, B6)
181) <i>Cuminum cyminum</i> L.	Jīraka	fruit	27 (S1, CS8, A9, C14, Sa8, B11)
182) <i>Ferula foetida</i> Regel	Hiṅgu	gum-resin	26 (S3, CS8, A14, C12, Sa5, B12)
183) <i>Heracleum rigens</i> Wall.	Sūksamela	seed	3 (CS3, A2, C1, B2)
184) <i>Peucedanum grande</i> C. B. Clarke	Ajagandhā	root	5 (S1, CS3, A2, C2, Sa2, B2)
185) <i>Peucedanum graveolens</i> L.	Satakusuma	root, entire	9 (CS1, A4, C6, Sa2, B5)
186) <i>Trachyspermum ammi</i> (L.) Sprague <i>Valerianaceae</i>	Ajamodā	fruit	28 (S2, CS8, A8, C13, Sa5, B12)
187) <i>Nardostachys jatamansi</i> DC.	Jāṭāmāṁsi	root	2 (A1, Sa1)
188) <i>Valeriana hardwickii</i> Wall. <i>Verbenaceae</i>	Tagara	root	3 (A1, Sa2)
189) <i>Clerodendrum serratum</i> (L.) Moon	Bhārṅgī	root	16 (S1, CS4, A8, C6, Sa2, B1)

(1)	(2)	(3)	(4)
190) <i>Gmelina arborea</i> L.	Kāśmarī	root, fruit	17 (S1, CS9, A6, C9, Sa1, B5)
191) <i>Lippia nodiflora</i> Mich.	Vanapippalī	root	31 (S2, CS12, A12, C12, Sa6, B2)
192) <i>Premna herbacea</i> Roxb.	Agnimantha	root, entire	11 (S1, CS5, A4, C7, Sa1, B3)
193) <i>Vitex negundo</i> L. Vitaceae	Nirgundi	root	2 (A1, B1)
194) <i>Vitis vinifera</i> L. Zingiberaceae	Drāksā	fruit	15 (CS9, A9, C3, B3)
195) <i>Alpinia galanga</i> (L.) Willd.	Elāparnī	rhizome	17 (CS8, A6, C4, B6)
196) <i>Amomum subulatum</i> Roxb.	Bṛhadēlā	seed	2 (Sa1, B1)
197) <i>Curcuma aromaticā</i> Salisb.	Āmrāgandha-haridrā	rhizome	1 (A1)
198) <i>Curcuma longa</i> Roxb.	Haridrā	rhizome	13 (S2, CS4, A3, C7, Sa2, B2)
199) <i>Elettaria cardamomum</i> (L.) Maton	Elā	seed	27 (S1, CS8, A14, C12, Sa4, B2)
200) <i>Kaempferia galanga</i> L.	Śalādhi	tuber	31 (S2, CS16, A13, C10, Sa6, B19)
201) <i>Kaempferia rotunda</i> L.	Bhūcampaka	tuber	3 (A2, C2, B1)
202) <i>Zingiber officinale</i> Rosc. Zyngophyllaceae	Sunthī	rhizome	91 (S3, CS34, A41, C41, Sa13, B32)
203) <i>Tribulus terrestris</i> L.	Gokṣura	fruit, entire	19 (S1, CS10, A7, C11, Sa2, B7)

(1)	(2)	(3)	(4)
-----	-----	-----	-----

**II Animal Drugs**

204) Conch shell	Śaṅkha	.....	2 (B2)
205) Coral	Pravāla	.....	1 (B1)
206) Deer horn	Mṛgaśringa	.....	3 (C1, Sa1, B1)
207) Oyster shell	Kapardikā	.....	1 (B1)
208) Pearl	Muktā	.....	1 (B1)
209) Pearl oyster shell	Śukti	.....	1 (B1)

**III Inorganic Substances**

210) Chalk of reddish brown colour	Gairika	.....	3 (CS1, A1, C1, Sa1)
211) Iron pyrites	Suvarṇamākṣika	.....	1 (B1)
212) Mica	Abhraka	.....	5 (B5)
213) Mineral bitumen	Śilājatu	.....	2 (CS1, A2, B2)
214) Sulphur	Gandhaka	.....	2 (B2)

**IV Metals**

215) Copper	Tāmra	.....	2 (B2)
216) Gold	Suvarṇa	.....	1 (B1)
217) Iron	Loha	.....	7 (A1, C1, Sa2, B6)
218) Lead	Nāga	.....	1 (B1)
219) Mercury	Rasa	.....	3 (B3)

(1)	(2)	(3)	(4)
220) Silver	Rajata	.....	1 (B1)
221) Tin	Vāṅga	.....	1 (B1)
<b>V Salts and Ashes</b>			
222) Common salt	Sāmudra	.....	9 (CS4, A4, C5, Sa5)
223) Efflorescent salt	Audbhīda	.....	9 (CS4, A4, C3, Sa4)
224) Rock-salt	Saindhava	.....	37 (S4, CS14, A15, C19, Sa8, B12)
225) Salt petre	Saurakṣāra	... .	12 (S1, CS4, A4, C7, Sa6, B3)
226) Alkali from kelp	Sarjikṣāra	.....	4 (S1, C1, B3)
227) Sonchal salt	Sauvarcalā	... .	29 (S2, CS8, A14, C22, Sa6, B12)
228) Vida salt	Viḍalavāṇa	... .	25 (S2, CS7, A11, C11, Sa6, B7)
229) Alkali from barley straw	Yavakṣāra	... .	25 (S2, CS8, A9, C16, Sa7, B14)

\*S = Suśruta Saṁhitā

CS = Caraka Saṁhitā

A = Aṣṭāṅgahṛdaya & Aṣṭāṅgasaṅgraha

C = Cakradatta

Sa = Śāṅgadharma Saṁhitā

B = Bhaiṣajyaratnāvalī

Numerals following the abbreviations indicate incidence in corresponding texts in chronological order.

## BIBLIOGRAPHY

1. Satyavati, G.V. 1982 Some traditional medical systems and practices of global importance. *Ind. J. Med. Res.* 76 (Suppl.), 1-26.
2. Kumar, D.S. and Prabhakar, Y.S. 1987 On the ethnomedical significance of the Arjun tree, *Terminalia arjuna* (Roxb.) Wight & Arn. *J. Ethnopharmacol.* 20, 173-190.
3. Prabhakar, Y.S. and Kumar, D.S. 1988 The chemistry of *Terminalia arjuna* (Roxb.) Wight & Arn. with reference to its medicinal uses. *Plant Méd. Phytothéér.* 22, 30-39.
4. Prabhakar, Y.S. and Kumar, D.S. 1990 The Varuna tree, *Crataeva nurvala* Buch.-Ham., a promising plant in the treatment of urinary stones, *Fitoterapia* (in press).
5. Kumar, D.S. and Prabhakar, Y.S. 1990 A survey of cardioactive drug formulations from Ayurveda I: Single drugs, decoctions, spirituous liquors, powders and tablets. (Communicated to *J. Ethnopharmacol.*)
6. Prabhakar, Y.S. and Kumar, D.S. 1990 A survey of cardioactive drug formulations from Ayurveda II: Porridges, oils, clarified butters, electuaries, pastes, ash preparations and calcined powders. (Communicated to *Fitoterapia*).
7. Sharma, R.K. and Dash, B. 1983 *Caraka Saṁhitā*, Vol. 1, Chowkhamba Sanskrit Studies No. 94, Chowkhamba Sanskrit Series Office, Varanasi, p. xxxviii.
8. Meulenbeld, G.J. 1974 The Mādhavanidānam and its Chief Commentary: Chapters 1-10, E.J. Brill, Leiden, pp. 389-436.
9. Meulenbeld, G.J. 1984 The surveying of Sanskrit medical literature. In: G.J. Meulenbeld (Ed.) *Proceedings of the International Workshop on Priorities in the Study of Indian Medicine*, Rijksuniversiteit te Groningen, Groningen, pp. 31-114.

10. Ray, P. and Gupta, H.N. 1980 *Caraka Saṁhitā* (a scientific synopsis), Indian National Science Academy, New Delhi, p. 1.
11. Vidyalankar, J. 1979 *Caraka Saṁhitā*, Vol. 2, Motilal Banarsidas Delhi, pp. 404-438.
12. Atridev 1981 *Suśruta Saṁhitā*, Motilal Banarsidas, Delhi, pp. 727-728.
13. Upadhyaya, Y. 1975 *Aṣṭāṅgahṛdaya*, Chowkhamba Sanskrit Sansthan, Varanasi, pp. 326-332.
14. Athavale, A.D. 1980 *Aṣṭāṅgasaṅgraha*, Śrimad Atreya Prakashan, Pune, India, pp. 455-460.
15. Mishra, B. 1983 Cakradatta, Haridas Sanskrit Series No. 107, Chowkhamba Sanskrit Series Office, Varanasi, pp. 271-275.
16. Ayurvedacharya, R.S. 1951 *Bhaiṣajyaratnāvalī*, Kashi Sanskrit Granthmaja No. 152, Chowkhamba Sanskrit Pusthakalaya, Varanasi, pp. 485-492.
17. Sharma, R.K. and Dash, B. 1985 *Caraka Saṁhitā*, Vol. 2, Chowkhamba Sanskrit Series Office, Varanasi, pp. 174-176.
18. Sharma, R.K. and Dash, B. 1983 *Caraka Saṁhitā*, Vol. 1, Chowkhamba Sanskrit Series Office, Varanasi, pp. 315-317.
19. Sharma, R.K. and Dash, B. 1983 *Caraka Saṁhitā*, Vol. 1, Chowkhamba Sanskrit Series Office, Varanasi, p. 576
20. Vidyalankar, J. 1979 *Caraka Saṁhitā*, Part 2, Motilal Banarsidas, Delhi, pp. 444-466.
21. Upadhyaya, Y. 1975 *Aṣṭāṅgahṛdaya*, Chowkhamba Sanskrit Sansthan, Varanasi, pp. 234-240.
22. Upadhyaya, Y. 1975 *Aṣṭāṅgahṛdaya*, Chowkhamba Sanskrit Sansthan, Varanasi, pp. 257-263.

23. Athavale, A.D. 1980 *Aṣṭāṅgasaṅgraha*, Srimad Atreya Prakashan, Pune, pp. 363-367.
24. Athavale, A.D. 1980 *Aṣṭāṅgasaṅgraha*, Srimad Atreya Prakashan, Pune, pp. 381-384.
25. Upadhyaya, Y. 1976 *Mādhavanidānam*, Part 1, The Kashi Sanskrit Series No. 158, Chowkhamba Sanskrit Sansthan, Varanasi, p. 495.
26. Swan, H.J.C. and Parmley, W.W. 1974 Congestive heart failure. In : W.A. Sodeman Jr. and W.A. Sodeman (Eds.) *Pathologic Physiology*, 5th eds, W.B. Saunders Co., Toronto, pp. 273-294.
27. Wynne, J. and Braunwald, E. 1984 The cardiomyopathies and myocarditides. In : E. Braunwald (Ed.), *Heart Disease*, Vol. 2, W.B. Saunders Co., Toronto, pp. 1399-1456.
28. Goth, A. 1976 *Medical Pharmacology*, The C.V. Mosby Company, Saint Louis, p. 263.
29. Bahadur, S.R.R. 1961 *Śabdakalpadruma*, Vol. 2, Motilal Banarsidas, Delhi, p. 178.
30. Atridev 1981 *Suśruta Saṃhitā*, Motilal Banarsidas, Delhi, pp. 769-772.
31. Meulenbeld, G.J. 1974 *The Mādhavanidānam and Its Chief Commentary: Chapters 1-10*, E.J. Brill, Leiden, p. 289.
32. Weinstein, L. 1984 Infective endocarditis, In : E. Braunwald (Ed.) *Heart Disease*, Vol. 2, W.B. Saunders Co., Philadelphia, pp. 1136-1182.
33. Sharma, R.K. and Dash, B. 1983 *Caraka Saṃhitā*, Vol. 1, Chowkhamba Sanskrit Series Office, Varanasi pp. 345-346.
34. Atal, C.K., Zutshi, U. and Rao, P.G. 1981 Scientific evidence on the role of ayurvedic herbals on bioavailability of drugs. *J. Ethnopharmacol.* 4, 229-232.

35. Atal, C.K., Dubey, R.K. and Singh, J. 1985 Biochemical basis of enhanced drug bioavailability by piperine: evidence that piperine is a potent inhibitor of drug metabolism. *J. Pharmacol. Exp. Ther.* 232, 258-262.
36. Singh, J., Dubey, R.K. and Atal, C.K. 1986 Piperine mediated inhibition of glucuronidation activity in isolated epithelial cells of the guinea-pig small intestine: evidence that piperine lowers the endogenous UDP-glucuronic acid content. *J. Pharmacol. Exp. Ther.* 236, 488-493.
37. Sharma, R.K. and Dash, B. 1985 *Caṭaka Saṁhitā*, Vol. 2, Chowkhamba Sanskrit Series Office, Varanasi, p. 120.
38. Panikkassery, V. 1982 *Patināyiram Pazhañcollukal* (ten thousand proverbs), National Bookstall, Kottayam Kerala, p. 7.
39. Sharma, P.V. 1956 *Dravyaguṇa Vijñāna*, Vols. 2&3, Chowkhamba Vidya Bhavan, Varanasi, pp. 197-593.
40. Handa, S.S., Sharma, A. and Chakraborti, K.K. 1986 Natural products and plants as liver protecting drugs. *Fitoterapia* 57, 307-351.
41. Giri, J., Devi, T.K.S. and Meerarani, S. 1984 Effect of ginger on serum cholesterol levels. *Ind. J. Nutr. Dietet.* 21, 433-436.
42. Tariq, M., Hussain, S.J., Asif, M. and Jahan, M. 1977 Protective effect of fruit extracts of *Emblica officinalis* (Gaertn.) and *Terminalia belerica* (Roxb.) in experimental myocardial necrosis in rats. *Ind. J. Exp. Biol.* 15, 485-486.
43. Shoji, N., Umeyama, A., Saito, N., Takemoto, T., Kajiwara, A. and Ohizumi, Y. 1986 Dehydropiperonaline, an emide possessing coronary vasodilating activity isolated from *Piper longum*. *J. Pharmac. Sci.* 75, 1188-1189.

44. Sharma, J.N., Rajpal, M.N., Rao, T.S. and Gupta, S.K. 1988 Some pharmacological investigations on the alcoholic extract of Triphala alone and in combination with petroleum ether extract of oleogum resin of *Commiphora mukul*. Indian Drugs 25, 220-223.
45. Iyer, M.A. and Pendse, G.S., 1966 *Plumbago zeylanica L.* (chitraka)- A gastrointestinal flora normaliser Planta Medica 14, 337-351.
46. Apte, V.S. 1964 The Student's English-Sanskrit Dictionary, S. Jain, Delhi, p. 112.
47. Anderson, J.R. 1985 The heart In : J. R. Anderson (Ed.) Muir's Textbook of Pathology, 12th end.. English Language Book Society / Edward Arnold, London, pp. 15. 1-15, 45.
48. Kumar, D.S. 1989 Reflections on some problems faced by contemporary Ayurveda in India. Ayurveda Newsletter (Strasbourg), 9 & 10, 15-24.
49. Atridev 1981 *Suśruta Saṃhitā*, Motilal Banarsidas, Delhi, pp. 248-249.
50. Atridev 1975 *Suśruta Saṃhitā*, Motilal Banarsidas, Delhi, p. 256.
51. Upadhyaya, Y. 1975 *Aṣṭāṅgahṛdaya*, Chowkhamba Sanskrit Sansthan, Varansi, p. 258.
52. Colucci, W.S. and Braunwald, E. 1984 Primary tumours of the heart In : E. Braunwald (Ed.), Heart Disease, Vol. 2, W.B. Saunders Co., Philadelphia, pp. 1457-1469.
53. Pandey, G.S. 1984 *Bhāvaprakāśa Nighaṇṭu*, 7th end., The V. Ayurveda Series No. 28, Chowkhamba Bharati Academy, Varanasi, pp. 1-986.
54. Prabhakar, Y.S. and Kumar, D.S. 1990 A model to quantity disease state based on the ayurvedic concept of tridoṣa. (Communicated to J. Theor. Biol.)

— ३८ —

卷之三

Digitized by srujanika@gmail.com

ପ୍ରକାଶ କାନ୍ତିମାଳା ଏବଂ ପାତ୍ର ହେଉଥିଲା

۱۵۱۱۸