

# MATH 648: Homework #5

April 25, 2024

**Grant Saggars**

- Section 10.3 #1, #2.
- Section 10.4 #2, #3.
- Section 10.6 #3, #4.

## Section 10.3

### Problem 1

Geodesics on a sphere of radius  $R > 0$  correspond to the extremals of the functional

$$J(\phi) = \int_{\theta_0}^{\theta_1} \sqrt{1 + \sin^2 \theta \phi'^2} d\theta,$$

where  $\phi$  is the polar angle,  $\theta$  is the azimuth angle, and  $\phi'$  denotes  $d\phi/d\theta$ . Show that  $J$  satisfies the Legendre condition (10.7).

**Solution:**

Working with the assumption that the Legendre condition, as given, holds in polar coordinates, we find:

$$f_{\phi'} = \frac{\phi' \sin^2(\theta)}{\sqrt{1 + \phi'^2 \sin^2(\theta)}}$$

$$f_{\phi' \phi'} = \frac{\sin^2(\theta)}{(1 + \phi'^2 \sin^2(\theta))^{3/2}} \geq 0$$

(This is never negative for real values, implying convexity)

## Problem 2

Let

$$J(y) = \int_{x_0}^{x_1} \frac{1+y^2}{y'^2} dx.$$

Suppose that  $J$  has a local extremum at  $y$ . Use the Legendre condition to determine the nature of the extremum.

**Solution:**

$$p(x) = f_{y'y'} = \frac{6(1+y^2)}{y'^4}$$

(This is always greater than zero by the even powers, implying a minimal)

## Section 10.4

### Problem 1

Derive the Riccati equation (10.11) associated with the functional of Example 10.3.3. Solve the Riccati equation directly and show that there are no solutions  $w$  defined for all  $x \in [0, \ell]$  if  $\ell > \pi$ .

**Solution:**

#### Part I

Our Riccati equation is:

$$w' + q(x) - \frac{w^2}{p(x)} = 0$$

To get to this, we start with the second variation, defined as:

$$\delta^2 J(\eta, y) = \int_{x_0}^{x_1} (p(x)\eta'^2 + q(x)\eta^2) dx$$

Well behaved functions (consider  $w(x)$ , for example) have  $\eta$  at endpoints which vanish to zero. Jacobi makes the observation that for ANY smooth function  $w$ , one has

$$\int_{x_0}^{x_1} (w\eta^2)' dx$$

by the product rule,  $2w\eta\eta' + w'\eta^2 = (w\eta^2)'$

$$\delta^2 J(\eta, y) = \int_{x_0}^{x_1} (p\eta'^2 + 2w\eta\eta' + (w' + q)\eta^2) dx$$

We know:

1. For  $y$  to be a minimal,  $p(x) \geq 0$  for  $x \in [x_0, x_1]$
2. By the Legendre necessary condition  $p(x)$  will be greater than zero for this domain.

Our integrand can therefore be rewritten as

$$\begin{aligned} p \left( \eta'^2 + 2\frac{w}{p}\eta\eta' + \frac{w^2}{p^2}\eta^2 \right) + \left( w' + q - \frac{w^2}{p} \right) \eta^2 \\ = p \left( \eta' + \frac{w}{p}\eta \right)^2 + \left( w' + q - \frac{w^2}{p} \right) \eta^2 \end{aligned}$$

If we can find a function  $w = w(x)$  so that

$$w' + q(x) - \frac{w^2}{p(x)} = 0, \quad x \in [x_0, x_1]$$

Then the second variation  $\delta^2 J(\eta, y) \geq 0$  for any  $\eta$  since  $p(x) > 0$

**Part II**

To solve we convert the riccati equation into a 2nd order linear system by introducing  $u = u(x)$ :

$$w(x) = -\frac{p(x)u'(x)}{u(x)}, \quad \text{or} \quad u(x) = u_0 e^{-\int_{x_0}^x \frac{w(z)}{p(z)} dz}$$

This gives us our beloved Jacobi Accessory Equation:

$$(p(x)u')' - q(x)u = 0, \quad x \in [x_0, x_1]$$

**Part III**

I am not sure how to prove that there are no solutions for all  $x \in [0, \ell]$  if  $\ell > \pi$ .

## Problem 2

Let

$$f(x, y, y') = y'^2 - y'y + y^2.$$

Show, using elementary arguments, that  $\delta^2 J(\eta, y) \geq 0$  for all  $\eta \in H$ . Derive the Jacobi accessory equation and show by solving this equation that any nontrivial solution  $u$  can have at most one zero.

**Solution:**

We've derived the Jacobi accessory equation in the prior problem. We can also apply the same arguments to show that the second variation is greater than zero for all  $\eta$ :

$$p(x) = f_{y'y'} = 2 \geq 0$$

If we also want to show that there exists no conjugate points for any nontrivial solution, we ought to apply the Jacobi accessory equation.

$$\begin{aligned} (p(x)u(x))' - q(x)u &= 0; \quad q(x) = f_{yy} - \frac{d}{dx}f_{yy'} = 2 \\ (2u)' - 2u &= 0 \end{aligned}$$

This is evidently an exponential function which has only one root.

## Section 10.6

### Problem 3

Let

$$J(y) = \int_0^{\pi/4} (y^2 - y'^2 - 2y \cosh x) dx.$$

Find the extremals for  $J$  and show that for the fixed endpoint problem these extremals produce weak local maxima.

**Solution:**

**E-L:**

$$\begin{aligned} 2y - 2 \cosh x - \frac{d}{dx} 2y' &= 0 \\ y'' - y &= -\cosh x \\ y_c &= c_1 e^x + c_2 e^{-x} \\ y_p &= -\frac{1}{2} x \sinh(x) \\ y(x) &= c_1 e^x + c_2 e^{-x} - \frac{1}{2} x \sinh(x) \end{aligned}$$

**Categorizing Extrema:**

$$\begin{aligned} p(x) &= f_{y'y'} = -2 \\ q(x) &= f_{yy} - \frac{d}{dx} f_{yy'} = 2 \\ (p(x)u')' + q(x)u &= -2u'' + 2u = 0 \\ u(x) &= c_1 e^x + c_2 e^{-x} \end{aligned}$$

Now,  $u(0) = 0$  implies:

$$c_1 + c_2 = 0$$

We choose  $c_1 = -c_2 = \frac{1}{2}$  which gives us:

$$u(x) = \sinh(x) > 0, \quad x > 0$$

Thereby showing it is a local maxima.

## Problem 4

Let

$$J(y) = \int_{x_0}^{x_1} y' (1 + x^2 y') dx,$$

where  $0 < x_0 < x_1$ . Find the extremals for  $J$  and the general solution to the Jacobi accessory equation. Find any conjugate points to  $x_0$  and determine the nature of the extremals for the fixed endpoint problem.

**Solution:**

**E-L:**

$$\begin{aligned} 1 + 2x^2 y' &= c_1 \\ y' &= \frac{dy}{dx} = \frac{c_1 - 1}{2x^2} \\ y &= -\frac{c_1 - 1}{2x} + c_2 \end{aligned}$$

**Categorizing Extremals:**

$$\begin{aligned} p(x) &= f_{y'y'} = x^2 > 0 \\ q(x) &= f_{yy} - \frac{d}{dx} f_{yy'} = 0 \\ x^2 u''(x) &= 0 \end{aligned}$$

We get  $u(x) = c_1 x + c_2$ . For  $u(0) = 0$ ;

$$c_2 = 0$$

There are no other zeroes of this function except for when  $c_1$  is 0, which has undefined behavior as  $u(x) = 0$  for all  $x$ , so this must be a local minimal (meaning no conjugate points).