

# Linear Algebra

—HOFFMAN AND RAY KUNZE

ManJack

2023 年 9 月 18 日

# 前言

这是数学系线性代数的笔记，写给自己。如有错误请见谅，这些只是作为分享。

MaJack

2023 年 9 月 18 日

# 目录

<b>第一章 Linear Equations</b>	<b>1</b>
1.1 环和域 . . . . .	1
1.1.1 加群和环的定义 . . . . .	1
1.1.2 交换元、单位元、零因子、整环 . . . . .	1
1.1.3 除环、域 . . . . .	2
1.2 System of linear Equations . . . . .	3
1.3 Matrix and Elementary Row Operations . . . . .	4

# 第一章 Linear Equations

## 1.1 环和域

### 1.1.1 加群和环的定义

**定义 1.1.1 (加群).** 假如一个 *Abel* 群的代数运算为加法, 并且用符号 ‘+’ 表示, 则该群叫做加群。

**注 1.1.1.** 加群的单位元  $e$  是唯一的, 且  $e=0$ 。称作零元, 我们有以下的计算规则:

$$0 + a = a + 0 = a$$

**定义 1.1.2 (环).** 一个集合  $R$  称之为环满足:

1.  $R$  是一个加群
2.  $R$  对一个乘法来说是一个半群 (半群是一个群胚 + 结合律)
3. 在集合  $R$  上, 乘法对加法满足分配率  $a(b+c)=(ab+ac)$

### 1.1.2 交换元、单位元、零因子、整环

**定义 1.1.3 (交换环).** 一个环叫做一个交换环, 假如:

$$ab = ba$$

在环中乘法运算下的单位元, 叫做环的单位元。

**定义 1.1.4 (单位元).** 环中的单位元  $e$ , 假如对于  $R$  的任意元素  $a$  来说, 有:

$$\boxed{e} a = a \boxed{e} = a$$

↑                      ↑  
单位元

**定义 1.1.5 (零因子).** 一个环中的两个元素  $a, b$  之间如果有一个是  $0$ , 那么  $ab=0$ . 但反之不成立.

$$ab = 0 \not\Rightarrow a = 0 \text{ or } b = 0$$

**例 1.1.1.** 例如模  $n$  的剩余类环: 假设  $n=ab$

若  $n$  不是素数, 假设:

$$[\alpha] \neq [0], [b] \neq [0], [\alpha][b] = [\alpha b] = [n] = [0]$$

则我们可以得知  $ab = 0 \xrightarrow{\text{不成立}} a = 0 \text{ or } b = 0$

**注 1.1.2.** 若是在一个环里,

$$a \neq 0, b \neq 0, ab = 0$$

则  $a$  被称为左零因子,  $b$  被称为右零因子

**定义 1.1.6** (整环). 一个环叫做整环, 满足:

1. 乘法交换律:

$$ab = ba$$

2.  $R$  有单位元  $1$ :

$$1a = a1 = a$$

3.  $R$  没有零因子:

$$ab = 0 \implies a = 0 \text{ or } b = 0$$

**注 1.1.3.**  $a, b$  可以是任意  $R$  中的元.

### 1.1.3 除环、域

**定义 1.1.7** (除环). 一个环  $R$  叫做一个除环, 满足:

1.  $R$  至少包含一个不为  $0$  的元

2.  $R$  有一个单位元

3.  $R$  的每个不等于  $0$  的元有一个逆元

**定义 1.1.8** (域). 一个集合  $F$  被称为域, 如果满足以下条件:

1. 加法封闭性:  $\forall a, b \in F$ , 有  $a + b \in F$ 。

2. 加法可交换性:  $\forall a, b \in F$ , 有  $a + b = b + a$ 。

3. 加法单位元素: 存在加法单位元素  $0$ , 使得  $\forall a \in F$ , 有  $a + 0 = a$ 。

4. 加法逆元素:  $\forall a \in F$ , 存在加法逆元素  $-a$ , 使得  $a + (-a) = 0$ 。

5. 乘法封闭性:  $\forall a, b \in F$ , 有  $a \cdot b \in F$ 。

6. 乘法可交换性:  $\forall a, b \in F$ , 有  $a \cdot b = b \cdot a$ 。

7. 乘法单位元素: 存在乘法单位元素  $1$ , 使得  $\forall a \in F$ , 有  $a \cdot 1 = a$ 。
8. 乘法逆元素:  $\forall a \in F$ , 对于非零元素, 存在乘法逆元素  $a^{-1}$ , 使得  $a \cdot a^{-1} = 1$ 。
9. 分配律:  $\forall a, b, c \in F$ , 满足  $(a + b) \cdot c = a \cdot c + b \cdot c$ 。

**定义 1.1.9** (Subfield). 设  $F$  是一个域。如果  $K \subseteq F$  满足以下条件, 则称  $K$  是  $F$  的子域:

1.  $K$  非空, 并且包含域  $F$  中的加法单位元素  $0$  和乘法单位元素  $1$ 。
2. 对于任意的  $a$  和  $b$  属于  $K$ ,  $a + b$  和  $a \cdot b$  也都属于  $K$  (其中  $+$  和  $\cdot$  分别表示域  $F$  中的加法和乘法运算)。
3. 对于任意的  $a$  属于  $K$ , 它的相反元素  $-a$  也属于  $K$ 。
4. 对于任意的非零元素  $a$  属于  $K$ , 它的乘法逆元素  $a^{-1}$  也属于  $K$ 。

**定义 1.1.10** (Characteristic). *In abstract algebra, "characteristic" is an important concept for a ring or a field. The characteristic is used to describe the smallest positive integer  $n$  for which  $n$  times the multiplicative identity  $1$  equals the additive identity (usually denoted as  $0$ ) in the algebraic structure.*

*For a ring (a set with addition and multiplication operations, satisfying certain algebraic rules), the characteristic refers to the smallest positive integer  $n$  such that  $n$  times  $1$  equals  $0$  (or defined as  $0$  if there is no such  $n$ ).*

*For a field (a special type of ring where every non-zero element has a multiplicative inverse), the characteristic is also a positive integer  $n$  or zero, representing  $n$  times  $1$  equals  $0$  or having characteristic zero if there is no such  $n$ .*

*The significance of the characteristic lies in its impact on the properties and structure of the ring or field. Particularly, in the case of a field, the characteristic is either a prime number or zero. This distinction is useful as it allows us to differentiate between fields of different characteristics and has important applications in properties of algebraic equations and polynomials.*

## 1.2 System of linear Equations

Suppose  $F$  is a field, We consider the problem of finding  $n$  scalars (element of  $F$ )  $x_1, \dots, x_n$  which satisfy the conditions

$$\begin{aligned}
 A_{11} + A_{12}x_1 + \cdots + A_{1n}x_n &= 0 \\
 A_{21} + A_{22}x_1 + \cdots + A_{2n}x_n &= 0 \\
 &\vdots \\
 A_{m1} + A_{m2}x_1 + \cdots + A_{mn}x_n &= 0
 \end{aligned} \tag{1.1}$$

where  $y_1, y_2, \dots, y_n$  and  $A_{ij}, 1 \leq i, j \leq n$  are given elements of  $F$ . We call 1.1 this a **system of  $m$  linear equations** in  $n$  unknowns. Any  $n$ -tuple  $(x_1, x_2, \dots, x_n)$  of elements of  $F$  which satisfies each of the equation in 1.1 is called a solution of the system. If  $y_1 = y_2 = \dots = y_m = 0$ , we say that the system is **homogeneous**, or that each of equations is homogenous,

**定义 1.2.1 (linear combination).** For the 1.1, suppose we select  $m$  scalars  $c_1, \dots, c_m$ , multiply the  $j$ th equation by  $c_j$  and then add.

$$(c_1 A_{11} + \dots + c_m A_{m1})x_1 + \dots + (c_1 A_{m1} + \dots + c_m A_{mn})x_n = c_1 y_1 + \dots + c_m y_m$$

**Note:** Evidently, any solution of the entire system of equations 1.1 will also be a solution of this new equation

**定义 1.2.2 (Linear equivalent).** Let us say that two systems of linear equations are **linearly equivalent** if each equation of one is a linear combination of the equations of the other.

$$\begin{aligned} B_{11} + B_{12}x_1 + \dots + B_{1n}x_n &= z_1 \\ B_{21} + B_{22}x_1 + \dots + B_{2n}x_n &= z_2 \\ &\vdots \\ B_{m1} + B_{m2}x_1 + \dots + B_{mn}x_n &= z_m \end{aligned} \tag{1.2}$$

**定理 1.2.1.** Equivalent system of linear equations have exactly the same solutions.

### 1.3 Matrix and Elementary Row Operations

there is no need to continue writing the 'unknowns'  $x_1, x_2, \dots, x_n$  in the system of linear equations 1.1, since one actually compute only with the coefficient  $A_{ij}$  and the scalars  $y_i$

We shall now abbreviate the system 1.1 by writing:

$$AX = Y$$

where:

$$A = \begin{bmatrix} A_{11} & A_{12} & \cdots & A_{1n} \\ A_{21} & A_{22} & \cdots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{m1} & A_{m2} & \cdots & A_{mn} \end{bmatrix} \tag{1.3}$$

A is the matrix of coefficient of the system

$$X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{and} \quad Y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$



**注 1.3.1.** 1. The entries of the matrix  $A$  are the scalars  $A(i,j) = A_{ij}$

2. The matrix  $A$  is an  $m \times n$  matrix

3. The matrix  $X$  is an  $n \times 1$  matrix

4. The matrix  $Y$  is an  $m \times 1$  matrix

5. The  $AX = Y$  is nothing more than a compact way of writing the system of linear equations 1.3

The elementary row operations on an  $m \times n$  matrix  $A$  over the field  $F$ :

1. multiply of one row of  $A$  by a none-zero scalar  $c$ ;
2. interchange of two rows of  $A$ ;
3. replacement of the  $r$ th row of  $A$  by row  $r$  plus  $c$  time row  $s$ ,  $c$  any scalar and  $r \neq s$ ,

An elementary row operation is thus a special type of function with domain the set of all  $m \times n$  matrices over  $F$  and range the same set. One can describe  $e$  in the three cases as follow:

1.  $e(A)_{ij} = A_{ij}$  if  $i \neq r, e(A)_{rj} = cA_{rj}$
2.  $e(A)_{ij} = A_{ij}$  if  $i \neq r, s, e(A)_{rj} = A_{sj}, e(A)_{sj} = A_{rj}$
3.  $e(A)_{ij} = A_{ij}$  if  $i \neq r, e(A)_{rj} = A_{rj} + cA_{sj}$