

# Bayesian Inference of COVID-19 spreading rates in South Africa - A summary



Jonas Wildberger  
University of Oxford

Problem Description

Model

Results

Discussion

## Problem Description

---

- ▶ Limited data at the beginning of the pandemic made it difficult to grasp the dynamic of the spread
  - ▶ How to account for the high uncertainty in model parameters?

- ▶ Limited data at the beginning of the pandemic made it difficult to grasp the dynamic of the spread
  - ▶ How to account for the high uncertainty in model parameters?
- ▶ How to analyse the efficacy of non-pharmaceutical interventions to minimise the spread of COVID-19?

- ▶ Limited data at the beginning of the pandemic made it difficult to grasp the dynamic of the spread
  - ▶ How to account for the high uncertainty in model parameters?
- ▶ How to analyse the efficacy of non-pharmaceutical interventions to minimise the spread of COVID-19?

Bayesian inference for the rescue!

Model

---




Figure: SEIR model [MM20]






Figure: SEIR model [MM20]

$$\begin{aligned}\frac{dS}{dt} &= -\frac{\lambda SI}{N} \\ \frac{dE}{dt} &= \frac{\lambda SI}{N} - \sigma E \\ \frac{dI}{dt} &= \sigma E - \mu I \\ \frac{dR}{dt} &= \mu I\end{aligned}$$




Figure: SEIR model [MM20]

$$\begin{aligned}\frac{dS}{dt} &= -\frac{\lambda SI}{N} \\ \frac{dE}{dt} &= \frac{\lambda SI}{N} - \sigma E \\ \frac{dI}{dt} &= \sigma E - \mu I \\ \frac{dR}{dt} &= \mu I\end{aligned}$$

- ▶ Basic reproductive number  $R_0 = \frac{\lambda}{\mu}$
- ▶ Delay  $D$  in becoming infected ( $I^{\text{new}}$ ) and being reported
- ▶  $\lambda$  time-varying with change points corresponding to NIPs

Parameter	Prior Distribution
Spreading rate $\lambda_0$	LogNormal(log(0.4),0.5)
Spreading rate $\lambda_1$	LogNormal(log(0.4),0.7)
Spreading rate $\lambda_2$	LogNormal(log(0.4),0.7)
Incubation to infectious rate $\sigma$	LogNormal(log(1/5),0.5)
Recovery rate $\mu$	LogNormal(log(1/8),0.2)
Reporting Delay $D$	LogNormal(log(8),0.2)
Initial Infectious $I_0$	Half-Cauchy(20)
Initial Exposed $E_0$	Half-Cauchy(20)
Change Point $t_1$	Normal(2020/03/18,1)
Change Point $t_2$	Normal(2020/03/28,1)

<https://doi.org/10.1371/journal.pone.0237126.t001>

Parameter	Prior Distribution
Spreading rate $\lambda_0$	LogNormal(log(0.4),0.5)
Spreading rate $\lambda_1$	LogNormal(log(0.4),0.7)
Spreading rate $\lambda_2$	LogNormal(log(0.4),0.7)
Incubation to infectious rate $\sigma$	LogNormal(log(1/5),0.5)
Recovery rate $\mu$	LogNormal(log(1/8),0.2)
Reporting Delay $D$	LogNormal(log(8),0.2)
Initial Infectious $I_0$	Half-Cauchy(20)
Initial Exposed $E_0$	Half-Cauchy(20)
Change Point $t_1$	Normal(2020/03/18,1)
Change Point $t_2$	Normal(2020/03/28,1)

<https://doi.org/10.1371/journal.pone.0237126.t001>

- Likelihood function Student-T distribution

Parameter	Prior Distribution
Spreading rate $\lambda_0$	LogNormal(log(0.4),0.5)
Spreading rate $\lambda_1$	LogNormal(log(0.4),0.7)
Spreading rate $\lambda_2$	LogNormal(log(0.4),0.7)
Incubation to infectious rate $\sigma$	LogNormal(log(1/5),0.5)
Recovery rate $\mu$	LogNormal(log(1/8),0.2)
Reporting Delay $D$	LogNormal(log(8),0.2)
Initial Infectious $I_0$	Half-Cauchy(20)
Initial Exposed $E_0$	Half-Cauchy(20)
Change Point $t_1$	Normal(2020/03/18,1)
Change Point $t_2$	Normal(2020/03/28,1)

<https://doi.org/10.1371/journal.pone.0237126.t001>

- Likelihood function Student-T distribution
- Initial  $R_0 = 3.278$  (CI[2.715, 3.73])

Parameter	Prior Distribution
Spreading rate $\lambda_0$	LogNormal(log(0.4),0.5)
Spreading rate $\lambda_1$	LogNormal(log(0.4),0.7)
Spreading rate $\lambda_2$	LogNormal(log(0.4),0.7)
Incubation to infectious rate $\sigma$	LogNormal(log(1/5),0.5)
Recovery rate $\mu$	LogNormal(log(1/8),0.2)
Reporting Delay $D$	LogNormal(log(8),0.2)
Initial Infectious $I_0$	Half-Cauchy(20)
Initial Exposed $E_0$	Half-Cauchy(20)
Change Point $t_1$	Normal(2020/03/18,1)
Change Point $t_2$	Normal(2020/03/28,1)

<https://doi.org/10.1371/journal.pone.0237126.t001>

- ▶ Likelihood function Student-T distribution
- ▶ Initial  $R_0 = 3.278$  (CI[2.715, 3.73])
- ▶ Hamiltonian Monte Carlo (HMC) method to sample from posterior distribution
  - ▶ No-U-Turn Sampling (NUTS)
  - ▶ 5000 samples with 1000 burn-in

Model	Change Points	LOO	Effective Parameters
SIR	2	448.00	10.27
SEIR	1	457.77	11.60
SEIR	2	459.94	12.00
SIR	1	463.03	8.51
SEIR	0	464.69	16.14
SIR	0	517.72	4.72



<https://doi.org/10.1371/journal.pone.0237126.t002>


Model	Change Points	LOO	Effective Parameters
SIR	2	448.00	10.27
SEIR	1	457.77	11.60
SEIR	2	459.94	12.00
SIR	1	463.03	8.51
SEIR	0	464.69	16.14
SIR	0	517.72	4.72

<https://doi.org/10.1371/journal.pone.0237126.t002>


- SIR model with two change points yields lowest LOO cross entropy loss









- Change points peak on 18/03 (travel ban, social distancing) and 28/03 (mass testing)






- Change points peak on 18/03 (travel ban, social distancing) and 28/03 (mass testing)
- Drop of spreading rate by 80% ( $R_0 = 0.655$  (CI[0.430, 0.960])) and 60% ( $R_0 = 1.304$  (CI[0.887, 1.7748])) respectively



- Change points peak on 18/03 (travel ban, social distancing) and 28/03 (mass testing)
- Drop of spreading rate by 80% ( $R_0 = 0.655$  (CI[0.430, 0.960])) and 60% ( $R_0 = 1.304$  (CI[0.887, 1.7748])) respectively
- Delay  $D \approx 6.848$  including incubation period of 4.537 days, laboratory delay of 2.311 days



- Change points peak on 18/03 (travel ban, social distancing) and 28/03 (mass testing)
- Drop of spreading rate by 80% ( $R_0 = 0.655$  (CI[0.430, 0.960])) and 60% ( $R_0 = 1.304$  (CI[0.887, 1.7748])) respectively
- Delay  $D \approx 6.848$  including incubation period of 4.537 days, laboratory delay of 2.311 days
- Recovery rate  $\mu \approx 0.151$  implies mean infectious period of 6.62 days

# Discussion

---

- ▶ Results confirm the impact of governmental measures to reduce the spread of COVID-19 in South Africa



- ▶ Results confirm the impact of governmental measures to reduce the spread of COVID-19 in South Africa
- ▶ Parameter estimates help quantify the dynamic of the pandemic and account for uncertainty

- ▶ Results confirm the impact of governmental measures to reduce the spread of COVID-19 in South Africa
- ▶ Parameter estimates help quantify the dynamic of the pandemic and account for uncertainty
- ▶ Assumptions on prior distributions?

- ▶ Results confirm the impact of governmental measures to reduce the spread of COVID-19 in South Africa
- ▶ Parameter estimates help quantify the dynamic of the pandemic and account for uncertainty
- ▶ Assumptions on prior distributions?
- ▶ Take into account local, provincial developments



Rendani Mbuyha and Tshilidzi Marwala.  
Bayesian inference of covid-19 spreading rates in south  
africa.  
04 2020.