

Convergence of shearing rate $\omega_{E \times B}$ with boxsize in gradient driven simulation

M. Lippert,^{1, a)} F. Rath,^{1, b)} and A. G. Peeters¹

Physics Department, University of Bayreuth, 95440 Bayreuth, Germany

(*Electronic mail: Florian.Rath1@uni-bayreuth.de)

(Dated: 22 November 2022)

An article usually includes an abstract, a concise summary of the work covered at length in the main body of the article. It is used for secondary publications and for information retrieval purposes.

It is known that radially sheared zonal flows plays a significant role in nonlinear stabilization in tokamak plasmas.¹⁻³. Through advection on the sheared zonal flows the turbulent structure in plasma gets deformed and tilted that causes an $E \times B$ nonlinearity.^{2,4,5} Zonal flows mediate spectral energy transfer to larger radial wave vectors.⁶⁻⁸ The strength of the shearing process is the $E \times B$ shearing rate $\omega_{E \times B}$ which is the radial derivative of the advecting zonal flow velocity.^{4,9} The shearing rate $\omega_{E \times B}$ is defined as

$$\omega_{E \times B} = \frac{1}{2} \frac{\partial^2 \langle \Phi \rangle}{\partial \psi^2} \quad (1)$$

where $\langle \Phi \rangle$ is the zonal electrostatic potential and ψ the radial coordinate that labels the flux surfaces.¹⁰⁻¹² It was shown that the nonlinear threshold for turbulence is directly related to shear stabilization.³ Often the shear stabilization is expressed in the empirical Waltz rule $\omega_{E \times B} \sim \gamma^{9,13}$ where γ is defined as the maximum linear growth rate in the unstable mode. In the

discovered zonal flows, also known as $E \times B$ staircase¹⁴, exhibit amplitudes in terms of the $E \times B$ shearing rate satisfying the stabilization criteria.^{10,15}




FIG. 1: Test




FIG. 4: Test

^{a)}GitHub Repository of this work:
<https://github.com/ManeLippert/Bachelorthesis-Shearingrate-Wavelength>

^{b)}Author to whom correspondence should be addressed:
Florian.Rath1@uni-bayreuth.de




FIG. 2: Test




FIG. 3: Test




FIG. 5: Test




FIG. 6: Test

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

	Counter		Words	
	1 Col	2 Col	1 Col	2 Col
Words			342	
Figure	3	4	200	400
Table	0	0	13	26
Table Row	0	0	5	13
Eq Row	0	0	7	13
Pages			3	
Total			2542	
Remain			958	

¹W. A. Cooper, Plasma Physics and Controlled Fusion **30**, 1805 (1988).

²H. Biglari, P. H. Diamond, and P. W. Terry, Phys. Fluids B: Plasma Physics **2**, 1–4 (1990).

³A. M. Dimits, G. Bateman, M. A. Beer, B. I. Cohen, W. Dorland, G. W. Hammett, C. Kim, J. E. Kinsey, M. Kotschenreuther, A. H. Kritiz, L. L. Lao, J. Mandrekas, W. M. Nevins, S. E. Parker, A. J. Redd, D. E. Shumaker, R. Sydora, and J. Weiland, “Comparisons and physics basis of tokamak transport models and turbulence simulations,” Phys. of Plasmas **7**, 969–983 (2000).

⁴T. S. Hahm and K. H. Burrell, “Flow shear induced fluctuation suppression in finite aspect ratio shaped tokamak plasma,” Phys. Plasmas **2**, 1648–1651 (1995).

⁵K. H. Burrell, Phys. Plasmas **4**, 1499–1518 (1997).

⁶R. E. Waltz and C. Holland, Phys. Plasmas **15**, 122503 (2008).

⁷M. Nakata, T.-H. Watanabe, and H. Sugama, Phys. Plasmas **19**, 022303 (2012).

⁸G. G. Whelan, M. J. Pueschel, and P. W. Terry, Phys. Rev. Lett. **120**, 175002 (2018).

⁹R. E. Waltz, R. L. Dewar, and X. Garbet, Phys. Plasmas **5**, 1784–1792 (1998).

¹⁰F. Rath, A. G. Peeters, R. Buchholz, S. R. Grosshauser, P. Miglano, A. Weikl, and D. Strintzi, “Comparison of gradient and flux driven gyrokinetic turbulent transport,” Phys. Plasmas **23**, 052309 (2016).

¹¹A. Weikl, A. G. Peeters, F. Rath, F. Seiferling, R. Buchholz, S. R. Grosshauser, and D. Strintzi, Phys. of Plasmas **25**, 072305 (2018).

¹²M. J. Pueschel, M. Kammerer, and F. Jenko, Physics of Plasmas **15**, 102310 (2008).

¹³R. E. Waltz, G. D. Kerbel, and J. Milovich, Phys. Plasmas **1**, 2229 (1994).

¹⁴G. Dif-Pradalier, P. H. Diamond, V. Grandgirard, Y. Sarazin, J. Abiteboul, X. Garbet, P. Ghendrih, A. Strugarek, S. Ku, and C. S. Chang, Phys. Rev. E **82**, 025401 (2010).

¹⁵A. G. Peeters, F. Rath, R. Buchholz, Y. Camenen, J. Candy, F. J. Casson, S. R. Grosshauser, W. A. Hornsby, D. Strintzi, and A. Weikl, “Gradient-driven flux-tube simulations of ion temperature gradient turbulence close to the non-linear threshold,” Phys. Plasmas **23**, 082517 (2016).