
Introduction to Feynman-Integrals

Manfred Kraus
mkraus@fisica.unam.mx

Problem 1: Massless Bubble integral

Derive the following solution to the integral in $d = 4 - 2\epsilon$

$$\begin{aligned} \int \frac{d^d \ell}{i\pi^{d/2}} \frac{1}{[\ell^2]^{\nu_1} [(\ell - p)^2]^{\nu_2}} \\ = (-1)^{\nu_1 + \nu_2} (-p^2)^{2 - \epsilon - \nu_1 - \nu_2} \frac{\Gamma(2 - \epsilon - \nu_1) \Gamma(2 - \epsilon - \nu_2) \Gamma(\nu_1 + \nu_2 - 2 + \epsilon)}{\Gamma(\nu_1) \Gamma(\nu_2) \Gamma(4 - 2\epsilon - \nu_1 - \nu_2)}, \end{aligned}$$

1. Combine the two propagators by using *Feynman Parameters*

$$\frac{1}{A^\alpha B^\beta} = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} \int_0^\infty dx_1 dx_2 \frac{\delta(1 - x_1 - x_2) x_1^{\alpha-1} x_2^{\beta-1}}{[x_1 A + x_2 B]^{\alpha+\beta}}$$

and perform the x_2 integration.

2. Complete the square in the denominator, i.e find Q and Δ such that

$$x_1[\ell^2] + (1 - x_1)[(\ell - p)^2] = (\ell - Q)^2 - \Delta.$$

Here Δ is a generalized mass term, i.e it can only depend on x_1, x_2 and p^2 .

3. Shift the loop momentum $\ell \rightarrow \ell + Q$
4. Use the result of the lecture to perform the loop momentum integration

$$\int \frac{d^d \ell}{i\pi^{d/2}} \frac{1}{[\ell^2 - \Delta]^n} = (-1)^n (\Delta)^{d/2 - n} \frac{\Gamma(n - d/2)}{\Gamma(n)}$$

5. Integrate the remaining x_1 integral via Euler's beta function

$$B(x, y) = \int_0^1 dt t^{x-1} (1 - t)^{y-1} = \frac{\Gamma(x) \Gamma(y)}{\Gamma(x + y)}.$$

Problem 2: (**) Two-loop massless sunrise integral

Compute the following integral in $d = 4 - 2\epsilon$

$$\int \frac{d^d \ell_1}{i\pi^{d/2}} \frac{d^d \ell_2}{i\pi^{d/2}} \frac{1}{\ell_1^2 (\ell_1 + \ell_2)^2 (\ell_2 - p)^2} = -(-p^2)^{1-2\epsilon} \frac{\Gamma^3(1 - \epsilon) \Gamma(-1 + 2\epsilon)}{\Gamma(3 - 3\epsilon)}.$$

1. Use the result of Problem 1 to perform the ℓ_1 integration.
2. Use the result of Problem 1 again to perform the remaining ℓ_2 integration.

Problem 3: (5P) Feynman Diagrams