

## **Probabilità**

Marco Isopi

24. Variabili di Poisson

Poisson 1837 1898 Bortkeric morti de celcio di cevello 1) prob. molto bossa del singolo evento 2) numero molto alto di eventi potenziali 3) indipendenza

# incidenti = 
$$X$$
 $E(X) = \frac{1}{2}$ ;

 $X = \sum_{i=1}^{N} Y_i$ 
 $Y(X) = \sum_{i=1}^{N} X_i = \sum_{i=1}^{N} X_i$ 
 $Y(X) = \sum_{i=1}^{N} E(Y_i) = N \cdot \frac{\lambda}{N} = \lambda$ 

$$X \sim \mathbb{E}(N; \frac{\lambda}{N})$$

$$P(X = K) = \binom{N}{K} \left(\frac{\lambda}{N}\right)^{K} \left(1 - \frac{\lambda}{N}\right)^{N-K}$$

$$\lim_{N \to \infty} P_{N}(X = K)$$

$$\binom{N}{K} = \frac{N!}{K!(N-K)!} = \frac{N(N-1) - \cdots (N-K+1)}{K!}$$

$$\frac{N(N-1)-(N-K+1)}{K!} \frac{\lambda^{\kappa}}{N^{\kappa}} \left(1-\frac{\lambda}{N}\right)^{\kappa} \left(1-\frac{\lambda}{N}\right)^{\kappa}$$

$$\frac{\lambda^{\kappa}}{k!} \frac{\left(N(N-1)-(N-K+1)\right)}{N^{\kappa}} \left(1-\frac{\lambda}{N}\right)^{N}$$

$$1 \xrightarrow{N} \frac{\lambda^{\kappa}}{N^{\kappa}} \left(1-\frac{\lambda}{N}\right)^{N}$$

Prisson~ Binomiale Applicazioni: 2 sticurozia hi Il incidenti outomobilistici 

$$X \sim \mathcal{O}(\Lambda) \qquad \lambda \in \mathbb{R}^{+}$$

$$E(X) = \sum_{k=0}^{\infty} X e^{-\lambda} \frac{\lambda^{k}}{k!} = e^{-\lambda} \sum_{k=1}^{\infty} K \frac{\lambda^{k}}{k!} = e^{-\lambda} \sum_{k=1}^{\infty} \frac{\lambda^{k-1}}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^{k-1}}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^{k}}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^{k}}{$$

$$V_{dr}(X) = E(X^{2}) - E(X)^{2}$$

$$E(X^{2}) = \sum_{\kappa=0}^{\infty} K^{2} e^{-\lambda} \frac{\lambda^{\kappa}}{k!} = \lambda^{2} + \lambda$$

$$V_{dr}(X) = \lambda$$
Se  $X \in Li \text{ Poisson} = \sum_{\kappa=0}^{\infty} E(X) = V_{dr}(X)$ 

$$X \sim \mathcal{O}(\lambda)$$
,  $Y \sim \mathcal{O}(\mu)$   
 $X$ ,  $Y$  indipendenti  
 $\longrightarrow X + Y \sim \mathcal{O}(\lambda)$   
 $2 = X + Y$   
 $\{2 = X\} = \bigcup_{j=0}^{N} \{X = j \cap Y = K - j\}$ 

$$P(Z=K) = \sum_{j=0}^{K} P(X=j) P(Y=K-j) =$$

$$= \sum_{j=0}^{K} e^{-\lambda} \frac{\lambda^{j}}{\lambda^{j}} e^{-\lambda} \frac{\lambda^{k-j}}{(K-j)!} =$$

$$e^{-(\lambda+\mu)} \sum_{j=0}^{K} \frac{K!}{j!(K-j)!} \lambda^{j} M^{K-j} =$$

$$e^{-(\lambda+\mu)} \sum_{j=0}^{K} \frac{K!}{j!(K-j)!} \lambda^{j} M^{K-j} =$$

$$e^{-(\lambda+\mu)} \sum_{j=0}^{K} \frac{(\lambda+\mu)^{K}}{j!(K-j)!} \lambda^{j} M^{K-j} =$$

$$X \sim \mathcal{O}(\lambda)$$

$$X = Y_1 + Y_2 + Y_3$$

$$Y_1, Y_2, Y_3 \text{ in dip.} \quad Y_i \sim \mathcal{O}(\frac{\lambda}{3})$$

# eventi che si veni fica no in un intervallo di tempo X= # eventi in a ngior ho e X2(1) => # eventi in une sett. ~ O(71)# eventi in un'ord ~  $O(\frac{1}{24})$ # eventi in 5 orc ~ ( [ ] ( )