Pizza Sales Analysis

1. Retrieve the total number of orders placed.

```
SELECT
COUNT(order_id) AS total_orders
FROM
orders;
```

	total_orders	
•	21350	

2. Calculate the total revenue generated from pizza sales.

```
SELECT
ROUND(SUM(order_details.quantity * pizzas.price), 2) AS total_sales
FROM
order_details
JOIN
pizzas ON pizzas.pizza_id = order_details.pizza_id
```

```
total_sales
> 817860.05
```

3. Identify the highest-priced pizza.

```
SELECT
pizza_types.name, pizzas.price
FROM
pizza_types
JOIN
pizzas ON pizzas.pizza_type_id = pizza_types.pizza_type_id
```

ORDER BY pizzas.price DESC LIMIT 1;

	name	price
•	The Greek Pizza	35.95

4. Identify the most common pizza size ordered.

```
SELECT
pizzas.size,
COUNT(order_details.order_details_id) AS order_count
FROM
pizzas
JOIN
order_details ON pizzas.pizza_id = order_details.pizza_id
GROUP BY pizzas.size
ORDER BY order_count DESC;
```

	size	order_count
•	L	18526
	M	15385
	S	14137
	XL	544
	XXL	28

5. List the top 5 most ordered pizza types along with their quantities.

```
SELECT
pizza_types.name, SUM(order_details.quantity) AS quantity
FROM
pizza_types
JOIN
```

```
pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id

JOIN

order_details ON order_details.pizza_id = pizzas.pizza_id

GROUP BY pizza_types.name

ORDER BY quantity DESC

LIMIT 5;
```

	name	quantity
٠	The Classic Deluxe Pizza	2453
	The Barbecue Chicken Pizza	2432
	The Hawaiian Pizza	2422
	The Pepperoni Pizza	2418
	The Thai Chicken Pizza	2371

6. Join the necessary tables to find the total quantity of each pizza category ordered.

```
SELECT
pizza_types.category,
SUM(order_details.quantity) AS quantity
FROM
pizza_types
JOIN
pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
JOIN
order_details ON pizzas.pizza_id = order_details.pizza_id
GROUP BY pizza_types.category
ORDER BY quantity DESC;
```

	category	quantity
٠	Classic	14888
	Supreme	11987
	Veggie	11649
	Chicken	11050

7. Determine the distribution of orders by hour of the day.

```
SELECT
HOUR(order_time) AS hour, COUNT(order_id) AS order_count
FROM
orders
GROUP BY HOUR(order_time);
```

	hour	order_count
•	11	1231
	12	2520
	13	2455
	14	1472
	15	1468
	16	1920
	17	2336
	18	2399
	19	2009
	20	1642
	21	1198
	22	663
	23	28
	10	8
	9	1

8. Join relevant tables to find the category-wise distribution of pizzas.

```
SELECT
category, COUNT(name)
FROM
pizza_types
GROUP BY category;
```

	category	COUNT(name)
٠	Chicken	6
	Classic	8
	Supreme	9
	Veggie	9

9. Group the orders by date and calculate the average number of pizzas ordered per day

```
SELECT
ROUND(AVG(quantity), 0) as avg_pizza_ordered_per_day
FROM
(SELECT
orders.order_date, SUM(order_details.quantity) AS quantity
FROM
orders
JOIN order_details ON orders.order_id = order_details.order_id
GROUP BY orders.order_date) AS order_quantity;

avg_pizza_ordered_per_day

138
```

10. Determine the top 3 most ordered pizza types based on revenue.

```
SELECT
pizza_types.name,
SUM(order_details.quantity * pizzas.price) AS revenue
FROM
pizza_types
JOIN
pizzas ON pizzas.pizza_type_id = pizza_types.pizza_type_id
JOIN
order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.name
ORDER BY revenue DESC
LIMIT 3;
```

	name	revenue
•	The Thai Chicken Pizza	43434.25
	The Barbecue Chicken Pizza	42768
	The California Chicken Pizza	41409.5

11. Calculate the percentage contribution of each pizza type to total revenue.

```
SELECT
  pizza_types.category,
  ROUND(SUM(order_details.quantity * pizzas.price) / (SELECT
           ROUND(SUM(order_details.quantity * pizzas.price),
                  2) AS total sales
         FROM
           order details
              JOIN
           pizzas ON pizzas.pizza_id = order_details.pizza_id) * 100,
       2) AS revenue
FROM
  pizza_types
    JOIN
  pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
  order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza types.category
ORDER BY revenue DESC;
```

	category	revenue
٠	Classic	26.91
	Supreme	25.46
	Chicken	23.96
	Veggie	23.68

12. Analyze the cumulative revenue generated over time.

```
select order_date,
sum(revenue) over(order by order_date) as cum_revenue
from
(select orders.order_date,
sum(order_details.quantity * pizzas.price) as revenue
from order_details join pizzas
on order_details.pizza_id = pizzas.pizza_id
```

join orders on orders.order_id = order_details.order_id group by orders.order_date) as sales;

	order_date	cum_revenue
•	2015-01-01	2713.85000000
	2015-01-02	5445.75
	2015-01-03	8108.15
	2015-01-04	9863.6
	2015-01-05	11929.55
	2015-01-06	14358.5
	2015-01-07	16560.7

13. Determine the top 3 most ordered pizza types based on revenue for each pizza category.

select name, revenue from (select category, name, revenue, rank() over(partition by category order by revenue desc) as rn from (select pizza_types.category, pizza_types.name, sum((order_details.quantity) * pizzas.price) as revenue from pizza_types join pizzas on pizza_types.pizza_type_id = pizzas.pizza_type_id join order_details on order_details.pizza_id = pizzas.pizza_id group by pizza_types.category, pizza_types.name) as a) as b where rn<=3;

	I	
	name	revenue
١	The Thai Chicken Pizza	43434.25
	The Barbecue Chicken Pizza	42768
	The California Chicken Pizza	41409.5
	The Classic Deluxe Pizza	38180.5
	The Hawaiian Pizza	32273.25
	The Pepperoni Pizza	30161.75
	The Spicy Italian Pizza	34831.25
	The Italian Supreme Pizza	33476.75
	The Sicilian Pizza	30940.5
	The Four Cheese Pizza	32265.70000000065
	The Mexicana Pizza	26780.75
	The Five Cheese Pizza	26066.5