

1. TRIAL SYNOPSIS

1.1 Trial Summary

Sponsor:	PharmaCo International, Inc.
Protocol Number:	ABC-1234-001
Protocol Title:	A Phase III, Randomized, Double-Blind, Placebo-Controlled Study to Evaluate the Efficacy and Safety of Drug ABC in Patients with Moderate to Severe Condition X
Short Title:	Drug ABC Phase III in Condition X
Phase:	Phase III
Indication:	Condition X (Moderate to Severe)
Study Type:	Interventional
Number of Subjects:	450 (planned)
Number of Sites:	60 sites across 12 countries
Planned Duration:	24 months enrollment + 12 months treatment + 4 weeks follow-up
Treatment Duration:	52 weeks
Randomization Ratio:	1:1:1
Blinding:	Double-Blind
Comparator:	Placebo
Primary Endpoint:	Change from baseline in Total Score at Week 52
Key Secondary Endpoint:	Proportion of responders ($\geq 50\%$ improvement) at Week 52
Statistical Method:	MMRM for primary; logistic regression for key secondary
Regulatory Agency:	FDA, EMA

2. STUDY DESIGN

2.1 Trial Arms

Subjects will be randomized in a 1:1:1 ratio to one of three treatment arms:

#	Arm Code	Description
1	ARM A	Drug ABC 10 mg once daily for 52 weeks
2	ARM B	Drug ABC 20 mg once daily for 52 weeks
3	ARM C	Matching placebo once daily for 52 weeks

2.2 Trial Elements and Epochs

The study consists of the following sequential elements within each epoch:

#	Code	Element	Epoch	Duration	Timing
1	SCRN	Screening	SCREENING	Up to 28 days	Day -28 to Day -1
2	LEAD	Lead-in Period	LEAD-IN	14 days	Day -14 to Day -1
3	TRT	Treatment Period	TREATMENT	52 weeks	Day 1 to Week 52
4	TAPER	Dose Taper	TREATMENT	2 weeks	Week 52 to Week 54
5	FU	Follow-up	FOLLOW-UP	4 weeks	Week 54 to Week 58

2.3 Arm-Element Sequence

All arms follow the same element sequence: Screening -> Lead-in -> Treatment -> Taper -> Follow-up
The only difference between arms is the treatment administered during TRT and TAPER elements.

2.4 Study Schema

Screening (28d) --> Lead-in (14d) --> Randomization --> Treatment (52wk) --> Taper (2wk) --> Follow-up (4wk)

3. SCHEDULE OF ASSESSMENTS

3.1 Visit Schedule

The following table summarizes the planned visits, target study days, and visit windows:

Visit#	Visit Name	Target Day	Window (Days)	Epoch
1	Screening	-28	-28 to -1	SCREENING
2	Lead-in	-14	-14 to -1	LEAD-IN
3	Baseline/Randomization	1	1	TREATMENT
4	Week 2	15	12 to 18	TREATMENT
5	Week 4	29	26 to 32	TREATMENT
6	Week 8	57	54 to 60	TREATMENT
7	Week 12	85	82 to 88	TREATMENT
8	Week 16	113	110 to 116	TREATMENT
9	Week 24	169	166 to 172	TREATMENT
10	Week 36	253	250 to 256	TREATMENT
11	Week 52 / EOT	365	362 to 368	TREATMENT
12	Taper Visit 1	372	370 to 375	TREATMENT
13	Taper Visit 2	379	377 to 382	TREATMENT
14	Follow-up	393	390 to 400	FOLLOW-UP
15	Early Termination		Unscheduled	

3.2 Assessments per Visit

- Demographics and Medical History: Screening only
- Vital Signs: All visits except Early Termination
- ECG: Screening, Baseline, Week 12, Week 24, Week 52, Follow-up
- Adverse Events: Continuous from Lead-in through Follow-up
- Tumor Assessments: Baseline, Week 12, Week 24, Week 36, Week 52 (oncology sites only)
- Lab Tests: Screening, Baseline, Week 4, Week 12, Week 24, Week 52, Follow-up
- Concomitant Medications: Continuous from Screening through Follow-up

4. INCLUSION CRITERIA

Subjects must meet ALL of the following criteria to be eligible for the study:

IN01.

Age ≥ 18 years and ≤ 75 years at time of informed consent

IN02.

Documented diagnosis of Condition X for at least 6 months prior to screening

IN03.

Moderate to severe disease activity defined as Total Score ≥ 6 at screening

IN04.

Inadequate response or intolerance to at least one prior standard therapy

IN05.

Body Mass Index (BMI) between 18.0 and 40.0 kg/m² at screening

IN06.

Female subjects of childbearing potential must use highly effective contraception

IN07.

Willing and able to provide written informed consent

IN08.

Able to comply with study procedures and visit schedule

IN09.

Stable doses of permitted concomitant medications for at least 4 weeks prior to baseline

IN10.

Adequate organ function as defined by laboratory values within protocol-specified ranges

5. EXCLUSION CRITERIA

Subjects meeting ANY of the following criteria will be excluded:

EX01.

Known hypersensitivity to Drug ABC or any excipient

EX02.

Current or history of malignancy within the past 5 years (except adequately treated basal cell carcinoma)

EX03.

Active or latent tuberculosis

EX04.

Known HIV, Hepatitis B, or Hepatitis C infection

EX05.

Severe hepatic impairment (Child-Pugh C)

EX06.

Estimated GFR < 30 mL/min/1.73m² at screening

EX07.

Pregnant or breastfeeding women

EX08.

Use of prohibited medications within 5 half-lives or 4 weeks prior to baseline, whichever is longer

EX09.

Participation in another interventional clinical study within 30 days or 5 half-lives prior to screening

EX10.

History of major surgical procedure within 4 weeks prior to screening or planned surgery during the study

EX11.

Any condition that, in the opinion of the investigator, would compromise the safety of the subject or interfere with study assessments

EX12.

QTcF interval > 500 msec at screening ECG