
SKIN & STDs

Macule: A flat, colored lesion, **<2 cm** in diameter, not raised above the surface of the surrounding skin. A "freckle," or ephelid, is a prototype pigmented macule.
Patch: A large (>2 cm) flat lesion with a color different from the surrounding skin. This differs from a macule only in size.
Papule: A small, solid lesion, **<0.5** cm in diameter, raised above the surface of the surrounding skin and, hence, palpable (e.g., a closed comedone, or whitehead, in acne).
Nodule: A larger (0.5–5.0 cm), firm lesion raised above the surface of the surrounding skin. This differs from a papule only in size (e.g., a dermal nevomelanocytic nevus).
Tumor: A solid, raised growth **>5 cm** in diameter.
Plaque: A large (>1 cm), **flat-topped**, **raised lesion**; edges may either be distinct (e.g., in psoriasis) or gradually blend with surrounding skin (e.g., in eczematous dermatitis).
Vesicle: A small, fluid-filled lesion, <0.5 cm in diameter, raised above the plane of surrounding skin. Fluid is often visible, and the lesions are translucent [e.g., vesicles in allergic contact dermatitis caused by *Toxicodendron* (poison ivy)].
Pustule: A vesicle filled with leukocytes. Note: The presence of pustules does not necessarily signify the existence of an infection.
Bulla: A fluid-filled, raised, often translucent lesion **>0.5 cm** in diameter.
Wheal: A raised, erythematous, edematous papule or plaque, usually representing short-lived vasodilatation and vasopermeability.
Telangiectasia: A dilated, superficial blood vessel.



▲ **FIGURE 84-2** Hair cycle and anatomy. The hair follicle cycle consists of stages of rest (telogen), hair growth (anagen), follicle regression (catagen), and hair shedding (exogen). The entire lower epithelial structure is formed during anagen and regresses during catagen. The transient portion of the follicle consists of matrix cells in the bulb that generate seven different cell lineages, three in the hair shaft and four in the inner root sheath. B = bulge; E = epidermis; DP = dermal papilla; H = hair shaft; M = matrix; ORS = outer root sheath; S = sebaceous gland; Sec Grm = secondary germ.

Epithelium	Tissue
Simple squamous epith	Blood vessels,Lymph vessels,mesothelium of body cavities,kidney ascending thin limb
Simple cuboidal epith	Thyroid follicles,respiratory bronchioles,ovaries, rete testes,ependema,kidneys-pct,dct,collecting ducts
Simple columnar epith	Fallopian tubules,cervix,endometrium,ejaculatory duct,bulbourethral glands,stomach,small intestine,large intestine,rectum,gall bladder
Pseudostratified epith	Epididymis,vas deference,seminal vesicle,larynx,trachea,nose,membranous urethra,penile urethra
Keratinized stratified epithelium	Tongue,hardpalate,oesophagus,oropharynx,larynx, ectocervix,vagina,cornea,skin,anus
Transitional stratified epithelium	Renal pelvis,ureter,bladder,prostatic urethra

Disorders of nail	Disease condition responsible
Onycholysis	Hyperthyroidism,psoriasis,fungal inf.
Koilonychia	Iron def. Anemia,syphilis,IHD
Splinter hemorhages	Trauma,nailpsoriasis,subacute bacterial endocarditis
Pterygium of nails	Lichen planus,idiopathic
Half and half nails	Chronic renal failure
Pitting of nails	Psoriasis,alopecia ariata,hand eczema
Beau's lines	Acute systemic illness
Terry's lines	Cirhosis
Hippocratic nails	Clubbing of nails
Mee's lines on nails	Chronic arsenic poisoning,hypoalbunemia
Hyperpigments on nails	Zidovudin,mephalan,daunorubicin,bleomycin
Blue nails	Chloroquine,wilson disease,subungual hematoma
Yellow nail syndrome	Pleural effusion,sinusitis,ankle oedema
Nail folding telengectasia	Dermatomyositis

Hints - Mouth lesions -

Lichen planus

Facial lesions - Pityriasis alba

Elbow,Knee, - Psoriasis

Extremeties - Atopic dermatitis

Trunk - P.Rosea

Umblicated Nodule/Umblication - Molluscum Contagiosum

Herpes Zoster - Thorax

Herpes Simplex - Face

1. Butter fly rash is seen in - SLE
2. Icthyosis is associated with - AIDS,Hypothyroidism,Hodgkin's disease
3. Drug causing Acanthosis negricans - Nicotinic acid
4. Target/Iris lesions r seen in - Erythema multi forme
5. Causative agent of Chancroid - Hemophillus Ducrei
6. Important feature of psoriasis is - Silver scales
7. Most important in diagnosis of leprosy is - slit skin smears for AFB
8. As per WHO minimum duration of treatment for leprosy is - 12 months
9. Slit skin smear is negative in - Neuritic leprosy
10. Come dones are characteristic of - Acne vulgaris
11. Most common cause of pemphigus in India - Pemphigus vulgaris
12. Lesions of childhood atopic derma located in - Cubital & Popliteal fossa
13. Melanocytes No. r reduced except - Albinism(vitiligo,piebaldism,hypopigmentation)
14. Increased incidence of vitiligo is found in - T2DM
15. In a patch of vitiligo - Melanocytes r absent
16. Vitiligo - genetic predisposition/PUVA(best)/Coaltar/Topical steroids
17. Sev itching over webs of fingers,> at night,l/e burrows - Scabies
18. Molluscum contagiosum is caused by - Virus
19. Fungus affecting hair is - Trichophyton/Microsporum
20. Apple jelly nodules are seen in - Lupus vulgaris
21. Scabies oral drug - Ivermectin
22. Keroin is a form of - Fungal infection
23. Most common cause of cellulitis is streptococcus pyogenes
24. Flag sign in Hair is charecteristic of - PEM
25. H.Ducreyi causes - Soft chancre

26. Mycosis fungoides is a - T-cell lymphoma
27. Alopecia areata - spontaneous recovery will occur within 3-6 months in major pts.
28. Cradle cap in babies is - seborrheic dermatitis
29. Leprosy will affect all organs except - Uterus
30. Freis test is diagnostic of - Lymphogranuloma venerium
31. Which is known as Gift spots - Leuconychia
32. Causative organism for common warts - Human papilloma virus
33. Acanthosis nigricans is most common due to carcinoma of - Colon
34. Most specific test for Syphilis - FTA-ABS
35. Photochemotherapy is useful in - Psoriasis
36. Skin condition treated with PUVA therapy is - psoriasis
37. Munro's micro abscess are seen in - Psoriasis
38. Treatment of psoriasis - PUVA/Methotrexate
39. Auspitz sign is found in - Psoriasis
40. Auspitz sign is seen in - plaque type psoriasis
41. Auspitz sign is characteristically seen n - Plaque psoriasis
42. Recurrent painful oro genital ulcers with arthritis is due to - Reiter's syndrome
43. Commonest cutaeneous eruption in SLE is - Erythema of light exposed area
44. Pitting of nails can be seen in - Alopecia areata
45. Young female c/o of genital wart the agent implicated is - Human papilloma virus.
46. Commonest veneral disease in india - Gonorrhoea
47. Tines versicolor is caused by - Malasia Furfur
48. **Helitrope rash is seen on face in - Dermatomyositis**
49. Lesions of **pityriasis rosea** are distributed mostly on - **Trunk**
50. **PUVA therapy is used for - psoriasis/vitiligo/Atopic dermatitis**
51. 5yrs/m with hypopigmented **scaly macule on cheek** some of his classmates also has it probable diagnosis - **Pityriasis Alba**
52. 10yr child- hypopigmented multiple **scaly patch in face** diagnosis - **Pityriasis alba**
53. 3yrchild-eczematous dermatitis on **extensor surface** mother-BA - **Atopic dermatitis**
54. **Mouth lesions** r seen n - **Lichen planus**
55. Nail changes in psoriasis - Oncholysis/pitting/subungual hyperkeratosis
56. Burrow is scabies in the - Stratum corneum
57. Itch is disease is true for - Atopic dermatitis
58. Drug producing erythema nodosum is - Sulphonamides

59. Tzank test is +ve in-Pemphigusfoliacious,P.vegetans,P.vulgaris,senear usher syndrom
60. Tzank cell is a - Keratinocyte
61. Acantholytic bullae are found in - Burns
62. Bulla spread sign is seen in - Pemphigus vulgaris
63. DOC for dermatitis herpatiformis - Dapsone
64. DOC in Gonorrhoea is - Procaine penicillin
65. DOC in Chancroid - Erythromycin
66. Drug of choice in systemic candidiasis is - **Amphotericin-B**
67. DOC in ENL is - Steroids
68. DOC for pustular psoriasis - methotrexate
69. DOC for pustular psoriasis in pregnant lady is - Prednisolone
70. DOC for Herpes Zoster - Acyclovir
71. DOC for Granuloma venerium - Erythromycin
72. DOC for Tinea Unguam - Griseofulvin
73. DOC for systemic candidiasis - Amphotericin B
74. Koebners phenonmenon is seen in - Lichen planus
75. Most common type of scarring alopecia is - Androgenic
76. Patch test is done to document - Delayed type hypersensitivity
77. Mucosal lesion is common in - Lichen planus
78. Recurrent plaques on glans which heal with residual hyperpigmentation is
    -herpes gestationalis
79. Earliest sensation to be lost in Leprosy is - Thermal
80. Degeneration of Basal cells occurs in - Lichen planus
81. Sebaceous cyst r seen in - Gardeners syndrome
82. Abscess is not a primary skin lesion
83. Acne-on drug-presents withblue black pigmentation of nails cause-Tab.Minocycline
84. Most common side effect of retinoids - Skin rashes
85. Acne vulgaris is due to involvement of - Pilosebaceous glands
86. Apocrine gland - modified sweat gland,present in groin&axilla,
87. Rhinophyma (potato nose) is-Glandular form of acne rosacea/sebaceous gland hypertrophy
88. Anagen phase of the hair indicates - phase of activity & growth
89. Female with diffuse alopecia suffered frm typhoid 4 months back diagnosis
    -Telogen effluvium

90. Correct sequence of cell cycle is - G0-G1-S-G2-M
91. Neither raised nor depressed is - Macule
92. Flat discoloration on skin as 1cm is called - Macule
93. Oil drop is seen in - psoriasis of nails
94. Felon is seen in - Nail bed
95. Tinea ungum affects - nail plate
96. Characteristic feature of atopic dermatitis - Pruritis
97. Secondary syphilis manifested by
- painless lymphadenopathy/mucosal erosion/asymptomatic rash
98. Hypopigmented patches r seen in - Naevus anemicus
99. Fir tree type distribution is seen in - pityriasis rosea
100. Air-borne contact dermatitis is diagnosed by - patch test
101. Alopecia areata is treated by - Minoxidil
102. School of fish appearance is shown by - Haemophilus ducreyi
103. Cicatricial alopecia-DLE/Lichen planus/SLE
104. Non cicatrical alopecia - Psoriasis/Alopecia areata
105. Rain drop pigmentation is seen in - chronic Arsenic poisoning
106. Patch test is done for - Allergic contact dermatitis
107. Fluid filled in epidermis and dermis is called - **Vesicle**
108. Commonest site of Atopic dermatitis is - **Ante cubital fossa**
109. Berloque dermatitis is due to contact with: - **Cosmetics**
110. Least common site involvement in psoriasis is - **CNS involvement**
111. Civatte bodies are found in - **Lichen planus**
112. A washer man presents with thickness erosion & discoloration of web spaces of toes diagnosis is - **Candidiasis**
113. Dhobi's itch is - **Tinea cruris**
114. Spongiosis is seen in - Acute Eczema
115. Coin shaped eczema - Nummular Eczema
116. Atopic dermatitis is diagnosed by - Clinical examination
117. Commonest metal causing skin hypersensitivity - Nickel
118. MCC of allergic contact dermatitis in Indian female - Detergents
119. Commonest cause of air borne contact dermatitis in India is - Parthenium grass
120. Pt gets recurrent urticaria while doing exercise & on exposure to sunlight cause is –
-Cholinergic urticaria

121.Annular herald patch is seen in - P.Rosea

122.Nummular ring is caused due to - Idiopathic

123.Darrier's sign is seen in - urticaria pigmentosa

124.Pt with recurrent swelling on face & lips due to emotional stress cause is - CI esterase inhibitor deficiency

125.Quincke's disease is popularly known as - Angioneurotic oedema

126.Bleeding spots seen on removal of scales in psoriasis is called as - Auspitz sign

127.Ash leaf maculae is found in - Tuberous sclerosis

128.D important feature of psoriasis is - Scaling

129.Patch test is a type of - Delayed type hypersensitivity

130.Patch test is read after - 2 days

131.Morbilliform eruptions r seen in - Rubella/Measles

132.Koebner phenonmenon is present in - Psoriasis/Lichen planus/Warts

133.Which is pruritic - Lichen planus/sunburns/Pemphigoid/psoriasis

134.Characteristic feature of lichen planus is - Wickham striae

135.Lichen planus TOC - Systemic steroids

136.Psoralen-A is used in treatment of - Vitiligo

137.Row of Tomb stones appearance is seen in - Pemphigus

138.Commonest variety of Pemphigus - Pemphigus Vulgaris

139.Associated with Pemphigus - Thymoma/CLL/Myasthenia gravis/Atrophic gastritis

140.Blister formation in burn case is in - Subepidermal

141.Mucous lesions r seen n - Pemphigus

142. Intera cellular **IgG** deposition in epidermis s seen n - **Pemphigus**

143.Increased level of **IgE** is seen in - **Atopy**

144.Granular **IgA** deposits at dermal papilla r found n - **Dermatitis herpetiformis**

145.HLA associated with dermatitis herpatiformis - HLA B27

146.Pyoderma gangrenosum is seen in - Ulcerative colitis

147.Erythema nodosum is seen in- RA/PTB/Enteric fever/Leprosy

148.What can pt with gluten sensitive hypersensitivity consume as food -Rice,corn

149.Target or Iris lesion is seen in - Erythema multi forme

150.Causative organism of Molluscum contagiosum - Pox virus

151.Ballooning is characteristic of - Herpes Zoster

152.Acantholysis is characteristic of - Pemphigus vulgaris

153.Pt c/o abd pain i vomitting,psychiatric symt & visual hallucinations

- Intermittant porphyria

154.Porphyria cutanea tarda can be treated by - Phlebotomy

155.Non palpable purpura is seen in - Amyloid/ITP/Drug induced vasculitis

156.Palpable purpura is seen n - HSP/Sr.Sickness/Wegners granulamatosis

157.Treatment for kawasaki's disease - IVIg

158.Lish nodule is seen in - von reclinghausens disease

159.IgA deposits on skin biopsy - Henoch schonlein purpura

160.Triad of Reiters syndrome include - Conjunctivitis/Urethritis/Arthritis

161.Defective DNA repair is associated - Xeroderma pigmentosa

162.Pseudo bubo is seen in - Donovanosis

163.Satellite lesions are seen in - Borderline tuberculoid leprosy

164.Dermatitis & Alopecia are due to defeciency of - Zinc

165.Tuberculoid leprosy is characterised by - Non caseating granuloma in nerve

166.Crocadile skin or sauroderma is seen in - Icthyosis vulgaris

167.Casal's necklace is caused by - pellagra

168.Rash n Measles occurs first n - Post auricular region

169.Erythema marginatum is seen in - Rheumatic fever

170.Commonest nerve involved in Leprosy is - Ulnar

171.Lupus vulgaris - Skin & Mucosa TB/Apple jelly nodules at root of nose

172.Acrodermatitis enteropathica - decreased Zinc level/autosomal recessive

173.Rose spot are seen in - Typhoid fever

174.Slapped cheek appearance is seen in - Erythema infectiosum

175.Phrynoderma - deficiency of - vitamin A

176.Virchows cells are seen in - Leprosy

177.Lepromin test is used for - prognosis

178.Features of Lepromatous leprosy - Saddle nose/Gynaecomastia/Mydriasis

179.Commonest fungal infection in female genitalia in diabetes is - Candida

180.Most potent anti leprotic drug is - Rifampicin

181.Permethrin is used in treatment of - Scabies

182.Leprosy do not involve - CNS

183.Leprosy affects all organs except - Uterus

184.In leprosy nerves commonly involved are - High ulnar,low median

185.Skin hazards of swimming are - Verrucae/M.marinum infection

186.Most common type of leprosy in India - TT

187.Commonest skin infection in children is - Impetigo contagiosa

188.Characteristic lesions of scabies is - Burrow

189.Honey colour crusts are characteristics of - Impetigo

190.Most common organism causing Tinea capitis - Microsporum

191.Athelet's foot - 4th toe web is involved/hyperthyroidism(+)/caused by Trichophyton

192.Tinea incognito is seen with - Steroid treatment

193.Treatment of Acne - 13 cis-retionol/Minocycline/Erythromycin

194.Multi drug therapy is given for - Leprosy

195.Tinea versicolor is caused by - Malassezia Furfur

196.Griseofulvin is useful in - T.capitis,T.cruris,T.Pedis

197.Treatment for T.Versicolor - clotrimazole/sod thiosulphate/miconazole/selenium sulphide

198.Safety pin appearance is shown by - Donovanosis granulomatis

199.Black dot ring worm is caused by - Trichophyton

200.Washerman i thickness erosion & discolouration of web spaces & toes -Candidiasis

201.Medically most important form of UV - UV-B

202.Rangeof light which causes max damage to skin - 290-360nm

203.Griseofulvin is not used in - T.Versicolor & Candida

204.Dhobi's Itch is - T.Cruris

205.Incubation period of scabies - 4 wks

206.Antidote for T.warfarrin - vit-k

207.Circlre of Hebra is associated with - Scabies

208.Vagabond's disease is - Pediculosis corporis

209.Impetigo contagiosa most commonly due to - Staphylococcus

210.Condylomata acuminata is caused by - HPV

211.LGV is caused by - Chlamydia trochamatis

212.Genital elephantiasis is caused by - Lymphogranuloma venerum

213.Sign of groove is found in - LGV

214.Frie's test is done in - LGV

215.DOC for LGV - doxycycline

216.DOC for Herpes Zoster - Acyclovir

217.DOC in primary syphilis is - Benzathine penicilline

218.Pseudo-isomorphic phenonmenon is shown by - Warts

219.Most severe form of scabies - Norwegian scabies

220.Chancroid is caused by - Hemophilus ducreyi

221.Most frequent cause of recurrent genital ulceration in sexually active male is
- Herpes genitalis

222.Ivermectin is indicated in treatment of - Scabies

223.Sabre tibia is seen in - syphilis

224.Acanthosis nigricans is indicative of
- internal malig/endocrine dis/Blooms synd/pigmentation of face & neck

225.Langerhans cells in skin are - Antigen presenting cells

226.Shagreen patch may be seen in - Tuberous sclerosis

227.Rodent ulcer is a type of - basal cell ca

228.Hypervitaminosis A is associated i - hair loss/dry skin/hypopigmentation/# bones.

229.Blue line on Gums r seen in - Acute monocytic leukemia

230.Escharotomy is done for - circumferential scars.

231.Leprosy does not affect - Uterus & ovary

232.Erythema multiforme is caused by - Virus

233.Systemic treat ment of Scabies - Ivermectin

234.Wasserman test is done for - syphilis

235.Protective from ultra violet rays - Melanocytes

236.Child born i nasal discharge & rash all over the body - Congenital syphilis

237.25 yr GM i urticaria,recurrent skin infection,abd cramps & diarhea upon taking sea foods
- Atopic dermatitis

238.Male has painless ulcer 9 days after coitus i professional sex worker diag - Chancre

239.Multi bacillary leprosy is -12 months

240.Pauci bacillary leprosy is - 6 months

241.Silvery scaling is an important feature of - Psoriasis

242.Melasma is seen in - Face

243.Iron defeciency anemia - Koilonychias

244.Apple jelly nodules- lupus vulgaris

245.Butterfly rashes r seen n - SLE

246.Auspitz sign is seen n - Psoriasis

247.Row of Tomb stone appearance is seen in - Pemphigus

248.Kerion is a - Fungal infection

249.Anagen is the growth phase of - Hair

250.Ashleaf spots are seen in - Tuberous sclerosis