



Probability and Stochastic Processes

Multiple Random Variables, Joint CDF and its Properties, Jointly
Discrete Random Variables, Joint PMF, Conditional PMF, Jointly
Continuous Random Variables, Joint PDF

Karthik P. N.

Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

02/05 September 2024

Multiple Random Variables


Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Two Random Variables)

Given two \mathcal{F} -measurable random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$, we say $(X, Y) : \Omega \rightarrow \mathbb{R}^2$ is a **random variable** with respect to \mathcal{F} if

$$(X, Y)^{-1}(B) = \{\omega \in \Omega : (X(\omega), Y(\omega)) \in B\} \in \mathcal{F} \quad \forall B \in \mathcal{B}(\mathbb{R}^2).$$



$$\forall B \in \mathcal{B}(\mathbb{R}^2), \quad (X, Y)^{-1}(B) \in \mathcal{F}$$

Joint Probability Law of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Joint Probability Law of Two Random Variables)


Given two random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined with respect to \mathcal{F} , their **joint probability law** $\mathbb{P}_{X,Y} : \mathcal{B}(\mathbb{R}^2) \rightarrow [0, 1]$, is the probability measure defined as

$$\mathbb{P}_{X,Y}(B) = \mathbb{P}(\{\omega \in \Omega : (X(\omega), Y(\omega)) \in B\}), \quad B \in \mathcal{B}(\mathbb{R}^2).$$

Remarks:

- $\mathbb{P}_{X,Y}$ is called the **pushforward** of \mathbb{P} under the random variable (X, Y)
- $\mathbb{P}_{X,Y}$ is the **probability law** of the random variable (X, Y)
- $\mathbb{P}_{X,Y}$ gives the **full probabilistic description** of (X, Y)

The Picture to Have in Mind



$$\mathbb{P}_{X,Y}(B) = \mathbb{P}((X, Y)^{-1}(B)) \quad \forall B \in \mathcal{B}(\mathbb{R}^2)$$


Remarks

- A special class of sets in $\mathcal{B}(\mathbb{R}^2)$ are semi-infinite rectangles of the form

$$(-\infty, x] \times (-\infty, y], \quad x, y \in \mathbb{R}.$$

- $\mathcal{B}(\mathbb{R}^2) = \sigma(\{(-\infty, x] \times (-\infty, y] : x, y \in \mathbb{R}\})$

Joint CDF of Two Random Variables



$$F_{X,Y}(x, y) = \mathbb{P}_{X,Y}((-\infty, x] \times (-\infty, y]) = \mathbb{P}(\{X \leq x\} \cap \{Y \leq y\}), \quad x, y \in \mathbb{R}$$

Definition (Joint CDF)

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Given random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ with respect to \mathcal{F} , their **joint CDF** $F_{X,Y} : \mathbb{R}^2 \rightarrow [0, 1]$ is defined as

$$F_{X,Y}(\mathbf{x}, \mathbf{y}) = \mathbb{P}_{X,Y}((-\infty, \mathbf{x}] \times (-\infty, \mathbf{y}]) = \mathbb{P}(\{X \leq \mathbf{x}\} \cap \{Y \leq \mathbf{y}\}), \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}.$$

Notation

- $\{X \leq x\} \cap \{Y \leq y\} = \{X \leq x, Y \leq y\}$
- $\mathbb{P}(\{X \leq x\} \cap \{Y \leq y\}) = \mathbb{P}(X \leq x, Y \leq y)$

Joint CDF \longleftrightarrow Joint Probability Law

- If we know $\mathbb{P}_{X,Y} = \{\mathbb{P}_{X,Y}(B) : B \in \mathcal{B}(\mathbb{R}^2)\}$, then we can extract the CDF $F_{X,Y} : \mathbb{R}^2 \rightarrow [0, 1]$ by using the formula

$$F_{X,Y}(x, y) = \mathbb{P}_{X,Y}((-\infty, x] \times (-\infty, y]), \quad x, y \in \mathbb{R}.$$

- Given the joint CDF $F_{X,Y} : \mathbb{R}^2 \rightarrow [0, 1]$, let

$$\mathbb{P}_{X,Y}((-\infty, x] \times (-\infty, y]) = F_{X,Y}(x, y), \quad x, y \in \mathbb{R}.$$

Then, by **Caratheodory's extension theorem**, there exists a unique extension of $\mathbb{P}_{X,Y}$ to all Borel subsets of \mathbb{R}^2

Properties of Joint CDF

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$

Let $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ be random variables with respect to \mathcal{F} with joint CDF $F_{X,Y}$

- $\lim_{x,y \rightarrow -\infty} F_{X,Y}(x,y) = 0, \quad \lim_{x,y \rightarrow +\infty} F_{X,Y}(x,y) = 1$
- (**Monotonicity**) If $x_1 \leq x_2$ and $y_1 \leq y_2$, then $F_{X,Y}(x_1, y_1) \leq F_{X,Y}(x_2, y_2)$
- $F_{X,Y}$ is **continuous from the right and top**, i.e., for all $x, y \in \mathbb{R}$,
$$\lim_{u \downarrow 0, v \downarrow 0} F_{X,Y}(x+u, y+v) = F_{X,Y}(x, y).$$
- $\lim_{y \rightarrow \infty} F_{X,Y}(x, y) = F_X(x)$ for all $x \in \mathbb{R}$
 $\lim_{x \rightarrow \infty} F_{X,Y}(x, y) = F_Y(y)$ for all $y \in \mathbb{R}$

Marginal Law/CDF from Joint Law/CDF

Given joint CDF/law, we may extract the marginal CDFs/laws

The converse is not possible in general

CDF	Law
$F_{X,Y} = \{F_{X,Y}(x, y) : x, y \in \mathbb{R}\}$	$\mathbb{P}_{X,Y} = \{\mathbb{P}_{X,Y}(B) : B \in \mathcal{B}(\mathbb{R}^2)\}$
$F_X(x) = \lim_{y \rightarrow +\infty} F_{X,Y}(x, y)$	$\mathbb{P}_X(A) = \mathbb{P}_{X,Y}(A \times \mathbb{R})$
$F_Y(y) = \lim_{x \rightarrow +\infty} F_{X,Y}(x, y)$	$\mathbb{P}_Y(A) = \mathbb{P}_{X,Y}(\mathbb{R} \times A)$

Table: Marginal law/CDF from joint law/CDF.


Multiple Random Variables

Fix a measurable space (Ω, \mathcal{F}) .

Definition (Multiple Random Variables)

Given random variables X_1, \dots, X_n defined with respect to \mathcal{F} , we say $(X_1, \dots, X_n) : \Omega \rightarrow \mathbb{R}^n$ is a **random variable** with respect to \mathcal{F} if

$$(X_1, \dots, X_n)^{-1}(B) = \{\omega \in \Omega : (X_1(\omega), \dots, X_n(\omega)) \in B\} \in \mathcal{F} \quad \forall B \in \mathcal{B}(\mathbb{R}^n).$$



$$\forall B \in \mathcal{B}(\mathbb{R}^n), \quad (X_1, \dots, X_n)^{-1}(B) \in \mathcal{F}$$

Joint Probability Law of Multiple Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Joint Probability Law of Multiple Random Variables)


Given two random variables X_1, \dots, X_n defined with respect to \mathcal{F} , their **joint probability law** is the probability measure $\mathbb{P}_{X_1, \dots, X_n} : \mathcal{B}(\mathbb{R}^n) \rightarrow [0, 1]$ defined as

$$\mathbb{P}_{X_1, \dots, X_n}(B) = \mathbb{P}(\{\omega \in \Omega : (X_1(\omega), \dots, X_n(\omega)) \in B\}), \quad B \in \mathcal{B}(\mathbb{R}^n).$$

Remarks:

- $\mathbb{P}_{X_1, \dots, X_n}$ is the **probability law** of the random variable (X_1, \dots, X_n)
- $\mathbb{P}_{X_1, \dots, X_n}$ gives the **full probabilistic description** of (X_1, \dots, X_n)

Joint Probability Law of Multiple Random Variables



$$\mathbb{P}_{X_1, \dots, X_n}(B) = \mathbb{P}((X_1, \dots, X_n)^{-1}(B)) \quad \forall B \in \mathcal{B}(\mathbb{R}^n)$$

Marginal Law/CDF from Joint Law/CDF

Given joint CDF/law, we may extract the marginal CDFs/laws

The converse is not possible in general

CDF	Law
$F_{X_1, \dots, X_n} = \{F_{X_1, \dots, X_n}(x_1, \dots, x_n) : x_1, \dots, x_n \in \mathbb{R}\}$	$\mathbb{P}_{X_1, \dots, X_n} = \{\mathbb{P}_{X_1, \dots, X_n}(B) : B \in \mathcal{B}(\mathbb{R}^n)\}$
$F_{X_1}(x_1) = \lim_{\substack{x_2 \rightarrow +\infty \\ x_3 \rightarrow +\infty \\ \vdots \\ x_n \rightarrow +\infty}} F_{X_1, \dots, X_n}(x_1, x_2, \dots, x_n)$	$\mathbb{P}_{X_1}(A) = \mathbb{P}_{X_1, \dots, X_n}(A \times \mathbb{R} \times \dots \times \mathbb{R})$
$F_{X_2}(x_2) = \lim_{\substack{x_1 \rightarrow +\infty \\ x_3 \rightarrow +\infty \\ \vdots \\ x_n \rightarrow +\infty}} F_{X_1, \dots, X_n}(x_1, x_2, \dots, x_n)$	$\mathbb{P}_{X_2}(A) = \mathbb{P}_{X_1, \dots, X_n}(\mathbb{R} \times A \times \mathbb{R} \times \dots \times \mathbb{R})$

Table: Marginal law/CDF from joint law/CDF.

Independence of Random Variables

Independence of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Two Random Variables)

Two random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined with respect to \mathcal{F} are said to be **independent** if

$$\{X \in B_1\} \perp \{Y \in B_2\} \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

That is,

$$\mathbb{P}(\{X \in B_1\} \cap \{Y \in B_2\}) = \mathbb{P}(\{X \in B_1\}) \cdot \mathbb{P}(\{Y \in B_2\}) \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

Equivalently,

$$\mathbb{P}_{X,Y}(B_1 \times B_2) = \mathbb{P}_X(B_1) \cdot \mathbb{P}_X(B_2) \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

Independence of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Two Random Variables)

$$X \perp\!\!\!\perp Y \iff \{X \in B_1\} \perp\!\!\!\perp \{Y \in B_2\} \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

B_1	B_2	$\{X \in B_1\}$	$\{Y \in B_2\}$	Implication
$(-\infty, x]$	$(-\infty, y]$			

Independence of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Two Random Variables)

$$X \perp\!\!\!\perp Y \iff \{X \in B_1\} \perp\!\!\!\perp \{Y \in B_2\} \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

B_1	B_2	$\{X \in B_1\}$	$\{Y \in B_2\}$	Implication
$(-\infty, x]$	$(-\infty, y]$	$\{X \leq x\}$	$\{Y \leq y\}$	$F_{X,Y}(x, y) = F_X(x) \cdot F_Y(y)$
$(-\infty, x]$	$(y, +\infty)$			

Independence of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Two Random Variables)

$$X \perp\!\!\!\perp Y \iff \{X \in B_1\} \perp\!\!\!\perp \{Y \in B_2\} \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

B_1	B_2	$\{X \in B_1\}$	$\{Y \in B_2\}$	Implication
$(-\infty, x]$	$(-\infty, y]$	$\{X \leq x\}$	$\{Y \leq y\}$	$F_{X,Y}(x, y) = F_X(x) \cdot F_Y(y)$
$(-\infty, x]$	$(y, +\infty)$	$\{X \leq x\}$	$\{Y > y\}$	$\mathbb{P}(\{X \leq x, Y > y\}) = \mathbb{P}(\{X \leq x\}) \cdot \mathbb{P}(\{Y > y\})$
$(-\infty, x]$	$\{y\}$			

Independence of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Two Random Variables)

$$X \perp\!\!\!\perp Y \iff \{X \in B_1\} \perp\!\!\!\perp \{Y \in B_2\} \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

B_1	B_2	$\{X \in B_1\}$	$\{Y \in B_2\}$	Implication
$(-\infty, x]$	$(-\infty, y]$	$\{X \leq x\}$	$\{Y \leq y\}$	$F_{X,Y}(x, y) = F_X(x) \cdot F_Y(y)$
$(-\infty, x]$	$(y, +\infty)$	$\{X \leq x\}$	$\{Y > y\}$	$\mathbb{P}(\{X \leq x, Y > y\}) = \mathbb{P}(\{X \leq x\}) \cdot \mathbb{P}(\{Y > y\})$
$(-\infty, x]$	$\{y\}$	$\{X \leq x\}$	$\{Y = y\}$	$\mathbb{P}(\{X \leq x, Y = y\}) = \mathbb{P}(\{X \leq x\}) \cdot \mathbb{P}(\{Y = y\})$
$(-\infty, x]$	(a, b)			

Independence of Two Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Two Random Variables)

$$X \perp\!\!\!\perp Y \iff \{X \in B_1\} \perp\!\!\!\perp \{Y \in B_2\} \quad \forall B_1, B_2 \in \mathcal{B}(\mathbb{R}).$$

B_1	B_2	$\{X \in B_1\}$	$\{Y \in B_2\}$	Implication
$(-\infty, x]$	$(-\infty, y]$	$\{X \leq x\}$	$\{Y \leq y\}$	$F_{X,Y}(x, y) = F_X(x) \cdot F_Y(y)$
$(-\infty, x]$	$(y, +\infty)$	$\{X \leq x\}$	$\{Y > y\}$	$\mathbb{P}(\{X \leq x, Y > y\}) = \mathbb{P}(\{X \leq x\}) \cdot \mathbb{P}(\{Y > y\})$
$(-\infty, x]$	$\{y\}$	$\{X \leq x\}$	$\{Y = y\}$	$\mathbb{P}(\{X \leq x, Y = y\}) = \mathbb{P}(\{X \leq x\}) \cdot \mathbb{P}(\{Y = y\})$
$(-\infty, x]$	(a, b)	$\{X \leq x\}$	$\{a < Y < b\}$...

Table: Independence of two random variables from various angles.

Independence and Joint CDFs

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Let $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ be random variables defined with respect to \mathcal{F} .

Proposition (Independence and Joint CDFs)

The following statements are equivalent.

1. $\mathbb{P}_{X,Y}(B_1 \times B_2) = \mathbb{P}_X(B_1) \cdot \mathbb{P}_Y(B_2)$ for all $B_1, B_2 \in \mathcal{B}(\mathbb{R})$.
2. $F_{X,Y}(x, y) = F_X(x) \cdot F_Y(y)$ for all $x, y \in \mathbb{R}$.

Independence and Joint CDFs

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Let $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ be random variables defined with respect to \mathcal{F} .

Proposition (Independence and Joint CDFs)

The following statements are equivalent.

1. $\mathbb{P}_{X,Y}(B_1 \times B_2) = \mathbb{P}_X(B_1) \cdot \mathbb{P}_Y(B_2)$ for all $B_1, B_2 \in \mathcal{B}(\mathbb{R})$.
2. $F_{X,Y}(x, y) = F_X(x) \cdot F_Y(y)$ for all $x, y \in \mathbb{R}$.

Interpretation of $2 \implies 1$

If the joint probability law products out on the collection

$$\mathcal{D} = \left\{ (-\infty, x] \times (-\infty, y] : x, y \in \mathbb{R} \right\},$$

then it has to product out on $\sigma(\mathcal{D}) = \mathcal{B}(\mathbb{R}^2)$ (by Caratheodory's extension theorem)

Independence of Multiple Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Independence of Multiple Random Variables)

1. Random variables X_1, \dots, X_n , all defined with respect to \mathcal{F} , are **independent** if

$$\mathbb{P}_{X_1, \dots, X_n}(B_1 \times \dots \times B_n) = \prod_{i=1}^n \mathbb{P}_{X_i}(B_i) \quad \forall B_1, \dots, B_n \in \mathcal{B}(\mathbb{R}).$$

2. For an arbitrary index set \mathcal{I} , the collection of random variables $\{X_i : i \in \mathcal{I}\}$ is **independent** if every finite subset of them is independent.

Independent and Identically Distributed (i.i.d.) Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Let X_1, \dots, X_n be random variables defined with respect to \mathcal{F} .

Definition (i.i.d. Random Variables)

X_1, \dots, X_n are said to be **independent and identically distributed (i.i.d.)** if

1. X_1, \dots, X_n are independent.
2. $F_{X_i} = F_{X_j}$ for all $i \neq j$ (**identical CDFs**).

Jointly Discrete Random Variables

Jointly Discrete Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Jointly Discrete Random Variables)

Random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined with respect to \mathcal{F} are said to be **jointly discrete** if $(X, Y) : \Omega \rightarrow \mathbb{R}^2$ is a discrete random variable.

- X discrete \iff

Jointly Discrete Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Jointly Discrete Random Variables)

Random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined with respect to \mathcal{F} are said to be **jointly discrete** if $(X, Y) : \Omega \rightarrow \mathbb{R}^2$ is a discrete random variable.

- X discrete $\iff \mathbb{P}_X(E_1) = \mathbb{P}(\{X \in E_1\}) = 1$ for some countable set $E_1 \subset \mathbb{R}$
- Y discrete \iff

Jointly Discrete Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Jointly Discrete Random Variables)

Random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined with respect to \mathcal{F} are said to be **jointly discrete** if $(X, Y) : \Omega \rightarrow \mathbb{R}^2$ is a discrete random variable.

- X discrete $\iff \mathbb{P}_X(E_1) = \mathbb{P}(\{X \in E_1\}) = 1$ for some countable set $E_1 \subset \mathbb{R}$
- Y discrete $\iff \mathbb{P}_Y(E_2) = \mathbb{P}(\{Y \in E_2\}) = 1$ for some countable set $E_2 \subset \mathbb{R}$
- E_1 countable, E_2 countable $\implies E_1 \times E_2$ countable (**exercise!**)
- $\mathbb{P}_{X,Y}(E_1 \times E_2) = \mathbb{P}(\{X \in E_1\} \cap \{Y \in E_2\}) = 1$

Jointly Discrete Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.


Definition (Jointly Discrete Random Variables)

Random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined with respect to \mathcal{F} are said to be **jointly discrete** if $(X, Y) : \Omega \rightarrow \mathbb{R}^2$ is a discrete random variable.

- X discrete $\iff \mathbb{P}_X(E_1) = \mathbb{P}(\{X \in E_1\}) = 1$ for some countable set $E_1 \subset \mathbb{R}$
- Y discrete $\iff \mathbb{P}_Y(E_2) = \mathbb{P}(\{Y \in E_2\}) = 1$ for some countable set $E_2 \subset \mathbb{R}$
- E_1 countable, E_2 countable $\implies E_1 \times E_2$ countable (**exercise!**)
- $\mathbb{P}_{X,Y}(E_1 \times E_2) = \mathbb{P}(\{X \in E_1\} \cap \{Y \in E_2\}) = 1$

X discrete, Y discrete $\implies (X, Y)$ discrete

Picture of Jointly Discrete Random Variables



Joint PMF

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Joint PMF)

The joint PMF of jointly discrete random variables $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ defined on \mathcal{F} is a function $p_{X,Y} : \mathbb{R}^2 \rightarrow [0, 1]$ defined as

$$p_{X,Y}(\mathbf{x}, \mathbf{y}) = \mathbb{P}(\{X = \mathbf{x}\} \cap \{Y = \mathbf{y}\}), \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}.$$

Note:

$$\mathbb{P}(\{(X, Y) \in E_1 \times E_2\}) = \sum_{\mathbf{x} \in E_1} \sum_{\mathbf{y} \in E_2} p_{X,Y}(\mathbf{x}, \mathbf{y}) = 1,$$


$$\mathbb{P}(\{(X, Y) \in B\}) = \sum_{(\mathbf{x}, \mathbf{y}) \in B \cap (E_1 \times E_2)} p_{X,Y}(\mathbf{x}, \mathbf{y}), \quad B \subseteq \mathbb{R}^2.$$

Properties of Joint PMF

- $\sum_{x \in E_1} \sum_{y \in E_2} p_{X,Y}(x, y) = 1.$

- $p_X(x) = \sum_{y \in E_2} p_{X,Y}(x, y), \quad x \in \mathbb{R}$

- $p_Y(y) = \sum_{x \in E_1} p_{X,Y}(x, y), \quad y \in \mathbb{R}$



Conditional PMF

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Conditional PMF)

Let X, Y be jointly discrete random variables defined with respect to \mathcal{F} . Fix $y \in \mathbb{R}$ such that $p_Y(y) = \mathbb{P}(\{Y = y\}) > 0$. The **conditional PMF** of X , conditioned on the event $\{Y = y\}$, is a function $p_{X|Y=y} : \mathbb{R} \rightarrow [0, 1]$ defined as

$$p_{X|Y=y}(x) = \frac{\mathbb{P}(\{X = x\} \cap \{Y = y\})}{\mathbb{P}(\{Y = y\})} = \frac{p_{X,Y}(x, y)}{p_Y(y)}, \quad x \in \mathbb{R},$$

defined for all $y \in \mathbb{R}$ such that $p_Y(y) = \mathbb{P}(\{Y = y\}) > 0$.

Conditional PMF

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition (Conditional PMF)


Let X, Y be jointly discrete random variables defined with respect to \mathcal{F} . Fix $y \in \mathbb{R}$ such that $p_Y(y) = \mathbb{P}(\{Y = y\}) > 0$. The **conditional PMF** of X , conditioned on the event $\{Y = y\}$, is a function $p_{X|Y=y} : \mathbb{R} \rightarrow [0, 1]$ defined as

$$p_{X|Y=y}(x) = \frac{\mathbb{P}(\{X = x\} \cap \{Y = y\})}{\mathbb{P}(\{Y = y\})} = \frac{p_{X,Y}(x, y)}{p_Y(y)}, \quad x \in \mathbb{R},$$

defined for all $y \in \mathbb{R}$ such that $p_Y(y) = \mathbb{P}(\{Y = y\}) > 0$.

Remark: In textbooks, $p_{X|Y=y}(x)$ is commonly denoted as **$p_{X|Y}(x|y)$**

Conditional PMF



Independence of Two Discrete Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Theorem

Let $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ be discrete random variables with respect to \mathcal{F} . The following statements are equivalent.

1. $X \perp\!\!\!\perp Y$.
2. $\{X = x\} \perp\!\!\!\perp \{Y = y\}$ for all $x, y \in \mathbb{R}$.
3. $p_{X,Y}(x, y) = p_X(x) \cdot p_Y(y)$ for all $x, y \in \mathbb{R}$.
4. For all $y \in \mathbb{R}$ such that $p_Y(y) > 0$,

$$p_{X|Y=y}(x) = p_X(x) \quad \forall x \in \mathbb{R}.$$

Jointly Continuous Random Variables

Jointly Continuous Random Variables

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Let $X : \Omega \rightarrow \mathbb{R}$ and $Y : \Omega \rightarrow \mathbb{R}$ be random variables defined with respect to \mathcal{F} .

Definition (Jointly Continuous Random Variables)

X and Y are said to be **jointly continuous** if $(X, Y) : \Omega \rightarrow \mathbb{R}^2$ is a continuous random variable, i.e., there exists a function $f_{X,Y} : \mathbb{R}^2 \rightarrow [0, +\infty)$ such that the joint CDF of X and Y may be expressed as

$$F_{X,Y}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{X,Y}(u, v) dv du \quad \forall x, y \in \mathbb{R}.$$

The function $f_{X,Y}$ is called the **joint PDF** of X and Y .

Remark:

X continuous, Y continuous $\not\Rightarrow X, Y$ jointly continuous