



# Probability and Stochastic Processes

Lecture 02: Sample Space, Algebra,  $\sigma$ -Algebra

**Karthik P. N.**

**Assistant Professor, Department of AI**

**Email: [pnkarthik@ai.iith.ac.in](mailto:pnkarthik@ai.iith.ac.in)**

01 August 2024

## Probability Theory – Humble Beginnings

- Bernoulli (1713) and de Moivre (1718) gave the first definition of probability:

$$\text{probability of an event} = \frac{\text{\# favourable outcomes}}{\text{total number of outcomes}}.$$


- Cournot (1843):  
“An event with very small probability is morally impossible; an event with very high probability is morally certain.”
- French mathematicians of the day were satisfied with the “frequentist” approach to probability, but not the German and English mathematicians of the day
- Frequentist approach could not satisfactorily explain certain paradoxes

## Why Axiomatic Theory? Bertrand's Paradox

Take a circle with unit radius and inscribe an equilateral triangle in it. Draw a random chord. What is the probability that the length of the “random chord” is greater than  $\sqrt{3}$ ?

Bertrand's perfectly valid arguments:

- Mid-point of chord should lie inside incircle of radius  $1/2$



## Why Axiomatic Theory? Bertrand's Paradox


Take a circle with unit radius and inscribe an equilateral triangle in it. Draw a random chord. What is the probability that the length of the “random chord” is greater than  $\sqrt{3}$ ?

Bertrand's perfectly valid arguments:

- Mid-point of chord should lie inside incircle of radius  $1/2$

**Answer:  $1/4$**

- Angle between chord and tangent at A should be between  $\pi/3$  and  $2\pi/3$



## Why Axiomatic Theory? Bertrand's Paradox

Take a circle with unit radius and inscribe an equilateral triangle in it. Draw a random chord. What is the probability that the length of the “random chord” is greater than  $\sqrt{3}$ ?

Bertrand's perfectly valid arguments:


- Mid-point of chord should lie inside incircle of radius  $1/2$

**Answer:  $1/4$**

- Angle between chord and tangent at A should be between  $\pi/3$  and  $2\pi/3$

**Answer:  $1/3$**


- Mid-point of chord should be between O and projection of O onto side BC



## Why Axiomatic Theory? Bertrand's Paradox

Take a circle with unit radius and inscribe an equilateral triangle in it. Draw a random chord. What is the probability that the length of the “random chord” is greater than  $\sqrt{3}$ ?

Bertrand's perfectly valid arguments:



- Mid-point of chord should lie inside incircle of radius  $1/2$   
**Answer:  $1/4$**
- Angle between chord and tangent at A should be between  $\pi/3$  and  $2\pi/3$   
**Answer:  $1/3$**
- Mid-point of chord should be between O and projection of O onto side BC  
**Answer:  $1/2$**

## Borel to the Rescue

- Contributions to Measure Theory by Borel (1894) provided a shift in perspective
- Countable unions played a key role in Borel's theory
- Kolmogorov's genius was in applying Borel's theory to formalise the axioms of probability, laying the foundation stone for modern probability theory
- For more details on the history of probability, see [[Shafer and Vovk, 2018](#)] and [[Kolmogorov, 2004](#)]

## Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – **source of randomness**



# Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – **source of randomness**

## Definition (Sample Space)

The sample space of a random experiment is the set of all possible outcomes of the random experiment.

# Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – **source of randomness**

## Definition (Sample Space)

The sample space of a random experiment is the set of all possible outcomes of the random experiment.

Example: Tossing a coin once

- If our interest is in the face that shows, then  $\Omega = \{H, T\}$

# Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – **source of randomness**

## Definition (Sample Space)

The sample space of a random experiment is the set of all possible outcomes of the random experiment.

Example: Tossing a coin once

- If our interest is in the face that shows, then  $\Omega = \{H, T\}$

# Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – **source of randomness**

## Definition (Sample Space)

The sample space of a random experiment is the set of all possible outcomes of the random experiment.

Example: Tossing a coin once

- If our interest is in the face that shows, then  $\Omega = \{H, T\}$
- If our interest is in the velocity with which the coin lands on ground, then  $\Omega = [0, \infty) = \mathbb{R}_+$

# Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – **source of randomness**

## Definition (Sample Space)

The sample space of a random experiment is the set of all possible outcomes of the random experiment.

Example: Tossing a coin once

- If our interest is in the face that shows, then  $\Omega = \{H, T\}$
- If our interest is in the velocity with which the coin lands on ground, then  $\Omega = [0, \infty) = \mathbb{R}_+$

# Sample Space

We begin with two universally accepted entities:

- Random experiment
- Outcome (denoted by  $\omega$ ) – source of randomness

## Definition (Sample Space)

The sample space of a random experiment is the set of all possible outcomes of the random experiment.  $\Omega$

Example: Tossing a coin once

- If our interest is in the face that shows, then  $\Omega = \{H, T\}$
- If our interest is in the velocity with which the coin lands on ground, then  $\Omega = [0, \infty) = \mathbb{R}_+$
- If our interest is in the number of times coin flips in air, then  $\Omega = \mathbb{N}$



Example: Toss a coin  $n$  times, for some  $n < \infty$ .  
Interest: faces that show up

Example: Toss a coin  $n$  times, for some  $n < \infty$ .

Interest: faces that show up

$$\Omega = \{H, T\}^n$$



Example: Toss a coin  $n$  times, for some  $n < \infty$ .

Interest: faces that show up

$$\Omega = \{H, T\}^n$$

*countably*

Example: Toss a coin  $\underbrace{\hspace{1cm}}$  infinitely many times.

Interest: faces that show up

Example: Toss a coin  $n$  times, for some  $n < \infty$ .

Interest: faces that show up

$$\Omega = \{H, T\}^n$$

Example: Toss a coin infinitely many times.

Interest: faces that show up

$$\Omega = \{H, T\}^\infty \quad \{H, T\}^{\mathbb{N}}$$

# Event

## Informal Definition (Event)

Informally,<sup>a</sup> an event is a subset of outcomes “of interest” to us.

---

<sup>a</sup>We shall give a more formal definition of an event later.

Example: Toss a coin 3 times; interest is in the faces that show up

## Informal Definition (Event)

Informally,<sup>a</sup> an event is a subset of outcomes “of interest” to us.

---

<sup>a</sup>We shall give a more formal definition of an event later.

Example: Toss a coin 3 times; interest is in the faces that show up  
 $\Omega = \{H, T\}^3 = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$

## Informal Definition (Event)

Informally,<sup>a</sup> an event is a subset of outcomes “of interest” to us.

---

<sup>a</sup>We shall give a more formal definition of an event later.

Example: Toss a coin 3 times; interest is in the faces that show up

$$\Omega = \{H, T\}^3 = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$$

Event  $A$  of interest: at least 2 heads show up

## Informal Definition (Event)

Informally,<sup>a</sup> an event is a subset of outcomes “of interest” to us.

---

<sup>a</sup>We shall give a more formal definition of an event later.

Example: Toss a coin 3 times; interest is in the faces that show up

$$\Omega = \{H, T\}^3 = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$$

Event A of interest: at least 2 heads show up

$$E = \{HHH, THH, HTH, HHT\}$$

# Event

## Informal Definition (Event)

Informally,<sup>a</sup> an event is a subset of outcomes “of interest” to us.

---

<sup>a</sup>We shall give a more formal definition of an event later.

Example: Toss a coin 3 times; interest is in the faces that show up

$$\Omega = \{H, T\}^3 = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$$

Event  $A$  of interest: at least 2 heads show up

$$E = \{HHH, THH, HTH, HHT\}$$

## Note

If an outcome  $\omega \in E$  occurs, we say that the event  $A$  occurs.

## References



Kolmogorov, A. N. (2004).

From the heritage of AN Kolmogorov: The theory of probability.  
*Theory of Probability & Its Applications*, 48(2):191–220.



Shafer, G. and Vovk, V. (2018).

The origins and legacy of Kolmogorov's Grundbegriffe.  
*arXiv preprint arXiv:1802.06071*.