



Sri Chaitanya IIT Academy.,India.

☆ A.P ☆ T.S ☆ KARNATAKA ☆ TAMILNADU ☆ MAHARASTRA ☆ DELHI ☆ RANCHI

A right Choice for the Real Aspirant**ICON Central Office - Madhapur - Hyderabad****SEC: Sr.S60, ELITE ,TARGET & LIIT****JEE-MAIN****Date: 25-05-2022****Time: 09.00Am to 12.00Pm****GTM-15****Max. Marks: 300**

KEY SHEET

PHYSICS

1)	2	2)	2	3)	1	4)	3	5)	2
6)	2	7)	4	8)	1	9)	3	10)	3
11)	4	12)	2	13)	3	14)	4	15)	2
16)	1	17)	1	18)	3	19)	4	20)	1
21)	2	22)	20	23)	4	24)	35	25)	3
26)	4	27)	19	28)	1	29)	-113	30)	40

CHEMISTRY

31)	2	32)	2	33)	3	34)	1	35)	1
36)	2	37)	4	38)	1	39)	2	40)	4
41)	3	42)	2	43)	4	44)	4	45)	4
46)	2	47)	4	48)	3	49)	3	50)	1
51)	3	52)	8	53)	462	54)	6	55)	4
56)	3	57)	30	58)	2	59)	3	60)	8

MATHEMATICS

61)	3	62)	3	63)	3	64)	1	65)	4
66)	3	67)	2	68)	1	69)	1	70)	1
71)	2	72)	4	73)	3	74)	1	75)	3
76)	3	77)	3	78)	3	79)	4	80)	4
81)	1	82)	5	83)	3	84)	6	85)	1
86)	0	87)	13	88)	1	89)	11	90)	10



SOLUTIONS

PHYSICS

1. Conceptual

2. Conceptual

3.

$$t = \sqrt{\frac{L}{5}} + \frac{L}{300} \quad dt = \frac{1}{\sqrt{5}} \frac{1}{2} L^{-1/2} dL + \left(\frac{1}{300} dL \right) dt = \frac{1}{2\sqrt{5}} \frac{1}{\sqrt{20}} dL + \frac{dL}{300} = 0.01$$

$$dL \left(\frac{1}{20} + \frac{1}{300} \right) = 0.01 \quad dL \left[\frac{15}{300} \right] = 0.01 \quad dL = \frac{3}{16}$$

$$\frac{dL}{L} \times 100 = \frac{3}{16} \times \frac{1}{20} \times 100 = \frac{15}{16} \approx 1\%$$

4. Given, $g = 10 \text{ m/s}^2$

Equation of trajectory of the projectile. $y = 2x - 9x^2$...(i)

In projectile motion, equation of trajectory is given by

$$y = x \tan \theta_0 - \frac{gx^2}{2v_0^2 \cos^2 \theta_0} \quad \dots(ii)$$


By comparison of Eqs. (i) and (ii), we get

$$\tan \theta_0 = 2 \quad \dots(iii)$$

and $\frac{g}{2v_0^2 \cos^2 \theta_0} = 9$ or $v_0^2 = \frac{g}{9 \times 2 \cos^2 \theta_0} \quad \dots(iv)$

From Eq. (iii), we can get value of $\cos \theta$ and $\sin \theta$

$$\cos \theta_0 = \frac{1}{\sqrt{5}} \quad \text{and} \quad \sin \theta_0 = \frac{2}{\sqrt{5}} \quad (v)$$



Using value of $\cos \theta_0$ from Eq. (v) to Eq. (iv), we get

$$v_0^2 = \frac{10 \times (\sqrt{5})^2}{2 \times (1)^2 \times 9} = \frac{10 \times 5}{2 \times 9} \Rightarrow v_0^2 = \frac{25}{9} \text{ or } v_0 = \frac{5}{3} \text{ m/s} \quad \dots(vi)$$

From Eq. (v), we get $\theta_0 = \cos^{-1} \left(\frac{1}{\sqrt{5}} \right)$

5. Components of velocity at an instant of times t of a body projected at an angle θ is

$$v_x = u \cos \theta + g_x t \quad \text{and} \quad v_y = u \sin \theta + g_y t$$

Here, components of velocity at $t = 1 \text{ s}$, is

$$v_x = u \cos 60^\circ + 0 \quad (\text{as } g_x = 0) = 10 \times \frac{1}{2} = 5 \text{ m/s and } v_y = u \sin 60^\circ + (-10) \times (1)$$

$$= 5\sqrt{3} - 10 \Rightarrow |v_y| = |10 - 5\sqrt{3}| \text{ m/s}$$

Now, angle made by the velocity vector at time of $t = 1 \text{ s}$




$$|\tan \alpha| = \left| \frac{V_y}{V_x} \right| = \frac{|10 - 5\sqrt{3}|}{5} \Rightarrow \tan \alpha = |2 - \sqrt{3}| \text{ or } \alpha = 15^\circ$$

\therefore Radius of curvature of the trajectory of the projected body, $R = v^2 / g \cos \alpha$

$$= \frac{(5)^2 + (10 - 5\sqrt{3})^2}{10 \times 0.97} \left(\because v^2 = v_x^2 + v_y^2 \text{ and } \cos 15^\circ = 0.97 \right) \Rightarrow R = 2.77 \text{ m} \approx 2.8 \text{ m}$$

6. The given situation is shown below.



Here, range, $OB = 2h \tan \alpha = \frac{u^2 \sin 2\theta}{g}$

and $h = \frac{u^2 \sin^2 \theta}{2g} \Rightarrow \frac{2h \tan \alpha}{h} = \frac{\frac{u^2 \sin 2\theta}{g}}{\frac{u^2 \sin^2 \theta}{2g}} \Rightarrow \tan \theta = 2 \cot \alpha$

$$\Rightarrow \sin \theta = \frac{2 \cos \alpha}{\sqrt{1 + \cot^2 \alpha}} \Rightarrow h = \frac{u^2 \sin^2 \theta}{2g}$$

or $\frac{u^2}{2g} = \frac{h}{\sin^2 \theta} = \frac{h}{4 \cot^2 \alpha} (1 + 4 \cot^2 \alpha) = \frac{\tan^2 \alpha}{2} + 1$

$$\Rightarrow \frac{u^2}{gh} = \frac{\tan^2 \alpha}{2} + 2 \Rightarrow \frac{u^2}{gh} - \frac{\tan^2 \alpha}{2} = 2$$

7. Conceptual


8. Conceptual

9. Let acceleration of mass m relative to wedge down the plane is a_r . Its absolute acceleration in horizontal direction is $a_r \cos 60^\circ - a$ (towards right.) Hence, let N be the normal reaction between the mass and the wedge. Then

$$N \sin \theta = Ma = m(a_r \cos 60^\circ - a)$$

or $a_r = \frac{(M + m)a}{m \cos 60^\circ} = \frac{2(M + m)a}{m}$

10. $T_1 \geq \mu mg$



$$T_2 - 2T_1 \geq \mu mg$$

or $T_2 - 2\mu mg \geq \mu mg$ (Putting $T_1 = \mu mg$)

or $T_2 \geq 3\mu mg$

Now, $F - 2T_2 \geq \mu mg$

or $F - 6\mu mg \geq \mu mg$ (Putting $T_2 = 3\mu mg$)

or $F \geq 7\mu mg$



11. Conceptual

12. Let, N_1 = normal reaction at A and N_2 = normal reaction at B

$$\frac{\sqrt{3}N_1}{2} - \frac{mg}{2} - \frac{N_2\sqrt{3}}{2} = ma \quad \frac{\sqrt{3}N_1}{2} - \frac{\sqrt{3}N_2}{2} = ma + \frac{mg}{2} \Rightarrow \frac{N_1 + N_2}{2} = \frac{mg\sqrt{3}}{2}$$

$$\Rightarrow \frac{\sqrt{3}N_1}{2} + \frac{\sqrt{3}N_2}{2} = \frac{3mg}{2}$$

From Eqs. (i) and (ii), we have $\sqrt{3}N_1 = 2mg + ma$, $\sqrt{3}N_2 = mg - ma$

Since cylinder does not lose contact at B, so

$$N_2 \geq 0 \Rightarrow a \leq g \Rightarrow a_{\max} = g$$

13. Conceptual

14. $F \propto s^{-1/3}$

i.e. acceleration $a \propto s^{-1/3}$

$$\text{or } v \frac{dv}{ds} = Ks^{-1/3} \quad \text{or } v^2 \propto s^{2/3}$$

$$\text{or } v \propto s^{1/3}$$

Now, $P = F.v$

$$\text{or } P \propto s^{-1/3} \cdot s^{1/3} \quad \text{or } P \propto s^0$$

i.e. power is independent of s.

15. Conceptual

$$16. \quad a = \frac{F_{\text{net}}}{m} = \frac{mg \sin \theta - (\mu_0 x)(mg \cos \theta)}{m} = g \sin \theta - (\mu_0 g \cos \theta)x$$

$$\therefore v \cdot \frac{dv}{dx} = g \sin \theta - (\mu_0 g \cos \theta)x$$


$$\therefore \int_0^0 v dv = \int_0^{x_{\max}} [g \sin \theta - (\mu_0 g \cos \theta)x] dx$$

Solving this integration, we get

$$x_{\max} = \frac{2 \tan \theta}{\mu_0}$$

17. From work energy theorem,

$$W_g + W_{\text{pseudo force}} = K_f - K_i$$




$$mgR \cos \theta - maR(1 - \sin \theta) = \frac{1}{2}mv^2 \Rightarrow v = \sqrt{2gR(\sin \theta + \cos \theta - 1)}$$

$$v \text{ is maximum when } \frac{dv}{d\theta} = 0 \Rightarrow \theta = 45^\circ$$

18. Apply work energy theorem

19. Apply work energy theorem

20. In equilibrium, (with respect to cylinder)



$$\tan \theta = \frac{2mg}{mg}$$

$$\text{or } \theta = \tan^{-1}(2)$$

$$\therefore \text{Maximum angular displacement} = 2\theta = 2 \tan^{-1}(2)$$

21. Conceptual

22. Conceptual

$$23. \quad E_Q = \frac{G \left(\rho \cdot \frac{4}{3} \pi R^3 \right)}{(2R)^2} + \frac{G \left[2\rho \left\{ \frac{4}{3} \pi (2R)^3 - \frac{4}{3} \pi (R)^3 \right\} \right]}{(2R)^2}$$

$$E_P = \frac{G \left(\rho \cdot \frac{4}{3} \pi R^3 \right)}{R^2} \quad E_Q / E_P = 15 / 4 = 3.75$$

$$24. \quad \text{Density of wire, } d = 9 \times 10^{-3} \frac{\text{kg}}{\text{cm}^3} = 9 \times 10^3 \text{ kg} / \text{m}^3$$

$$\text{Strain in the wire, } \varepsilon = 4.9 \times 10^{-4}$$

$$\text{Young's modulus of wire is } Y = 9 \times 10^{10} \frac{\text{N}}{\text{m}^2}$$

$$\text{Lowest frequency of vibration in wire will be } f = \frac{1}{2L} \sqrt{\frac{T}{(M/L)}}$$

$$\text{Now, } \frac{T}{M/L} = \frac{T}{V\rho/L} = \frac{T}{\left(\frac{LA\rho}{L} \right)} = \frac{T}{A\rho}$$

$$\text{But } \frac{T}{A} = \text{stress } Y \times \text{strain} = Y \times \varepsilon \Rightarrow \frac{T}{M/L} = \frac{Y \times \varepsilon}{\rho}$$

So, from Eq (i), frequency will be

$$f = \frac{1}{2L} \sqrt{\frac{T}{(M/L)}} = \frac{1}{2L} \sqrt{\frac{Y \times \varepsilon}{\rho}} = \frac{1}{2 \times 1} \times \sqrt{\frac{9 \times 10^{10} \times 49 \times 10^{-4}}{9 \times 10^3}} = 35 \text{ Hz}$$

25. Conceptual

26. Conceptual

27. Conceptual

28. Conceptual

29. Conceptual

30. Conceptual

**CHEMISTRY**

31. Formation of NaCl is an exothermic reaction.

$$32. \frac{\lambda_1}{\lambda_2} = \frac{(mv)_2}{(mv)_1}$$

33. $[ClF_2O]^+ - sp^3, [ClF_4O]^- - sp^3d^2$

34. $C_2H_5OH + 2Cr_2O_7^{2-} \rightarrow 2CO_2 + 4Cr^{3+}$

Eq. of $Cr_2O_7^{2-}$ = eq of C_2H_5OH

$$\frac{8.0 \times .05 \times 6}{1000} = \frac{10}{46/12} \quad W = \frac{8.0 \times .05 \times 46}{1000 \times 12} \times 6 = .0092 \quad \% = \frac{0.0092}{10} \times 100 = 0.092\%$$

35. As per Le Chatelier's principle, equilibrium moves in forward direction.

36. $C(\text{graphite}) + \frac{1}{2}O_2(g) \rightarrow CO(g); \Delta H = -ve$ Is not a combustion reaction

$$37. \Delta S = 2.303nR \log \frac{V_2}{V_1}$$

38. De-excitation of electron from 5th orbit to 4th orbit gives first line in Brackett series.

39. $HI_n \rightleftharpoons H^+ + I_n^-$

Acidic basic

Red Blue


$$K_a = \frac{[H^+][I_n^-]}{[HI_n]}$$

Red from 75% blue from 25% $\therefore [H^+] = 3 \times 10^{-5} \times \frac{75}{25} = 9 \times 10^{-5}$

When 75% blue and 25% red, then

$$[H^+] = 3 \times 10^{-5} \times \frac{25}{75} = 1 \times 10^{-5} \therefore \text{change is } \therefore [H^+] = 9 \times 10^{-5} - 1 \times 10^{-5} = 8 \times 10^{-5}$$


40. Localised electron pair on nitrogen.



42. Most stable alkene is the major product.


43. BeF_2 Soluble in water due to greater hydration energy of Be^{+2} , as compared to L.E

44. Al^{+3} ion further can not lose electrons so it cannot act as reducing agent.





46. For lead, (+2) is more stable due to inert pair effect
47. As BOD values increases, pollution increase.
48. As the electronegativity of central atom increases, bond angle increases.
49. Friedel Crafts alkylation is generally not preferred for the preparation of alkyl benzenes because electrophile undergoes rearrangement giving mixture of products and poly alkylation takes place as alkyl group is activating
50. More hyper conjugations.
51. I, III and IV
52. $2S_4N_4 \rightarrow 4N_2(g) + S_8(g)$
53. $\Delta G = 0$ at equilibrium hence $\Delta H = T\Delta S$ and $\therefore T = \frac{\Delta H}{\Delta S}$
54. Borax- $Na_2B_4O_7 \cdot 10H_2O$ and Kernite- $Na_2B_4O_7 \cdot 4H_2O$
55. a, b, e, g



57. $P_1V_1 = P_2V_2$

58. $q = ms\Delta T$

59. i, ii, iii.

60. KCN	HCN
Volume = 'x' ml	volume = 10ml
Molarity = 5	Molarity = 2m

$$\therefore P^H = -\log Ka + \log \frac{[salt]}{[Acid]}$$

$$9.6020 = -\log 5 \times 10^{-10} + \log \frac{5x/(10+x)}{20/(10+x)}$$

$$9.6020 = 9.3010 + \log \frac{x}{4}$$

$$\log \frac{x}{4} = 0.3010 = \log 2$$

$$x = 8ml$$

**MATHEMATICS**

61. Lines are $y - 1 = \tan \theta (x - 1)$ and $(y - 1) = \cot \theta (x - 1)$ multiply and compare

$$\sin 2\theta = \frac{2}{a+2}$$

62. $\lim_{n \rightarrow \infty} \sum_{r=1}^n \tan^{-1}(r^2 + r) - \tan^{-1}(r^2 - r) = \frac{\pi}{2}$


63. Let $g(x) = e^{-x} f(x)$ As $g''(x) > 0$ so $g'(x)$ is increasing

So, for $x < 1/4$, $g'(x) < g'(1/4) = 0$

$$\Rightarrow (f(x) - f(x))e^{-x} < 0 \Rightarrow f(x) < f(x) \text{ in } (0, 1/4)$$

64. $ST^2 = AS^2 + AT^2 - 2AS \cdot AT \cdot \cos \alpha$ and $AS \cdot AT = \frac{1}{2}bc$

65. Use half angle formulae and Heron's formula



$$\text{Area of rectangle} = \alpha \left(\frac{6-3\alpha}{2} - \alpha \right) = \alpha \left(\frac{6-5\alpha}{2} \right) \quad \frac{dA}{d\alpha} = 0 \Rightarrow \alpha = \frac{3}{5}$$

$$A_{\text{Max}} = \frac{3}{5} \times \left(\frac{3}{2} \right) = \frac{9}{10} \text{ Sq units}$$

67. Since $f''(x) > 0$

$\Rightarrow f'(x)$ is always increasing.

$$g'(x) = 2f'(2x^3 - 3x^2) \times (6x^2 - 6x) + f'(6x^2 - 4x^3 - 3)(12x - 12x^2)$$

$$= 12(x^2 - x) \cdot (f'(2x^3 - 3x^2) - f'(6x^2 - 4x^3 - 3))$$

$$= 12x(x-1) [f'(2x^3 - 3x^2) - f'(6x^2 - 4x^3 - 3)]$$

For increasing $g'(x) > 0$

$$\Rightarrow 2x^3 - 3x^2 > 6x^2 - 4x^3 - 3 \{ \because f'(x) \text{ is increasing} \}$$

$$\Rightarrow (x-1)^2 \left(x + \frac{1}{2} \right) > 0 \rightarrow x > -\frac{1}{2} \therefore x \in \left(-\frac{1}{2}, 0 \right) \cup (1, \infty)$$

Case-II If $0 < x < 1$

$$f'(2x^3 - 3x^2) < f'(6x^2 - 4x^3 - 3)$$

$$(x-1)^2 \left(x + \frac{1}{2} \right) < 0 \quad x < -\frac{1}{2}, \text{ so there is no solution}$$

$$\Rightarrow \text{Hence the values are } x \in \left(-\frac{1}{2}, 0 \right) \cup (1, \infty)$$



$$68. \begin{vmatrix} 5-\alpha & 0 & 0 \\ 0 & 3-\alpha & -2 \\ 0 & -1 & 2-\alpha \end{vmatrix} = 0 \Rightarrow \alpha = 1, 4, 5$$

$$69. \lambda = \frac{m_{21} - m_{11} - m_{22}}{3^{2012}} = \frac{2012 \times 3^{2012} - 3^{2012} - 3^{2012}}{3^{2012}} = 2010$$

$$70. \text{ Given } y = 1 + \frac{1}{x-1} + \frac{2x}{(x-1)(x-2)} + \frac{3x^2}{(x-1)(x-2)(x-3)}$$

$$= \left(1 + \frac{1}{x-1}\right) + \frac{2x}{(x-1)(x-2)} + \frac{3x^2}{(x-1)(x-2)(x-3)}$$

$$y = \frac{x^3}{(x-1)(x-2)(x-3)}$$

$$\log y = \log\left(\frac{x}{x-1}\right) + \log\left(\frac{x}{x-2}\right) + \log\left(\frac{x}{x-3}\right)$$

Diff. both sides w.r. t x, we get

$$\frac{dy}{dx} = \frac{y}{x} \left[\frac{1}{1-x} + \frac{2}{2-x} + \frac{3}{3-x} \right] \therefore -9y'(4) = 104$$

$$71. m \sin 2\alpha = m \sin\{(\alpha + \beta) + (\alpha - \beta)\} = \cos^2(\alpha - \beta) + m \cos(\alpha + \beta) \sin(\alpha - \beta)$$

$$m \sin 2\beta = m \sin\{(\alpha + \beta) - (\alpha - \beta)\} = \cos^2(\alpha - \beta) - m \cos(\alpha + \beta) \sin(\alpha - \beta)$$

$$\text{Now, } \frac{1}{1 - m \sin 2\alpha} + \frac{1}{1 - m \sin 2\beta}$$

$$\frac{1}{\sin^2(\alpha - \beta) - m \cos(\alpha + \beta) \sin(\alpha - \beta)} + \frac{1}{\sin^2(\alpha - \beta) + m \cos(\alpha + \beta) \sin(\alpha - \beta)} = \frac{2}{1 - m^2}$$

$$72. \text{ Continuous at } x = 1 \quad p + q = -2 + \frac{\pi}{4}$$

$$\text{differentiable at } x = 1 \quad 3p = \frac{1}{1+1} = \frac{1}{2} \quad p = \frac{1}{6}; q = \frac{\pi}{4} - \frac{13}{6}$$

$$73. (1+y)\cos^2 2x + 4\cos 2x + 3y - 1 = 0$$

$$\text{Since } \cos 2x \text{ is real, } 16 - 4(3y - 1)(1 + y) \geq 0 \quad \text{Or } 3y^2 + 2y - 5 \leq 0$$

$$\text{But } y = 1 \Rightarrow \cos 2x = -1 \text{ i.e. } x = \frac{\pi}{2} \text{ which is not permissible.}$$

$$74. \text{ Total disks in } A_n \text{ is } S_n = 3n(n+1) \text{ and length of side of hexagon is}$$

$$1 = 2r_n(n-1) + \frac{2r_n}{\sqrt{3}} \Rightarrow r_n = \frac{1}{2n-2 + \frac{2}{\sqrt{3}}}$$

$$75. \bar{d}_1 \bar{d}_2 = 0 \Rightarrow (\bar{a} + 2\bar{b}) \cdot (2\bar{a} - \bar{b}) = 0 \Rightarrow 2|\bar{a}|^2 + 3\bar{a} \cdot \bar{b} - 2|\bar{b}|^2 = 0$$

$$\bar{d}_3 \bar{d}_4 = 0 \Rightarrow (2\bar{a} + \bar{b}) \cdot (\bar{a} - 2\bar{b}) = 0 \Rightarrow 2|\bar{a}|^2 - 3\bar{a} \cdot \bar{b} - 2|\bar{b}|^2 = 0$$

$$\Rightarrow \bar{a} \cdot \bar{b} = 0 \quad \text{Now } [\bar{a} \bar{b} \bar{a} \times 2\bar{b}] = (\bar{a} \times \bar{b}) \cdot (\bar{a} \times 2\bar{b}) = 2|\bar{a}|^2 |\bar{b}|^2$$

$$76. A = \begin{bmatrix} e^t & e^{-t} \cos t & e^{-t} \sin t \\ e^t & -e^{-t}(\sin t + \cos t) & e^{-t}(\cos t - \sin t) \\ e^t & 2e^{-t} \sin t & -2e^{-t} \cos t \end{bmatrix}$$



$$|A| = e^t \cdot e^{-2t} \begin{vmatrix} 1 & \cos t & \sin t \\ 1 & -\sin t - \cos t & \cos t - \sin t \\ 1 & 2\sin t & -2\cos t \end{vmatrix}$$

$$|A| = e^{-t} \begin{vmatrix} 1 & \cos t & \sin t \\ 0 & -\sin t - 2\cos t & \cos t - 2\sin t \\ 0 & 2\sin t - \cos t & -2\cos t - \sin t \end{vmatrix}$$

$$|A| = e^{-t} (\sin^2 t + 4\cos^2 t + 4\sin t \cos t - 4\cos t + 4\sin^2 t + \cos^2 t)$$

$$|A| = e^{-t} (5\sin^2 t + 5\cos^2 t) = 5 \cdot e^{-t} \neq 0 \forall t \in R$$

77.

$$A = \begin{bmatrix} 0 & 1 \\ 3 & 0 \end{bmatrix} \Rightarrow A^2 = A \cdot A = \begin{bmatrix} 0 & 1 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 3 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\Rightarrow A^4 = A^2 \cdot A^2 = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 3^2 & 0 \\ 0 & 3^2 \end{bmatrix}$$

$$\Rightarrow A^8 = \begin{bmatrix} 3^4 & 0 \\ 0 & 3^4 \end{bmatrix} \text{ and } A^6 = A^4 \cdot A^2 = \begin{bmatrix} 3^2 & 0 \\ 0 & 3^2 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 3^3 & 0 \\ 0 & 3^3 \end{bmatrix}$$

$$\text{Let } V = \begin{bmatrix} x \\ y \end{bmatrix}$$

$$A^8 + A^6 + A^4 + A^2 + I$$

$$\begin{bmatrix} 81 & 0 \\ 0 & 81 \end{bmatrix} + \begin{bmatrix} 27 & 0 \\ 0 & 27 \end{bmatrix} + \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix} + \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 121 & 0 \\ 0 & 121 \end{bmatrix}$$

$$(A^8 + A^6 + A^4 + A^2 + I)V = \begin{bmatrix} 0 \\ 11 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 121 & 0 \\ 0 & 121 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 11 \end{bmatrix} \Rightarrow \begin{bmatrix} 121x \\ 121y \end{bmatrix} = \begin{bmatrix} 0 \\ 11 \end{bmatrix}$$

78.

$$\text{Let } y = \frac{\cos^3 \theta \sin^3 \theta + 1}{(\sin \theta + \cos \theta + 1)^2} = \frac{(\cos \theta + \sin \theta)(1 - \sin \theta \cos \theta) + 1}{(\sin \theta + \cos \theta + 1)^2}$$

$$\text{Let } \cos \theta + \sin \theta = x \quad y = \frac{x \left(1 - \frac{x^2 - 1}{2} \right) + 1}{(x+1)^2} = \frac{2 + 3x - x^3}{2(x+1)^2} = \frac{(2-x)(x+1)^2}{2(x+1)^2} = 1 - \frac{x}{2}$$

$$\therefore y = 1 - \frac{\cos \theta + \sin \theta}{2} \geq 1 - \frac{\sqrt{2}}{2}$$

79.

$$\text{The lines are } \frac{x}{1} = \frac{y}{1} = \frac{z-1}{0}, \frac{x}{1} = \frac{y}{-1} = \frac{z+1}{0}$$

$$\text{Let } C = (\alpha, \beta, \gamma)$$

$$\text{The distance of C from } \vec{r} = \vec{a} + s\vec{b} \text{ is } \frac{|(\vec{c} - \vec{a}) \times \vec{b}|}{|\vec{b}|}$$

$$\text{Distance of C from the first line is } \sqrt{(\gamma-1)^2 + \frac{(\alpha-\beta)^2}{2}} \quad \dots(i)$$



The distance of C from the second line is $\sqrt{(\gamma+1)^2 + \frac{(\alpha+\beta)^2}{2}}$ (ii)

$$(i), (ii) \Rightarrow (\gamma+1)^2 = (\gamma-1)^2 + \frac{(\alpha-\beta)^2}{2} \quad 4\gamma = -2\alpha\beta. \text{ Locus of C is } xy = -2z.$$

80. $f'(x) = 2\sin(x+\alpha)\cos(x+\alpha) + 2\sin(x+\beta)\cos(x+\beta) - 2\cos(\alpha-\beta)\cos(x+\alpha)\sin(x+\beta) - 2\cos(\alpha-\beta)\sin(x+\beta)\cos(x+\beta)$
 $= \sin(2x+2\alpha) + \sin(2x+2\beta) - 2\cos(\alpha-\beta)\sin(2x+\alpha+\beta)$
 $= 2\sin(2x+\alpha+\beta)\cos(\alpha-\beta) - 2\cos(\alpha-\beta)\sin(2x+\alpha+\beta) = 0$
 $\therefore f(x)$ is a constant function.

81. Let equation of plane be $x+2y+3z-7+\lambda x=0$

Since $(-1, 0, 2)$ lies on it $\therefore \lambda = -2$

Equation of plane is $-x+2y+3z-7=0$

$$|\cos\theta| = \frac{6}{7} = \frac{p}{q}$$

82. Plane is $-x+z=1 \Rightarrow d = \frac{3}{\sqrt{2}}$

83. $f(x+y) = f(x)f(y)$
 $f(2) = (f(1))^2, f(3) = (f(1))^3$ and so on
 $\sum_{r=1}^n f(a+r) = f(a)f(1)(1+f(1)+(f(1))^2+\dots+(f(1))^{n-1})$
 $= f(a)\frac{3(3^n-1)}{2}$

84. If $f(\theta) = 0 \Rightarrow \sin\theta = 0$ or $\sin 2\theta = 0$ or $\cos 3\theta = 0$ or $\cos 4\theta = 0$ or $\cos 5\theta = 0$

85. $(t+1)x+8y-4t=0$
 $tx+(t+3)y-(3t-1)=0$
 $\frac{t+1}{t} = \frac{8}{t+3} = \frac{-4t}{-(3t-1)}$
 $t=3$

86. $\lim_{x \rightarrow \frac{\pi}{4}} \frac{4\sqrt{2} - (\cos x + \sin x)^5 - 10\cos x(1 - \sin 2x)}{1 - \sin 2x} = \lim_{x \rightarrow \frac{\pi}{4}} \left(\frac{4\sqrt{2} - \left(\sqrt{2} \cos\left(x - \frac{\pi}{4}\right) \right)^5}{1 - \sin 2x} \right) - \lim_{x \rightarrow \frac{\pi}{4}} 10\cos x$
 $= \lim_{x \rightarrow \frac{\pi}{4}} 4\sqrt{2} \left(\frac{1 - \cos^5\left(x - \frac{\pi}{4}\right)}{1 - \sin 2x} \right) - \frac{10}{\sqrt{2}} = \lim_{h \rightarrow 0} \frac{4\sqrt{2}(1 - \cos^5 h)}{1 - \cos 2h} - 5\sqrt{2}$
 $= \lim_{x \rightarrow 0} \frac{4\sqrt{2}(1 + \cosh + \cos^2 h + \cos^3 h + \cos^4 h)}{2(1 + \cosh)} - 5\sqrt{2} = 5\sqrt{2} - 5\sqrt{2} = 0$

87. Taking dot product with $\bar{a}, \bar{b}, \bar{c}$, we get $5[\bar{a} \bar{b} \bar{c}] = \bar{a} \cdot \bar{a}$, $3[\bar{a} \bar{b} \bar{c}] = \bar{a} \cdot \bar{c}$ and $\bar{a} \cdot \bar{b} = 0$



So. $\frac{(2\bar{a} + \bar{c} + 3\bar{d}) \cdot \bar{a}}{[\bar{a} \bar{b} \bar{c}]} = 13$ as $\bar{d} \cdot \bar{a} = (\lambda_1 \bar{b} + \lambda_2 \bar{a} \times \bar{b}) \cdot \bar{a} = 0$

88. Check the continuity and differentiability at integral points. At $x = 1$, it is not differentiable $f(x) = x \sin \pi(x-1) = -x \sin \pi x$ $x > 1 = x \sin \pi x$

It is continuous but not differentiable

89. Let P and Q be points on the lines, then

$$P(-2+2s, -6+3s, 34-10s), Q(-6+4t, 7-3t, 7-2t)$$

The d.r's of PQ are $(-4+4t-2s, 13-3t-3s, -27-2t+10s)$

PQ is perpendicular to the two lines with d.r's 2, 3, -10 and 4, -3, -2.


$$\therefore 2(-4+4t-2s) + 3(13-3t-3s) - 10(-27-2t+10s) = 0$$

$$4(-4+4t-2s) - 3(13-3t-3s) - 2(-27-2t+10s) = 0$$

$$\Rightarrow 113s - 19t = 301, 29 - 19s = 1$$

Solving $s = 3, t = 2, P = (4, 3, 4) = (a, b, c) \Rightarrow a + b + c = 11$

- 90.



$$\text{Area of base (PQRS)} = \frac{1}{2} |\overrightarrow{PR} \times \overrightarrow{SQ}| = \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & -2 \\ 1 & -3 & -4 \end{vmatrix}$$

$$= \frac{1}{2} |-10\hat{i} + 10\hat{j} - 10\hat{k}| = 5|\hat{i} - \hat{j} + \hat{k}| = 5\sqrt{3}$$

$$\text{Height} = \text{proj. of PT on } \hat{i} - \hat{j} + \hat{k} = \left| \frac{1-2+3}{\sqrt{3}} \right| = \frac{2}{\sqrt{3}}$$

$$\text{volume} = (5\sqrt{3}) \left(\frac{2}{\sqrt{3}} \right) = 10 \text{ cu. Units}$$