



**Sri Chaitanya**  
Educational Institutions

**Infinity**  
Learn



**JEE ADVANCED**



**Sri Chaitanya IIT Academy., India.**

✶ A.P ✶ T.S ✶ KARNATAKA ✶ TAMILNADU ✶ MAHARASTRA ✶ DELHI ✶ RANCHI

*A right Choice for the Real Aspirant*

**ICON Central Office - Madhapur - Hyderabad**

**Sec: Sr.Super60\_STERLING\_BT**

**Paper -2(Adv-2020-P2-Model)**

**Date: 01-10-2023**

**Time: 02.00Pm to 05.00Pm**

**CTA-08**

**Max. Marks: 198**

01-10-2023\_Sr.Super60\_STERLING\_BT\_Jee-Adv(2020-P2)\_CTA-08\_Syllabus

**PHYSICS**

: CTA-8 (RPTA.1 TO RPTA.9 SYLLABUS)

Thermal physics, Geometrical optics, Wave optics, General Physics, Kinematics, NLM, Friction, Circular motion, WPE, COM & Collisions, Rigid body Dynamics 1, Rigid body Dynamics 2, Fluid statics & Dynamics

**CHEMISTRY**

: CTA-8 (RPTA.1 TO RPTA.9 SYLLABUS)

Nomenclature, Isomerism, GOC, Alkanes, Alkene & Alkyne, Benzene, Alkyl Halides & Aryl Halides, Alcohols, Phenols, Ethers & Amines, Aldehydes & Ketones, Carboxylic acid & Derivatives, Biomolecules, Polymers, Chemistry in Everyday Life & POC, 1.Classification of Elements and Periodicity in Properties, 2.Chemical bonding and Molecular Structure, 3.Hydrogen, 1.s-Block Elements, 2.Group-13, 3.Group 14

**MATHEMATICS**

: CTA-8 (RPTA.1 TO RPTA.9 SYLLABUS)

Functions & LCD, Application of Differentiation (AOD), TOTAL DIFFERENTIAL CALCULUS, Indefinite Integration, Definite Integration, Areas & Differential Equations, Total Integral Calculus, Vectors & 3D-Geometry, Matrices & Determinants

**Name of the Student:** \_\_\_\_\_

**H.T. NO:**

--	--	--	--	--	--

**JEE-ADVANCE-2020-P2-Model**

Time: 3:00Hour's

**IMPORTANT INSTRUCTIONS**

Max Marks: 198

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 1 – 6)	Questions With Integer Answer Type	3	-1	6	18
Sec – II(Q.N : 7 – 12)	<b>Questions with Multiple Correct Choice +1 partial marks</b>	4	-2	6	24
Sec – III(Q.N : 13 – 18)	Questions with Numerical Value Answer Type	4	0	6	24
<b>Total</b>				<b>18</b>	<b>66</b>

**CHEMISTRY:**

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 19 – 24)	Questions With Integer Answer Type	3	-1	6	18
Sec – II(Q.N : 25 – 30)	<b>Questions with Multiple Correct Choice +1 partial marks</b>	4	-2	6	24
Sec – III(Q.N : 31 – 36)	Questions with Numerical Value Answer Type	4	0	6	24
<b>Total</b>				<b>18</b>	<b>66</b>

**MATHEMATICS:**

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 37 – 42)	Questions With Integer Answer Type	3	-1	6	18
Sec – II(Q.N : 43 – 48)	<b>Questions with Multiple Correct Choice +1 partial marks</b>	4	-2	6	24
Sec – III(Q.N : 49 – 54)	Questions with Numerical Value Answer Type	4	0	6	24
<b>Total</b>				<b>18</b>	<b>66</b>

Sri Chaitanya Educational Institutions  
INDIA

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 2

Sri Chaitanya  
Educational InstitutionsInfinity  
Learn

**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1  
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

JEE MAIN

RANK

JEE Advanced

RANK

NEET

RANK

SINGARAJU  
VENKAT KUMAR  
RANK 1  
300  
300



VAVILALA  
CHIDVILAS REDDY  
RANK 1  
341  
360



ROSA VARUN  
CHANDRASEKHAR  
RANK 1  
720  
720





## PHYSICS


Max Marks: 66

SECTION-I  
(INTEGER ANSWER TYPE)

- This section contains SIX (06) questions.
- The answer to each question is a SINGLE DIGIT INTEGER ranging from 0 TO 9, BOTH INCLUSIVE.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual Numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated **according to the following marking scheme:**


**Full Marks:** +3 If ONLY the correct integer is entered;**Zero Marks:** 0 If the question is unanswered;**Negative Marks:** -1 In all other cases

01. A side wall of a wide open tank is provided with a narrowing tube through which water flows out. The cross sectional area of the tube decreases from  $S=3.0 \text{ cm}^2$  to  $s=1.0 \text{ cm}^2$ . The water level in the tank is  $h = 4.6 \text{ m}$  higher than in the tube. Neglecting the viscosity of water, find the horizontal component of the force tending to pull the tube out of the tank. (Communicate the nearest integer).



02. A uniform disc of mass  $M$  and radius  $R$  is rotating freely about its stationary smooth central vertical axis with angular speed  $\omega_0$ . Another disc of mass  $m$  and radius  $r$  is free to rotate about a horizontal rod  $AB$ . Length of the rod  $AB$  is  $L$  ( $< R$ ). And its end  $A$  is rigidly attached to the vertical axis of the first disc. The disc of mass  $m$ , initially at rest, is placed gently on the disc of mass  $M$  as shown in the figure. Find the time in seconds after which the slipping between the two discs will cease. Assume that normal reaction between the two disc is equal to  $mg$ , coefficient of friction between the two discs is  $\mu$  given

$$M = 1\text{kg}, m = 1\text{kg}, L = 0.5 \text{ m}, \mu = 0.1, \omega_0 = 5\text{rad/sec}$$



JEE MAIN

2023  
SINGARAJU  
VENKAT KISHORHNYA  
RANK 1  
300

RANK

JEE Advanced

2023  
VAVILALA  
CHIDVILAS REDDY  
RANK 1  
360

RANK

NEET


2023  
KOTA VARUN  
CHANDRANATHAN  
RANK 1  
720

RANK






03. Two identical calorimeters A and B contain equal quantity of water at  $20^{\circ}\text{C}$ . A 5 gm piece of metal X of specific heat  $0.2\text{ cal g}^{-1}(\text{C}^{\circ})^{-1}$  is dropped into A and a 5 gm piece of metal Y into B. The equilibrium temperature in A is  $22^{\circ}\text{C}$  and in B is  $23^{\circ}\text{C}$ . The initial temperature of both the metals is  $40^{\circ}\text{C}$ . The specific heat of metal Y in  $\text{cal g}^{-1}(\text{C}^{\circ})^{-1}$  is  $\frac{27}{17x}$ . Find  $x$ .
04. A uniform U – Tube containing a liquid of uniform density  $\rho$ , rotates about a vertical axis with constant angular velocity  $\omega$ , as shown in the figure.



Length of the horizontal section is  $L$ . Difference in height of the liquid columns in the

two vertical sections of the tube is  $\Delta h = \frac{\omega^2 L^2}{xg}$ . Find  $x$ .

05. A uniform rod of mass  $m$  and  $L$  is rotated uniformly at angular velocity  $\omega$  about an axis passing through its end perpendicular to its length. The net force on the left half of the rod at the instant shown in the diagram is found to be  $\frac{n}{8}mL\omega^2$ . What is the value of  $n$ ?



JEE MAIN

2023

SINGARAJU

VENKAT KUMAR

RANK 1

300

300

RANK

1

JEE Advanced

2023

VAVILALA

CHIDVILAS REDDY

RANK 1

341

360

RANK

1

NEET

2023

RORA VARUN

RANK 1

720

720

RANK

1



06. Let  $S_1$  &  $S_2$  be the two slits in YDSE. If central maxima is observed at  $\rho$  and angle  $S_1\rho S_2 = \theta$  ( $\theta$  is small), the Y- coordinate of  $3^{rd}$  minima assuming the origin at the central maxima is  $\pm \frac{K\lambda}{2\theta}$ . Find 'K' ( $\lambda$ : wave length of mono chromatic light used).

**SECTION – II**  
**(ONE OR MORE CORRECT ANSWER TYPE)**

•This section contains SIX (06) questions.

•Each question has FOUR options. **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).

•For each question, choose the option(s) corresponding to (all) the correct answer(s).

•Answer to each question will be evaluated **according to the following marking scheme**:

**Full Marks:** +4 If only (all) the correct option(s) is(are) chosen; **Partial Marks** +3 If all the four options are correct but ONLY three options are chosen;


**Partial Marks:** +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;

**Partial Marks:** +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;

**Zero Marks:** 0 If none of the options is chosen (i.e. the question is unanswered);

**Negative Marks:** -2 In all other cases.

07. The pitch of screw gauge is 1mm and its circular scale is divided into 100 divisions. When nothing is put between the zero of main scale is not seen, but when circular scale is rotated by  $450^\circ$  the zero of main scale is seen and zero of main scale coincide with the zero of circular scale. When a glass plate is placed between the studs, the circular scale lies between  $18^{th}$  to  $19^{th}$  division of main scale and circular scale reads 34 division. Then,
- A) Error is positive zero error and its magnitude is 1.25 mm.  
B) Error is negative zero error and its magnitude is 1.25 mm  
C) The thickness of the glass plate is 19.59 mm.  
D) The thickness of the glass plate is 17.09 mm.
08. The diagram illustrates a right circular cone-shaped hypothetical mountain, of apex 'O'. If you build a shortest distance track for a sightseeing train around the mountain, in which the track starts at point A and ends at point B, the track will first go uphill, but then it will go downhill.  $AB=10\text{km}$ , radius of cone= $20\text{ km}$ . Slant height of the cone= $60\text{km}$ . The speed of train (consider it a particle) is set at a constant  $20\text{ kmph}$ . Mark the **CORRECT** statements(s).



JEE MAIN

2023

SINGARAJU

VENKAT KUMAR

RANK 1

300

300



RANK

1

JEE Advanced

2023

VAVILALA

CHIDVILAS REDDY

RANK 1

344

360



RANK

1

NEET

2023

ROOJA VARUN

CHANDRANATH

RANK 1

720





720



RANK


1

- A) At the instant of shortest distance from 'O' the angular velocity of train about 'O' is maximum
- B) The journey time of the train is  $\frac{\sqrt{91}}{2}$  hr
- C) The length of the downhill part of the journey is  $\frac{200}{\sqrt{91}}$  km
- D) The velocity of the train is always perpendicular to the line joining 'O' to the train
09. Which of the following is/are not possible equilibrium conditions?




10. Which one of the following statements is (are) FALSE?
- A) Temperature differing by  $25^\circ$  on the Fahrenheit scale must differ by  $45^\circ$  on the Celsius scale (Centigrade scale)
- B) Temperature which differ by  $10^\circ$  on the Celsius scale (Centigrade scale) must differ by  $18^\circ$  on the Fahrenheit scale
- C) Water at  $90^\circ C$  is warmer than water at  $202^\circ F$
- D)  $0^\circ F$  corresponds to  $-32^\circ C$
11. There is an infinite array of ideal springs as shown in figure, whose base row is attached to a fixed support. All the springs are identical and their spring constants are  $k$  each. Initially all the springs are unstretched. Then a constant force  $F$  is applied on the free end of topmost spring along its length. (Assume all the bars to be of negligible mass). Number of springs rows are increasing in GP, from top to bottom. When the system is in equilibrium, choose the CORRECT option(s).





- A) Displacement of free end is  $2F/K$
- B) Displacement of free end is  $F/2K$
- C) Displacement of topmost bar is  $F/K$
- D) Displacement of topmost bar is  $F/4K$

12. In two identical communicating vessel we poured water (see picture). In one of them we put an ice ball of volume  $V = 100\text{cm}^3$  which gets exactly half immersed in the water. The density of water  $\rho_w = 1000 \text{ kg / m}^3$ , the density of ice  $\rho_i = 900 \text{ kg / m}^3$ .



Select the correct statement(s). Soon after placing the ice ball in left vessel.


- A) The volume of water flowing to the right vessel is 25cc.
- B) The volume of water flowing to the right vessel is 50cc.
- C) The ice ball is resting on the bottom of vessel.
- D) The ice ball is floating on the water surface.

### SECTION – III (NUMERICAL VALUE TYPE)

- This section contains **SIX (06)** questions. The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual Numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:  
**Full Marks:** +4 If ONLY the correct numerical value is entered;  
**Zero Marks:** 0 In all other cases..




13.



Maximum force a to move these blocks together is \_\_\_ N if  $g=10\text{m/s}^2$

14. A light ray incident along  $2\vec{i} + 4\vec{j} + \sqrt{5}\vec{k}$  strikes on x-z plane from medium 1 of refractive index  $\sqrt{3}$  and enters in to medium 2 of refractive index  $\mu_2$ . The value of  $\mu_2$  for which the value of angle of refraction becomes  $90^\circ$ , is \_\_\_\_ ( $\sqrt{3} = 1.73$ )
15. A T-shaped system is made by using two identical rods AB and CD each of mass m and length 'l'. A liner impulse J is imparted to the system at midpoint of rod AB. Just after impulse, angular speed of rod AB is (1)  $\omega_1$  if midpoint of rod CD is connected to end B by a pin support so that rod CD can rotate freely about hinge at end B and (2)  $\omega_2$  if midpoint of CD is rigidly connected to end B. Find  $\frac{4\omega_1}{\omega_2}$



16. Tennis balls are usually rejected if they fail to rebound to waist level when dropped from shoulder level. If a ball just passes the test as indicated in the figure, determine the coefficient of restitution  $e$ .



JEE MAIN

2023

SINGARAJU

VENKAT KUMAR

RANK 1

300

300

RANK

1

JEE Advanced

2023

VAVILALA

CHIDVILAS REDDY

RANK 1

341

360

RANK

1

NEET

2023

RORA VARUN

CHANDRAVATHI

RANK 1


720

720


RANK

1





17. A uniform rod has mass  $m=1\text{kg}$  and length  $l=13\text{m}$ . One end of the rod is pulled with a constant velocity of  $v_0=1\text{m/s}$  along a frictionless horizontal floor in the negative  $x$  direction. The other end is moving along a parabolic fixed curve. The equation of the parabola is  $x^2=20y$ . If the angular velocity of the rod (in minutes/s) when the end point 'B' is at  $(10, 5)$  is  $\omega$ . Find  $\frac{\omega}{10}$



18. A sphere of mass  $m$  is moving with a velocity  $4\hat{i} - \hat{j}$  when it hits a smooth wall and rebounds with velocity  $\hat{i} + 3\hat{j}$ . Find the coefficient of restitution between the sphere and the wall.


## CHEMISTRY

Max Marks: 66

SECTION-I  
(INTEGER ANSWER TYPE)

- This section contains SIX (06) questions.
- The answer to each question is a SINGLE DIGIT INTEGER ranging from 0 TO 9, BOTH INCLUSIVE.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual Numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:  
**Full Marks:** +3 If ONLY the correct integer is entered;  
**Zero Marks:** 0 If the question is unanswered;  
**Negative Marks:** -1 In all other cases

19.




(Note: No Hydration across alkyne from A to B)

How many of the following are correct statements regarding above conversion

- A)  $x + y = 9$
- B) B consumes 6 mole NaOH for Acid – Base neutralization
- C) B gives silver mirror test
- D) One mole of B with excess Na liberates 2 moles of  $\text{H}_2$  gas
- E) B gives turbidity immediately with Lucas reagent.
20. How many of the following reactions produce carbonyl compound as one of the products.

A		F	
B		G	
C		H	
D		I	
E		J	

21.



How many of the following mechanisms are as principle mechanisms involved in above conversions; during formation of major product in respective reaction / step


- A) Electrophilic aromatic substitution [EAS]      B) Electrophilic Addition  
C) Free radical substitution      D) Free radical addition  
F) Nucleophile substitution      G) Nucleophilic Addition  
H) Elimination of water      I) dehydro halogenation

22. In how many of the following reactions, the product has inversion in configuration when compared with reactant at reaction center. [In formation of major product.]

A		F	
B		G	
C		H	
D		I	
E			



23.



Pick up correct statements about A to I in above conversions.

- (1) A is  $CH_3 - CH = CH_2$
- (2) B is non terminal (internal) alkyne
- (3) C & D are diastereomers
- (4) E is Racemic mixture of 4-Methyl Hept-2-yn-4-ol
- (5) F is ethoxide
- (6) G decolourises Baeyer's reagent
- (7) H is Aryl Alkyl ether
- (8) There are 4 possible isomeric Alkenes in the mixture I.

24.

Gly-Lys-Pro-Glu- His

Regarding above pentapeptide, chain, pick up the number of correct statements given below.

- A) It is also represented as GKPGH
- B) On complete hydrolysis gives 2 essential amino acids
- C) On complete hydrolysis gives one Acidic Amino acid & Two Basic Amino acid
- D) At one place of chain; Nitrogen atoms are arranged similar to Nitrogen atoms in Guanidine [Guanidyl group of Nitrogen atoms]
- E) Glycine end of chain is called as c-terminal
- F) On complete hydrolysis gives four primary Amino acids & one secondary amino acid.



## SECTION – II

### (ONE OR MORE CORRECT ANSWER TYPE)

• This section contains SIX (06) questions.

• Each question has FOUR options. **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).

• For each question, choose the option(s) corresponding to (all) the correct answer(s).

• Answer to each question will be evaluated **according to the following marking scheme:**

**Full Marks:** +4 If only (all) the correct option(s) is(are) chosen; **Partial Marks** +3 If all the four options are correct but ONLY three options are chosen;


**Partial Marks:** +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;

**Partial Marks:** +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;

**Zero Marks:** 0 If none of the options is chosen (i.e. the question is unanswered);

**Negative Marks:** -2 In all other cases.

25.



Pick up correct Statements about P;Q;X;Y;Z

A) The electrophiles Involved in P to Q and P to Z respectively are  $N=O^+$  &  $:CCl_2^+$


B)  $Q \xrightarrow[\text{(iii) } AC_2O]{\text{(i) } NaOH/CO_2, \text{ (ii) } H_3O^+} \text{Asprin (Major)}$

C)  $P \xrightarrow[H^+]{PhN_2^+ Cl^-} \text{yellow dye}$

$Q \xrightarrow[OH^-]{PhN_2^+ Cl^-} \text{Orange dye}$

D) Both P & Q form white PPT with Bromine water.

26.




Pick up correct statements among following.





- A) Among A, B, D, E & F only 3 compounds give  $\text{Cu}_2\text{O}$  with Fehling's solution  
 B) When A, B, D, E & F are treated separately with  $\text{KOI} / \text{I}_2 + \text{KOH}$  then only 2 compounds form  $\text{CHI}_3$ .  
 C)  $G \xrightarrow{\text{HBr}} X$  Formation of X is nucleophile addition at alkene.  
 D) Cannizzaro & Aldol condensation reactions are involved in above conversion

27.



Pick up correct statements about A ; B; & D



D) B to D Involves PERKIN reaction in one of the steps.

28. The following observations are recorded about 3 isomers of  $\text{C}_3\text{H}_9\text{N}$ .

$\Rightarrow$  Isomer X is prepared by Gabriel phthalimide synthesis with higher rate than other  $1^\circ$  amine isomer Y

$\Rightarrow$  Isomer Z reacts with Hinsberg's reagent and the product formed is insoluble in alkali

$\Rightarrow$  Pick up correct statements about X; Y & Z



JEE MAIN

2023  
 SINGARAJU  
 VENKAT KISHOREN  
 RANK 1  
 300



RANK

JEE Advanced

2023  
 VAVILALA  
 CHIRUVAAS REDDY  
 RANK 1  
 360



RANK


NEET

2023  
 KOTA VARUN  
 CHANDRANATHAN  
 RANK 1  
 720




RANK





B) Y is isopropyl amine


C) Z is N-Methyl Ethanamine



29. Decreasing order of boiling points of 4 isomeric BUTYL ALCOHOLS is  
 $P > Q > R > S$

Pick up correct statements about P, Q, R, S

A) Compound P is 1-Butanol



C) In Lucas test; "P" isomer gives red turbidity immediately

D) Formation of ether by bimolecular reaction is most suitable for isomer P; when P is treated with  $\text{H}_2\text{SO}_4$  with suitable temperature to form ether

30. How many of the following are correct [consider Neutral forms of molecules].

A) Oxime of Acetaldehyde exhibit Geometrical isomerism

B) Semicarbazone of Acetone has 5 lone pairs

C) Cyanohydrine of butanone is chiral molecule

D) Sodium Bisulphite adduct of propanal is chiral molecule


### SECTION – III (NUMERICAL VALUE TYPE)

- This section contains **SIX (06)** questions. The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual Numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:  
**Full Marks:** +4 If ONLY the correct numerical value is entered;  
**Zero Marks:** 0 In all other cases..




31. How many of the following can be oxidized by Tollen's reagent.

A) Glucose; B) Fructose; C) Propene; D) Mannose;




G) But-2-yne H) Ribose I) 2-deoxy Ribose

32. The sum of Bond orders in all the following combinedly is \_\_\_\_ (Rounding to the nearest natural number)



33. 2 Liter 56 volume  $H_2O_2$  can oxidise \_\_\_\_ mole  $KMnO_4$  in  $H_2SO_4$

34. How many of the following ionic hydrides liberate hydrogen gas on addition to water.



35. (i) Number of electron deficient species among following is 'x'



(ii) Number of chemical species that can get partial or complete hydrolyzed among following is y[ in any medium )  $SiF_4$ ;  $SiCl_4$ ;  $BCl_3$ ;  $BF_3$  Determine  $x + y$

36. In long form of periodic table with 7 periods & 18 groups [ Numbered 1 to 18 from left); Calculate sum of group number of metal ions present in Gilman reagent & Grignard's Reagent



**MATHEMATICS****Max Marks: 66****SECTION-I  
(INTEGER ANSWER TYPE)**

- This section contains SIX (06) questions.
- The answer to each question is a SINGLE DIGIT INTEGER ranging from 0 TO 9, BOTH INCLUSIVE.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual Numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated **according to the following marking scheme:**  
**Full Marks:** +3 If ONLY the correct integer is entered;  
**Zero Marks:** 0 If the question is unanswered;  
**Negative Marks:** -1 In all other cases

37. If  $\int \frac{(2-x^2)e^x}{(1-x)\sqrt{1-x^2}} dx = \mu e^x \left( \frac{1+x}{1-x} \right)^\lambda + C$  then  $2(\lambda + \mu)$  is equal to .....

38. If  $\lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{{}^n C_K}{n^K (K+3)} = e - l$ . Then  $l$  is

39. Let S denotes the sum of all the values of  $\lambda$  for which the system of equations

$$(1+\lambda)x_1 + x_2 + x_3 = 1$$

$$x_1 + (1+\lambda)x_2 + x_3 = \lambda$$

$$x_1 + x_2 + (1+\lambda)x_3 = \lambda^2$$

is inconsistent. Find  $|S|$ .

40. If a, b, c are three complex numbers such that  $a^2+b^2+c^2=0$  and

$$\Delta = \begin{vmatrix} b^2 + c^2 & ab & ac \\ ab & c^2 + a^2 & bc \\ ac & bc & a^2 + b^2 \end{vmatrix} = ka^2b^2c^2, \text{ then value of } k \text{ is}$$

41. A circle is inscribed in an n-sided regular polygon  $A_1, A_2, \dots, A_n$  having each side a unit for any arbitrary point P on the circle.

If  $\sum_{i=1}^n (PA_i)^2 = n \frac{a^2}{k} \left( \frac{\lambda + \cos^2 \pi/n}{\sin^2 \pi/n} \right)$ . Then the value of  $\left[ \frac{k+\lambda}{2} \right]$  is (where  $[x]$  is the greatest integer less than equal to x).

42. Let PM be the perpendicular from the point  $P(1, 2, 3)$  to XY-plane. If OP makes an angle  $\theta$  with the positive direction of the Z-axis and OM makes an angle  $\phi$  with the positive direction of X-axis, where O is the origin, then  $3\sqrt{5} \tan \theta + \tan \phi$  is \_\_\_\_\_.



JEE MAIN

2023

SINGARAJU

VENKAT KISHOREN

RANK 1

300

300

RANK

1

JEE Advanced

2023

VAVILALA

CHIDVILAS REDDY

RANK 1

341

360

RANK

1

NEET

2023

NORA VARUN

CHANDRANATHAN

RANK 1

720

720

RANK

1





## SECTION – II

### (ONE OR MORE CORRECT ANSWER TYPE)

• This section contains SIX (06) questions.

• Each question has FOUR options. **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).

• For each question, choose the option(s) corresponding to (all) the correct answer(s).

• Answer to each question will be evaluated **according to the following marking scheme**:

**Full Marks:** +4 If only (all) the correct option(s) is(are) chosen; **Partial Marks** +3 If all the four options are correct but ONLY three options are chosen;

**Partial Marks:** +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;

**Partial Marks:** +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;

**Zero Marks:** 0 If none of the options is chosen (i.e. the question is unanswered);

**Negative Marks:** -2 In all other cases.

43. If  $\int x \log(1+x^2) dx = \phi(x) \log(1+x^2) + x \psi(x) + C$ , then, which of the following option(s) is/are correct?

A)  $\phi(x) = \frac{1+x^2}{2}$     B)  $\psi(1) = \frac{-1}{2}$     C)  $\psi(x) = \frac{-(1+x^2)}{2}$     D)  $\phi(3) = 5$

44. If  $S$  be the area of the region enclosed by  $y = e^{-x^2}$ ,  $y = 0$ ,  $x = 0$  and  $x = 1$ . Then, which of the following option(s) is/are correct?

A)  $S \geq \frac{1}{e}$     B)  $S \geq 1 - \frac{1}{e}$     C)  $S \leq \frac{1}{4} \left( 1 + \frac{1}{\sqrt{e}} \right)$     D)  $S \leq \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{e}} \left( 1 - \frac{1}{\sqrt{2}} \right)$

45. A curve  $y=f(x)$  has the property that the perpendicular distance of the origin from the normal at any point P of the curve is equal to the distance of the point P from the x-axis. Then the differential equation of the curve

- A) is homogeneous  
B) Can be converted into linear differential equation with some suitable substitution  
C) is the family of circles touching the x-axis at the origin  
D) The family of circles touching the y-axis at the origin

46. If  $2f(x) + xf\left(\frac{1}{x}\right) - 2f\left(\sqrt{2} \sin \pi \left(x + \frac{1}{4}\right)\right) = 4 \cos^2\left(\frac{\pi x}{2}\right) + x \cos\left(\frac{\pi}{x}\right)$ ,  $\forall x \in \mathbb{R} - \{0\}$ , which of the following statement(s) is/are true?

A)  $f(2) + f\left(\frac{1}{2}\right) = 1$     B)  $f(2) + f(1) = 0$   
C)  $f(2) + f(1) = f\left(\frac{1}{2}\right)$     D)  $f(1) \cdot f\left(\frac{1}{2}\right) \cdot f(2) = 1$

47.  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  are three vectors such that  $\mathbf{a} \cdot \mathbf{a} = \mathbf{b} \cdot \mathbf{b} = \mathbf{c} \cdot \mathbf{c} = 3$  and  $|\mathbf{a} - \mathbf{b}|^2 + |\mathbf{b} - \mathbf{c}|^2 + |\mathbf{c} - \mathbf{a}|^2 = 27$ , then which of the following option(s) is/are correct?
- A)  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  are necessarily coplanar.  
B)  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  represent sides of a triangle in magnitude and direction



JEE MAIN

2023

SINGARAJU

VENKAT KUMAR

RANK 1

300

300

RANK

1

2023

JEE ADVANCED

2023

360

360

RANK

1

2023

NEET

2023

720

720

RANK

1

2023

NEET

2023

720

720



C)  $\mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{c} + \mathbf{c} \cdot \mathbf{a}$  has the least value  $-9/2$

D)  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  represent orthogonal triad of vectors

48. If  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$  and  $\mathbf{d}$  are the unit vectors such that  $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = 1$  and  $\mathbf{a} \cdot \mathbf{c} = \frac{1}{2}$ , then which of the following option(s) is/are correct?

A)  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$  are non-coplanar

B)  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{d}$  are non-coplanar

C)  $\mathbf{b}$ ,  $\mathbf{d}$  are non-parallel

D)  $\mathbf{a}$ ,  $\mathbf{d}$  are parallel and  $\mathbf{b}$ ,  $\mathbf{c}$  are parallel

### SECTION – III (NUMERICAL VALUE TYPE)

• This section contains **SIX (06)** questions. The answer to each question is a **NUMERICAL VALUE**.

• For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual

Numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.

• Answer to each question will be evaluated according to the following marking scheme:

**Full Marks:** +4 If ONLY the correct numerical value is entered;

**Zero Marks:** 0 In all other cases..

49. If  $I = \int_0^{\sqrt{3}} \frac{1}{1+x^2} \cdot \sin^{-1}\left(\frac{2x}{1+x^2}\right) dx$ . Then  $\frac{72I}{\pi^2}$  is \_\_\_\_\_

50. For positive integers  $K=1, 2, 3, \dots, n$ . Let  $S_K$  denotes the area of  $\Delta OAB_K$  (where 'O' is the origin) such that  $\angle AOB_K = \frac{K\pi}{2n}$ ,  $OA=1$  and  $OB_K = K$ . The value of the

$$\lim_{n \rightarrow \infty} \frac{\pi^2}{n^2} \cdot \sum_{K=1}^n \frac{S_K}{10} \text{ is}$$

51. Let  $f: R \rightarrow R$  is a function which satisfies condition  $f(x+y^3) = f(x) + [f(y)]^3$  for all  $x, y \in R$ . If  $f'(0) > 0$ , then  $\frac{1}{1000} f(100)$  is \_\_\_\_\_

52. Let  $f(x) = \begin{vmatrix} \sec x & \cos x & \sec^2 x + \cot x \operatorname{cosec} x \\ \cos^2 x & \cos^2 x & \operatorname{cosec}^2 x \\ 1 & \cos^2 x & \operatorname{cosec}^2 x \end{vmatrix}$  and  $I = \int_{\pi/4}^{\pi/2} f(x) dx$ . The value of

$$I + \frac{\pi}{8} + \frac{1}{6\sqrt{2}} \text{ is } \_\_\_\_\_\_$$

53. Let  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  be vectors with magnitudes 3, 4 and 5, respectively and  $\mathbf{a} + \mathbf{b} + \mathbf{c} = 0$ , then the values of  $|\mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{c} + \mathbf{c} \cdot \mathbf{a}|$  is

54. The distance between the plane  $Ax - 2y + z = d$  and the plane containing the lines

$$\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4} \text{ and } \frac{x-2}{3} = \frac{y-3}{4} = \frac{z-4}{5} \text{ is } \sqrt{6}, \text{ then } |d| \text{ is equal to } \dots$$



JEE MAIN

2023

SINGARAJU

VENKAT KUMAR

RANK 1

300

300

RANK

JEE Advanced

2023

VAVILALA

CHIDVILAS REDDY

RANK 1

344

360

RANK

NEET

2023

NORA VARUN

CHANDRANATH

RANK 1

720

720

RANK





# Sri Chaitanya

Junior Colleges & Coaching Centres

Infinity  
Learn BY SRI CHAITANYA



**VAVILALA CHIDVILAS REDDY**  
HT.No. 236165088  
Sri Chaitanya  
6<sup>th</sup> - 12<sup>th</sup> Class

**RANK**

WITH ALL INDIA **RANK 1** IN JEE ADVANCED 2023

# SRI CHAITANYA

STANDS AT THE TOP

SEIZES 5 RANKS IN TOP 10 IN ALL-INDIA OPEN CATEGORY

ANDHRA PRADESH STATE TOPPER



**RAMESH SURYA THEJA**  
HT.No. 236131971

8<sup>th</sup> - 12<sup>th</sup> Class  
Sri Chaitanya



**RANK**



**KALRA RISHI**  
HT.No. 237047209

AITS Student



**RANK**



**RAGHAV GOYAL**  
HT.No. 237020329

9<sup>th</sup> - 12<sup>th</sup> Class  
Sri Chaitanya



**RANK**



**BIKKINA ABHINAV CHOWDARY**  
HT.No. 236163128

4<sup>th</sup> - 12<sup>th</sup> Class  
Sri Chaitanya



**RANK**

32 TOP RANKS BELOW 100 IN ALL-INDIA OPEN CATEGORY

<b>RANK 12</b> DASARI SAKETH NAIDU HT.No. 236060448	<b>RANK 14</b> GUTHIKONDA ABHIRAM HT.No. 236120121	<b>RANK 17</b> P GNANA KOUSIK REDDY HT.No. 236133520	<b>RANK 18</b> D VENKATA YUGESH HT.No. 236046359	<b>RANK 19</b> MOULIK JINDAL HT.No. 237021004	<b>RANK 22</b> D PRATAP SINGH HT.No. 232006139	<b>RANK 23</b> SHAIK SAHIL CHANDA HT.No. 236123370	<b>RANK 26</b> SONI MAYANK HT.No. 232097025	<b>RANK 33</b> GOYAL NAMAN HT.No. 232029345
<b>RANK 37</b> SUMEDH SS HT.No. 231229088	<b>RANK 38</b> V DEERAJ SATHVIK REDDY HT.No. 236067191	<b>RANK 44</b> V VERGIYA KAUSHAL HT.No. 231004058	<b>RANK 46</b> SUTHAR HARSHAL HT.No. 231005060	<b>RANK 58</b> ABHI JAIN HT.No. 232091036	<b>RANK 61</b> RIDAM JAIN HT.No. 232046021	<b>RANK 62</b> SUTHAR HARSH HT.No. 232100028	<b>RANK 63</b> SOJITRA DIPEN HT.No. 231071001	<b>RANK 65</b> D PANEENDRANADHA REDDY HT.No. 236060288
<b>RANK 67</b> GUNDA SUSHANTH HT.No. 231219732	<b>RANK 68</b> HARSH TAYA HT.No. 237021176	<b>RANK 69</b> KRISH GUPTA HT.No. 232066189	<b>RANK 73</b> KUNCHAKURTHY VIVEK HT.No. 236134093	<b>RANK 81</b> N DHARMA TEJA REDDY HT.No. 236060138	<b>RANK 84</b> S VENKAT KOUNDINYA HT.No. 236130679	<b>RANK 87</b> MRUNAL SHRIKANT HT.No. 237044138	<b>RANK 92</b> SAMOTA APURVA HT.No. 232099210	<b>RANK 97</b> N PRINCE BRANHAM REDDY HT.No. 236064112

BELOW 20  
All India Open  
Category Ranks

**10**  
RANKS (50%)

BELOW 100  
All India Open  
Category Ranks

**32**

BELOW 1000  
All India Open  
Category Ranks

**181**

BELOW 100  
All India  
Category Ranks  
Count

**89**

BELOW 1000  
All India  
Category Ranks  
Count

**699**

NUMBER OF RANKS QUALIFIED  
**3,621**

ADMISSIONS OPEN » JEE ADVANCED LONG-TERM 2024

040 66 06 06 06



Scan QR Code for  
JEE Advanced Offers

www.srichaitanya.net