



Sri Chaitanya IIT Academy.,India.

✪ A.P ✪ T.S ✪ KARNATAKA ✪ TAMILNADU ✪ MAHARASTRA ✪ DELHI ✪ RANCHI

A right Choice for the Real Aspirant

ICON Central Office - Madhapur - Hyderabad

Sec: **Sr.Super60_NUCLEUS & STERLING_BT**

Paper -1(Adv-2021-P1-Model)

Date: 03-09-2023

Time: 09.00Am to 12.00Pm

RPTA-05

Max. Marks: 180

03-09-2023_Sr.Super60_NUCLEUS&STERLING_BT_Jee-Adv(2021-P1)_RPTA-05_Syllabus

PHYSICS

: NLM, Friction, Circular motion, WPE Newton's laws of motion; Inertial and uniformly accelerated frames of reference; Static and dynamic friction; Kinetic and potential energy; Work and power, Uniform circular motion, Conservation of mechanical energy

CHEMISTRY

: Alcohols, Phenols , Ethers & Amines:

Alcohols : Physical properties; Reactions: esterification, dehydration (formation of alkenes and ethers); Reactions with: sodium, phosphorus halides, $ZnCl_2$ /concentrated HCl, thionyl chloride; Conversion of alcohols into aldehydes, ketones and carboxylic acids

Phenols: Physical properties; Preparation, Electrophilic substitution reactions of phenol (halogenation, nitration, sulphonation); Reimer-Tiemann reaction, Kolbe reaction; Esterification; Etherification; Aspirin synthesis; Oxidation and reduction reactions of phenol.

Ethers : Preparation, Properties & Reactions

Amines: Basicity of substituted anilines and aliphatic amines, Preparation from nitro compounds, nitriles and amides; Reactions: Hoffmann bromamide degradation, Gabriel phthalimide synthesis; Reaction with nitrous acid, Azo coupling reaction of diazonium salts of aromatic amines; Sandmeyer and related reactions of diazonium salts; Carbylamine reaction, Hinsberg test, Alkylation and acylation reaction

MATHEMATICS

: Definite Integration

Name of the Student: _____

H.T. NO:

--	--	--	--	--	--

**JEE-ADVANCE-2021-P1-Model**

Time:3Hr's

IMPORTANT INSTRUCTIONS

Max Marks: 180

PHYSICS:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 1 – 4)	Questions with Single Correct Choice	+3	-1	4	12
Sec – II(Q.N : 5 – 10)	Paragraph Questions with Numerical Value Answer Type	+2	0	6	12
Sec – III(Q.N : 11 – 16)	Questions with Multiple Correct Choice with partial mark	+4	-2	6	24
Sec – IV(Q.N : 17 – 19)	Questions with Non-negative Integer Value Type	+4	0	3	12
Total				19	60

CHEMISTRY:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 20 – 23)	Questions with Single Correct Choice	+3	-1	4	12
Sec – II(Q.N : 24 – 29)	Paragraph Questions with Numerical Value Answer Type	+2	0	6	12
Sec – III(Q.N : 30 – 35)	Questions with Multiple Correct Choice with partial mark	+4	-2	6	24
Sec – IV(Q.N : 36– 38)	Questions with Non-negative Integer Value Type	+4	0	3	12
Total				19	60

MATHEMATICS:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 39 – 42)	Questions with Single Correct Choice	+3	-1	4	12
Sec – II(Q.N : 43 – 48)	Paragraph Questions with Numerical Value Answer Type	+2	0	6	12
Sec – III(Q.N : 49 – 54)	Questions with Multiple Correct Choice with partial mark	+4	-2	6	24
Sec – IV(Q.N : 55 – 57)	Questions with Non-negative Integer Value Type	+4	0	3	12
Total				19	60

Sec: Sr.Super60_ **NUCLEUS & STERLING_BT**

Space for rough work

Page 2

**Sri Chaitanya**
Educational Institutions**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1**
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN 2023
SINGARAJU
VENKAT KOUNDINYA
RANK 1
300
MARKS



RANK 1

JEE Advanced 2023
VAVILALA
CHIDVILAS REDDY
RANK 1
341
MARKS



RANK 1

NEET 2023
BORA VARUN
CHAKRAVARTHI
RANK 1
720
MARKS



RANK 1



PHYSICS

Max Marks: 60


SECTION – I
(SINGLE CORRECT ANSWER TYPE)

This section contains 4 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONLY ONE option can be correct.

Marking scheme: +3 for correct answer, 0 if not attempted and –1 in all other cases. Section 1 (Max Marks: 12)

- Section 1 contains Four questions
- Each Question has Four Options and Only One of these four will be the correct answer.
- For each question, choose the option corresponding to the correct answer
- The Marking scheme to evaluate Answer to each question will be :
- Full Marks: **+3** (If the answer is correct)
- Zero Marks: **0** (If the question is unanswered)
- Negative Marks: **-1** (In all other cases)


1. Block A of mass M is placed on an inclined plane, connected to a string, passing over a pulley as shown in the fig. The other end of the string also carries a block B of mass M . The system is held in the position shown such that triangle APQ lies in a vertical plane with horizontal line AQ in the plane of the inclined surface.



Find the minimum coefficient of friction between the inclined surface and block A such that the system remains at rest after it is released. Take $\theta = \alpha = 45^\circ$


- A) $\frac{\sqrt{3+2\sqrt{2}}}{\sqrt{2}-1}$ B) $\frac{\sqrt{3-2\sqrt{2}}}{\sqrt{2}+1}$ C) $\frac{\sqrt{5-2\sqrt{2}}}{\sqrt{2}-1}$ D) $\frac{\sqrt{5-2\sqrt{2}}}{\sqrt{2}+1}$
2. Three small discs are connected with two identical massless rods as shown in fig. the rods are pinned to the discs such that angle between them can change freely. The system is placed on a smooth horizontal surface with discs A and B touching a smooth wall and the angle ACB being 90° . A force F is applied to the disc C in a direction perpendicular to the wall. Find acceleration of disc B immediately after the force starts to act. Masses of discs are $m_A = m; m_B = 2m; m_C = m$
- [Wall is perpendicular to the plane of the fig.]





- A) $\frac{F}{m}$ B) $\frac{F}{3m}$ C) $\frac{F}{5m}$ D) $\frac{F}{7m}$


3. A small disc P is placed on an inclined plane forming an angle θ with the horizontal and imparted an initial velocity v_0 . Find how the velocity of disc depends on the angle ϕ which its velocity vector makes with x axis (see figure). The coefficient of friction is $\mu = \tan \theta$ and $\phi_{\text{initial}} = \frac{\pi}{2}$.



- A) $\frac{v_0}{1 + \cos \phi}$ B) $\frac{v_0}{1 + \sin \phi}$ C) $\frac{2v_0}{1 + \sin \phi}$ D) $\frac{2v_0}{1 + \cos \phi}$

4. A small object is sliding on a smooth horizontal floor along a vertical wall. The wall makes a smooth turn by angle θ_0 . Coefficient of friction between the wall and the block is μ . Speed of the object before the turn is u. find its speed (v) just after completing the turn. [The turn is smooth and there are no sharp corners]





- A) $ue^{-\mu\theta_0}$ B) $ue^{\frac{-\mu\theta_0}{2}}$ C) $ue^{-2\mu\theta_0}$ D) $ue^{-\mu(\ln 2)(\theta_0)}$


SECTION 2

- This section contains **THREE (03)** questions stems.
- There are **TWO (02)** questions corresponding to each question stem.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:
- **Full Marks: +2** If ONLY the correct numerical value is entered at the designated place;
- **Zero Marks:0** in all other cases

Question Stem for Question Nos. 5 and 6

Question Stem

A smooth cylinder is fixed with its axis horizontal. Radius of the cylinder is R . A uniform rope (ACB) of linear mass density λ (kg/m) is exactly of length πR and is held in semi-circular shape in vertical plane around the cylinders as shown in the figure. Two massless strings are connected at the ends of the rope and are pulled up vertically with 2 forces T_0 each to keep the rope in contact with cylinder. Take $g = 10 \text{ m/s}^2$






5. Find minimum value of T_0 so that rope does not lose contact with the cylinder at any point and express it as $k\lambda gR$. What is the value of k ?
6. If required T_0 is applied, then Find Tension in the rope at 30° measured anticlockwise from the location where there is maximum chance of string losing contact with the cylinder. Express it as $c\lambda gR$. Find c ?

Question Stem for Question Nos. 7 and 8

Question Stem

Three identical masses are attached to the ends of light strings, the other ends of which are connected together as shown in the figure. Each of the three strings has a length of 3m. The three masses are dropped through three holes (A, B, C) in a table and the system is allowed to reach equilibrium. Each mass is 3 kg. Take reference for potential energy at table level.



7. Locate a point K inside the $\triangle ABC$ such that $AK + BK + CK$ is minimum. If it is expressed as $\frac{1}{2} \left(1 + \sqrt{\frac{p}{q}} \right)$. Find p/q
8. Calculate minimum potential energy of system under equilibrium conditions. If it is expressed as $-255 \left(1 - \sqrt{\frac{r}{s}} \right)$. find s/r






Question Stem for Question Nos. 9 and 10

Question Stem

Mass(m) is released from top of fixed inclined plane having coefficient of friction μ from A to B as shown in figure.. Horizontal surface has coefficient 2μ from B to C. When block reaches C, it entangles in the ideal massless string of length L without loss of any energy and starts circular motion. At a later instant, motion ceases to be circular.

Given $\mu = \frac{1}{64}$. Assume all corners to be gradual enough so that there are no collisions and

block remains in contact with the incline surface from A to B. (Diagram not to scale)



9. The magnitude of the change in its velocity from C to the position, where the string is horizontal is $(K)\sqrt{gL}$. Find the K?
10. When the block leaves the circular path, its velocity vector is $\sqrt{\frac{gL}{8}}(-p\hat{i} + q\hat{j})$. Find the $p + q$

SECTION 3

- This section contains **SIX (06)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:
- Full Marks: +4** If only (all) the correct option(s) is (are) chosen;
- Partial Marks: +3** If all the four options are correct but **ONLY** three options are chosen,
- Partial Marks: +2** If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
- Partial Marks: +1** If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
- Zero Marks: 0** If unanswered;
- Negative Marks: -2** In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to the correct answer, then
 - Choosing **ONLY** (A), (B) and (D) will get +4 marks;
 - Choosing **ONLY** (A), will get +1 mark;
 - Choosing **ONLY** (B), will get +1 mark;
 - Choosing **ONLY** (D), will get +1 mark;
 - Choosing no option(s) (i.e. the question is unanswered) will get 0 marks and
 - Choosing any other option(s) will get -2 marks.





11. A balloon starts rising from the earth's surface. The ascension rate is constant and equal to v_0 . Due to the wind, the balloon gathers a horizontal velocity component $v_x = ky$, where k is constant and y is the height of ascent. Find the dependence of the following quantities on y .


A) Tangential acceleration at any y is $\frac{k^2 y v_0}{\sqrt{v_0^2 + k^2 y^2}}$

B) Centripetal acceleration at any y is $\frac{k v_0}{\sqrt{1 + \left(\frac{ky}{v_0}\right)^2}}$

C) Radius of curvature at an y is $R_C = \frac{v_0}{k} \left(1 + \left[\frac{ky}{v_0}\right]^2\right)^{3/2}$

D) The curvature of path decreases as the balloon rises.

12. A uniform rope of length l is held motionless on a frictionless hemisphere of radius r with one end of the rope at the top of the hemisphere. The hemisphere is made immobile by gluing it on a horizontal floor. The rope is now released. Immediately after the rope is released



A) Only the weight of corresponding vertical projection of string length is effectively responsible to cause the acceleration of rope.

B) The angular position measured from horizontal where maximum tensile force is

developed is $\cos^{-1}\left(\frac{r}{l}\left(1 - \cos\frac{l}{r}\right)\right)$






- C) If just sufficient uniform coefficient of friction is present between rope and hemisphere to prevent sliding of rope, then tension developed in rope at all points is zero.
- D) Minimum coefficient of friction required between rope & hemisphere to prevent slipping of rope is $\tan\left(\frac{\ell}{2r}\right)$. Ignore any component of Rope Tension in radial direction.

13. The incorrect options in the following is/are:

- A) Work done by kinetic friction is always negative.
- B) When an athlete runs with acceleration, the gain in kinetic energy of the athlete is due to work done by frictional force between his shoe and ground. Here work done by friction is positive.
- C) Total Work done by internal tension force in any system is always zero.
- D) In the elongation of spring, Work done by spring is negative irrespective of the reference frame.

14. A hemispherical bowl is placed in inverted position on frictionless horizontal floor and a small disk is placed on the top of the bowl as shown in the figure. The disk and the bowl are of equal mass and there is no friction between them. The disk is slightly pushed horizontally so that it starts sliding on the bowl with negligible speed.




- A) Momentum of hemisphere + disc system will remain conserved until they get separated
- B) At the time of separation, velocity of disc and hemisphere are same.
- C) At time of separation, the angle that disc makes with vertical is $\cos^{-1}(\sqrt{3}-1)$
- D) At time of separation, Pseudo force on disc in frame of centre of mass of hemisphere will be zero but Pseudo force on hemisphere in frame of disc is non-zero






15. The spring is compressed by a distance 'a' and released. The block again comes to rest when the spring is elongated by a distance 'b' from natural length. From the point of maximum compression to maximum elongation, Find



- A) Work done by the spring on the block $= \frac{1}{2}k(a+b)^2$
 B) Work done by the spring on the block $= \frac{1}{2}k(a^2 - b^2)$
 C) Coefficient of friction $= \frac{k(a-b)}{2mg}$
 D) Coefficient of friction $= \frac{k(a+b)}{2mg}$
16. All surfaces are rough and have same μ . Pulleys & strings are ideal & massless and smooth. In both cases block move together & have same acceleration i.e., $a_1 = a_2$ & same velocity i.e. $v_1 = v_2$. Man A, B, C, D remain stationary. A & C manage to keep pulley P_1 & P_2 at rest.






- A) Power supplied by man D > power by man B
 B) Friction force acting on man C from ground > friction force acting on man A from ground
 C) Friction below feet of man B does more work than friction below feet of man D
 D) Work done by static friction on M with respect to ground is positive.

SECTION 4


- This section contains **THREE (03)** questions.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer by using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:
- **Full Marks** : +4 If ONLY the correct integer is entered;
- **Zero Marks** : 0 In all other cases.

17. Three identical smooth cylinders, each of mass m and radius r are resting in equilibrium within a fixed smooth cylinder of radius R (only a part of this cylinder has been shown in the figure). Find the largest value of R in terms of r , for the small cylinders to remain in equilibrium. If R is expressed as $R = r(1 + 2\sqrt{K})$, what is the value of K ?




18. A rubber band has mass m and force constant k . When relaxed, the band forms a ring of radius r . The band is placed horizontally on a vertical frictionless cone as shown (angle 2θ is given). The radius R of the ring formed by the band is $R = r + \frac{mg}{\alpha\pi^2k} \cot \theta$. Find the value of α





19. A simple pendulum of length ℓ is suspended by a nail on a vertical wall at point O. Another nail is to be fixed on the wall such that if the pendulum is released from its initial horizontal position, it just completes a vertical circle round that nail as shown. The locus of the point where the second nail can be fixed is $k(x^2 + y^2) = (3\ell - 2y)^2$. Find the value of k .





CHEMISTRY

Max. Marks: 60

SECTION 1

- This section contains **Four (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:
- Full Marks : +3 If ONLY the correct option is chosen;
- Zero Marks : 0 If the none of the options is chosen (i.e. the question is unanswered);
- Negative Marks : -1 In all other cases.


20. I) benzene 1,2-diol II) benzene 1,3-diol

III) benzene 1,4-diol IV) Phenol


The increasing order of boiling points of above mentioned Phenol(s) is

A) I<II<III<IV B) I<II<IV<III C) IV<I<II<III D) IV<II<I<III

21.




22. Which of the given compound is the most reactive towards dehydration reaction?





23. Which of the following is Most basic in nature?



SECTION 2

- This section contains **THREE (03)** questions stems.
- There are **TWO (02)** questions corresponding to each question stem.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:
- **Full Marks** : +2 If ONLY the correct numerical value is entered at the designated place;
- **Zero Marks** : 0 In all other cases.

Question Stem for Question Nos. 24 and 25

Question Stem

Nitrous acid reacts with all classes of amines. The nature of products obtained from these reactions depends on whether the amine is primary, secondary or tertiary. It also depends on whether the amine is aliphatic or aromatic. Aliphatic primary amines react with nitrous acid ($\text{NaNO}_2 + \text{HCl}$) to form alcohol as the major product. In addition to alcohol, alkene and alkyl halide are also formed as minor products. Certain cyclic primary amines undergo either ring expansion or ring contraction on treatment with nitrous acid. This reaction is called Demjanov ring expansion or contraction.

24.



What is the molecular weight in gm/mol

25. 14.75 g of n-propyl amine reacts with HNO_2 to give product (B) what amount of product is obtained in grams ?





Question Stem for Question Nos. 26 and 27

Question Stem:


3-ethyl -2-Pentene(X) under go reaction with $Hg(OAc)_2 / H_2O$ followed by $NaBH_4$ gives product (Y); but when (X) is reacted with Br_2 / H_2O gives product (Z) and (X) with HBr / H_2O_2 gives product (P).

26. How many sigma (σ) bonds are present in product (P)?

27. How many rearrangement steps involved during the product (Y) formation?

Question Stem for Question Nos. 28 and 29

Question Stem:



28. What is the value of D.B.E (double bound equivalent) of (A)?





29. How many Lone pairs are present in product (B)?

SECTION 3

- This section contains **SIX (06)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:
- Full Marks: +4** If only (all) the correct option(s) is (are) chosen;
- Partial Marks: +3** If all the four options are correct but **ONLY** three options are chosen,
- Partial Marks: +2** If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
- Partial Marks: +1** If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
- Zero Marks: 0** If unanswered;
- Negative Marks: -2** In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to the correct answer, then
 Choosing ONLY (A), (B) and (D) will get +4 marks;
 Choosing ONLY (A), will get +1 mark;
 Choosing ONLY (B), will get +1 mark;
 Choosing ONLY (D), will get +1 mark;
 Choosing no option(s) (i.e. the question is unanswered) will get 0 marks and
 Choosing any other option(s) will get -2 marks.

30. Which of the following reaction do Not give phenol major product ?






31. Ethanol on reaction with $(CH_3CO)_2O$ gives

- A) $CH_3 - COOH$ B) $CH_3COOC_2H_5$
C) $C_2H_5 - O - \underset{\underset{O}{||}}{C} - OH$ D) $HCOOH$

32.




Identify X, Y, Z

- A) Y is $CH_3 - \underset{\underset{O-CH_3}{|}}{CH} - CH_2C \equiv CCH_3$
B) Y is $CH_3 - CH_2 - \underset{\underset{OCH_3}{|}}{CH}C \equiv CCH_3$
C) Z is $CH_3 - \underset{\underset{O-CH_3}{|}}{CH} - CH_2 - CH = CH - CH_3$
D) X is $CH_3 - \underset{\underset{O^-}{|}}{CH}CH_2C \equiv CCH_3$



33. Which of the following reagent(s) may be used to distinguish between phenol and $\text{CH}_3\text{CH}_2\text{OH}$.
- A) Aq. NaOH B) Neutral FeCl_3 C) Tollen's reagent D) NaHCO_3
34. A positive carbylamine test is given by
- A) N, N-dimethylaniline B) 2,4- dimethylaniline
C) 2,N-dimethylaniline D) p-methylbenzylamine
35. In the following reaction, the product S is




SECTION 4

- This section contains **THREE (03)** question.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer the using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:
- **Full Marks : +4** If ONLY the correct integer is entered;
- **Zero Marks : 0** In all other cases.






36.




If reagent used in step-3 is $\text{CH}_3\text{ONa} / \text{CH}_3\text{OH}$ So, $\text{S}_\text{N} / \text{HCl}$ Should be used at which step?

37.



On which carbon bromine will be present in the major product (A)

38. Number of Hydroxyl group(s) present in the product Q is ____





MATHEMATICS

Max. Marks: 60

SECTION 1

- This section contains **Four (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:
- Full Marks** : +3 If **ONLY** the correct option is chosen;
- Zero Marks** : 0 If the none of the options is chosen (i.e. the question is unanswered);
- Negative Marks** : -1 In all other cases.

39. $\int_0^4 \left\{ \frac{x}{4} \right\} \left(1 + \left[\tan \left(\frac{\{x\}}{1 + \{x\}} \right) \right] \right) dx =$ (The greatest integer function is denoted by $[x]$)

where as the fractional part function is denoted by $\{x\}$.

- A) 0 B) 1 C) 2 D) 4

40. $\int_0^2 e^x (x^4 + 8x^3 + 18x^2 + 16x + 5) dx$

- A) $81e^2 - 1$ B) $81e^2 - 8$ C) $81e^2 - 27$ D) $81e^2 + 1$

41. $\int_{-3/2}^{-1/2} (x^5 + 5x^4 + 10x^3 + 8x^2 + x) dx$

- A) $\frac{5}{6}$ B) $\frac{4}{5}$ C) $\frac{6}{7}$ D) $\frac{7}{8}$

42. $\int_2^{12} \frac{\sqrt{x + \sqrt{x + \sqrt{x + \dots \infty}}}}{\sqrt{x} \sqrt{x} \sqrt{x} \dots \infty} dx$

- A) $\ln 6 + \int_3^7 \frac{x^2}{x^2 - 1} dx$ B) $\frac{1}{2} \ln 6 + \int_3^7 \frac{x^2}{x^2 - 1} dx$
- C) $\frac{1}{2} \ln 6 + \int_3^7 \frac{x^2}{x^2 + 1} dx$ D) $\frac{1}{2} \ln 6 + \int_3^7 \frac{x}{x^2 + 1} dx$

SECTION 2

- This section contains **THREE (03)** questions stems.
- There are **TWO (02)** questions corresponding to each question stem.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:
- Full Marks**: +2 If **ONLY** the correct numerical value is entered at the designated place;
- Zero Marks**: 0 In all other cases.

Sec: Sr.Super60_ NUCLEUS & STERLING_BT

Space for rough work

Page 19

Sri Chaitanya
Educational Institutions

THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN
2023
SINGARAJU
VENKAT KOUNDINYA
RANK 1
300
300
MARKS



RANK
1

JEE Advanced
2023
VAVILALA
CHIDVILAS REDDY
RANK 1
341
360
MARKS



RANK
1

NEET
2023
BORA VARUN
CHAKRAVARTHI
RANK 1
720
720
MARKS



RANK
1

**Question Stem for Question Nos. 43 and 44****Question Stem:**

Given that $\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi} = 1.77$

43. $\int_{-\infty}^{\infty} e^{4-x^2 - \frac{4}{x^2}} dx$

44. $\lim_{n \rightarrow \infty} \sqrt{n} \cdot \int_{-1/2}^{1/2} (1 - 4x^2 + x^4)^n dx$

Question Stem for Question Nos. 45 and 46**Question Stem:**

Use coefficient of t^r in the expansion $\int_0^1 ((1-x) + tx)^n dx$, where $r \leq n$

and $\binom{x}{k} = \frac{x(x-1)(x-2)\dots(x-k+1)}{k!}$

45. $100 \int_0^1 \binom{207}{7} x^{200} (1-x)^7 dx$

46. $\int_0^{2024} \binom{x}{2025} dx$

Question Stem for Question Nos. 47 and 48**Question Stem:**

Given that $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} = 1.64$

47. $\int_0^1 x \left[\frac{1}{x} \right] dx$ (The greatest integer function is denoted by $[x]$)

48. $\int_0^1 \frac{(x+1) \log(x)}{x^3 - 1} dx$

**JEE MAIN****2023**
SINGARAJU
VENKAT KOUNDINYA
RANK 1
300
300
MARKS**RANK****1****JEE Advanced****2023**
VAVILALA
CHIDVILAS REDDY
RANK 1
341
360
MARKS**RANK****1****NEET****2023**
BORA VARUN
CHAKRAVARTHI
RANK 1
720
720
MARKS**RANK****1**



SECTION 3

- This section contains **SIX (06)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:
- **Full Marks** : +4 If only (all) the correct option(s) is (are) chosen;
- **Partial Marks** : +3 If all the four options are correct but **ONLY** three options are chosen,
- **Partial Marks** : +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
- **Partial Marks** : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
- **Zero Marks** : 0 If unanswered;
- **Negative Marks**: -2 In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to the correct answer, then
 Choosing ONLY (A), (B) and (D) will get +4 marks;
 Choosing ONLY (A), will get +1 mark;
 Choosing ONLY (B), will get +1 mark;
 Choosing ONLY (D), will get +1 mark;
 Choosing no option(s) (i.e. the question is unanswered) will get 0 marks and
 Choosing any other option(s) will get -2 marks.

49. $\int_0^{\frac{\pi}{2}} x \tan 2x \ln(\tan x) dx =$

A) $\frac{\pi}{2} \int_0^{\frac{\pi}{2}} \tan 2x \ln(\tan x) dx$

B) $\frac{\pi}{4} \int_0^{\frac{\pi}{2}} \tan 2x \ln(\tan x) dx$

C) $\frac{\pi}{8} \int_{-1}^1 \frac{\ln(1-x)}{x} dx$

D) $\frac{\pi}{8} \int_0^{\pi} \tan x \ln(1 - \cos x) dx$

50. $\int_0^{\infty} \frac{\tan^{-1}(x)}{(1+x)\sqrt{x}} dx =$

A) $\int_0^{\infty} \frac{\arctan(1/x)}{\sqrt{x}(1+x)} dx$

B) $\frac{\pi}{4} \int_0^{\infty} \frac{dx}{\sqrt{x}(1+x)}$

C) $\frac{\pi}{2} \int_0^{\infty} \frac{dx}{\sqrt{x}(1+x)}$

D) $\frac{\pi^2}{4}$

51. $\int_0^{\infty} \frac{x-1}{x^8-1} dx =$

A) $\frac{1}{2} \int_0^{\infty} \frac{dx}{x^4-1} - \int_0^{\infty} \frac{dx}{x^8-1}$

B) $\int_0^{\infty} \frac{1}{x^4+1} dx$

C) $\frac{1}{2} \int_0^{\infty} \frac{1}{x^4+1} dx$

D) $\frac{\pi}{4\sqrt{2}}$

JEE MAIN
2023SINGARAJU
VENKAT KOUNDINYA
RANK 1
Sri Chaitanya
JEE Prep Class300
300
MARKS

RANK

1

JEE Advanced
2023VAVILALA
CHIDVILAS REDDY
RANK 1
Sri Chaitanya
JEE Prep Class341
360
MARKS

RANK

1

NEET
2023BORA VARUN
CHAKRAVARTHI
RANK 1
Sri Chaitanya
JEE Prep Class720
720
MARKS

RANK

1



52. $\int_0^{\infty} \frac{1}{(1+x^{2023})(1+x^2)} dx =$

A) $\frac{\pi}{2}$

B) $\frac{\pi}{4}$

C) $\int_0^{\infty} \frac{x^{2023}}{(1+x^{2023})(1+x^2)} dx$

D) $\int_0^{\infty} \frac{1}{1+4x^2} dx$

53. $\int_{\frac{1}{2}}^2 \log \left(\frac{\log \left(x + \frac{1}{x} \right)}{\log \left(x^2 - x + \frac{17}{4} \right)} \right) dx =$

A) $\int_{1/2}^2 \log \log \left(x + \frac{1}{x} \right) dx - \int_0^{3/2} \log \log (x^2 + 4) dx$

B) $\int_{1/2}^2 \log \log \left(x + \frac{1}{x} \right) dx - \int_{1/2}^2 \log \log \left(x^2 - x + \frac{17}{4} \right) dx$

C) $\int_{1/2}^2 \left(\log \log \left(x + \frac{1}{x} \right) - \log \log \sqrt{x^2 - x + \frac{17}{4}} - \log 2 \right) dx$

D) $-\frac{3}{2} \log 2$

54. $\int_0^{1/\sqrt{3}} \sqrt{x + \sqrt{x^2 + 1}} dx =$

A) $\int_0^{\pi/6} \sqrt{\tan(\theta) + \sec(\theta)} \sec^2(\theta) d\theta$

B) $\frac{1}{2} + \frac{1}{4} \int_0^{\pi/6} \sqrt{\tan(\theta) + \sec(\theta)} \sec^2(\theta) d\theta$

C) $\frac{1}{\sqrt{3}} - \frac{1}{2} \int_0^{\pi/6} \sqrt{\tan(\theta) + \sec(\theta)} d(\sec(\theta))$

D) $\frac{2}{3}$





SECTION 4

- This section contains **THREE (03)** question.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer the using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:
- **Full Marks** : +4 If ONLY the correct integer is entered;
- **Zero Marks** : 0 In all other cases.

55. $\int_{-9}^9 \underbrace{\left| \left| \left| \left| x \right| - 1 \right| - 1 \right| - 1 \dots \dots \dots - 1 \right|}_{9 \text{ times}} dx =$

56. $\lim_{n \rightarrow \infty} \int_0^{\infty} \left(1 + \frac{t}{n}\right)^{-n} \cdot \cos\left(\frac{t}{n}\right) dt =$

57. $\int_0^9 \{x[x]\} dx = \left(\begin{array}{l} \text{The greatest integer function is denoted by } [x] \\ \text{where as the fractional part function is denoted by } \{x\}. \end{array} \right)$





Sri Chaitanya

Junior Colleges & Coaching Centres

Infinity
Learn BY SRI CHAITANYA



VAVILALA CHIDVILAS REDDY
HT.NO. 236165088
Sri Chaitanya
6th - 12th Class

RANK

WITH ALL INDIA **RANK 1** IN JEE ADVANCED 2023

SRI CHAITANYA

STANDS AT THE TOP

SEIZES 5 RANKS IN TOP 10 IN ALL-INDIA OPEN CATEGORY

ANDHRA PRADESH STATE TOPPER



RAMESH SURYA THEJA
HT.NO. 236131371

8th - 12th Class

Sri Chaitanya



RANK



KALRA RISHI
HT.No. 237047209

AITS Student



RANK



RAGHAV GOYAL
HT.NO. 237020529

9th - 12th Class

Sri Chaitanya



RANK



BIKKINA ABHINAV CHOWDARY
HT.NO. 236165128

4th - 12th Class

Sri Chaitanya



RANK

32 TOP RANKS BELOW 100 IN ALL-INDIA OPEN CATEGORY

RANK 12 DASARI SAKETH NAIDU HT.No. 236060149	RANK 14 GUTHIKONDA ABHIRAM HT.No. 236128721	RANK 17 P GNANA KOUSIK REDDY HT.No. 236135530	RANK 18 D VENKATA YUGESH HT.No. 236145359	RANK 19 MOULIK JINDAL HT.No. 237021004	RANK 22 D PRATAP SINGH HT.No. 232006139	RANK 23 SHAIK SAHIL CHANDA HT.No. 236132870	RANK 26 SONI MAYANK HT.No. 232097025	RANK 33 GOYAL NAMAN HT.No. 232029345
RANK 37 SUMEDH SS HT.No. 231028080	RANK 38 V DEERAJ SATHVIK REDDY HT.No. 236060751	RANK 44 V VERGIYA KAUSHAL HT.No. 233004058	RANK 46 SUTHAR HARSHAL HT.No. 233009060	RANK 58 ABHI JAIN HT.No. 232091036	RANK 61 RIDAM JAIN HT.No. 232068021	RANK 62 SUTHAR HARSH HT.No. 232100028	RANK 63 SOJITRA DIPEN HT.No. 231077001	RANK 65 D PANEENDRANADHA REDDY HT.No. 236080208
RANK 67 GUNDA SUSHANTH HT.No. 236128732	RANK 68 HARSH TAYA HT.No. 237021778	RANK 69 KRISH GUPTA HT.No. 232008189	RANK 73 KUNCHAKURTHY VIVEK HT.No. 236134083	RANK 81 N DHARMA TEJA REDDY HT.No. 236050138	RANK 84 S VENKAT KOUNDINYA HT.No. 230130678	RANK 87 MRUNAL SHRIKANT HT.No. 232104138	RANK 92 SAMOTA APURVA HT.No. 232099210	RANK 97 N PRINCE BRANHAM REDDY HT.No. 236064112

BELOW
20
All India Open
Category Ranks

10
RANKS (50%)

BELOW
100
All India Open
Category Ranks

32

BELOW
1000
All India Open
Category Ranks

181

BELOW
100
All India
Category Ranks
Count

89

BELOW
1000
All India
Category Ranks
Count

699

NUMBER OF RANKS QUALIFIED
3,621

ADMISSIONS OPEN » JEE ADVANCED LONG-TERM 2024

040 66 06 06 06



Scan QR Code for
JEE Advanced Offers

www.srichaitanya.net