



Sri Chaitanya IIT Academy.,India.

✪ A.P ✪ T.S ✪ KARNATAKA ✪ TAMILNADU ✪ MAHARASTRA ✪ DELHI ✪ RANCHI

A right Choice for the Real Aspirant

ICON Central Office - Madhapur - Hyderabad

SEC: **Sr.Super60_NUCLEUS&STERLING_BT**

JEE-MAIN

Date: 02-09-2023

Time: 09.00Am to 12.00Pm

RPTM-05

Max. Marks: 300

IMPORTANT INSTRUCTION:

1. Immediately fill in the Admission number on this page of the Test Booklet with **Blue/Black Ball Point Pen** only.
2. The candidates should not write their Admission Number anywhere (except in the specified space) on the Test Booklet/ Answer Sheet.
3. The test is of **3 hours** duration.
4. The Test Booklet consists of 90 questions. The maximum marks are **300**.
5. There are **three** parts in the question paper 1,2,3 consisting of **Physics, Chemistry and Mathematics** having **30 questions** in each subject and subject having **two sections**.

(I) Section –I contains 20 **multiple choice** questions with only one correct option.


Marking scheme: +4 for correct answer, 0 if not attempt and -1 in all other cases.

(II) Section-II contains 10 **Numerical Value Type** questions. Attempt any 5 questions only, if more than 5 questions attempted, First 5 attempted questions will be considered.

- The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).

To cancel any attempted question bubble on the question number box.

For example: To cancel attempted question 21. Bubble on 21 as shown below



Question Answered for Marking

Question Cancelled for Marking

Marking scheme: +4 for correct answer, 0 if **not attempt** and -1 in all other cases.

6. Use **Blue / Black Point Pen only** for writing particulars / marking responses on the Answer Sheet. **Use of pencil is strictly prohibited.**
7. No candidate is allowed to carry any textual material, printed or written, bits of papers, mobile phone any electron device etc, except the Identity Card inside the examination hall.
8. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
9. On completion of the test, the candidate must hand over the Answer Sheet to the invigilator on duty in the Hall. **However, the candidate are allowed to take away this Test Booklet with them.**
10. **Do not fold of make any stray marks on the Answer Sheet**

Name of the Candidate (in Capital): _____

Admission Number:

--	--	--	--	--	--	--	--	--	--

Candidate's Signature: _____

Invigilator's Signature: _____

**02-09-23_Sr.Super60_NUCLEUS&STERLING_BT_Jee-Main_RPTM-05_Test Syllabus****PHYSICS**

: NLM, Friction, Circular motion, WPE Newton's laws of motion; Inertial and uniformly accelerated frames of reference; Static and dynamic friction; Kinetic and potential energy; Work and power, Uniform circular motion, Conservation of mechanical energy

RPTM-05 (15Q-RPTM.5 SYLLABUS+15Q CUMULATIVE SYLLABUS)**CHEMISTRY**

: Alcohols, Phenols, Ethers & Amines: Alcohols : Physical properties; Reactions: esterification, dehydration (formation of alkenes and ethers); Reactions with: sodium, phosphorus halides, ZnCl_2 /concentrated HCl, thionyl chloride; Conversion of alcohols into aldehydes, ketones and carboxylic acids
Phenols: Physical properties; Preparation, Electrophilic substitution reactions of phenol (halogenation, nitration, sulphonation); Reimer-Tiemann reaction, Kolbe reaction; Esterification; Etherification; Aspirin synthesis; Oxidation and reduction reactions of phenol.
Ethers : Preparation, Properties & Reactions
Amines: Basicity of substituted anilines and aliphatic amines, Preparation from nitro compounds, nitriles and amides; Reactions: Hoffmann bromamide degradation, Gabriel phthalimide synthesis; Reaction with nitrous acid, Azo coupling reaction of diazonium salts of aromatic amines; Sandmeyer and related reactions of diazonium salts; Carbylamine reaction, Hinsberg test, Alkylation and acylation reaction

RPTM-05 (15Q-RPTM.5 SYLLABUS+15Q CUMULATIVE SYLLABUS)**MATHEMATICS : Definite Integration****RPTM-05 (15Q-RPTM.5 SYLLABUS+15Q CUMULATIVE SYLLABUS)**


**PHYSICS****Max Marks: 100****(SINGLE CORRECT ANSWER TYPE)**

This section contains 20 multiple choice questions. Each question has 4 options (1), (2), (3) and (4) for its answer, out of which ONLY ONE option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.


1. Statement I: A cyclist is moving on an unbanked road with a speed of 7 kmh^{-1} and takes a sharp circular turn along a path of radius of 2 m without reducing the speed. The static friction coefficient is 0.2. The cyclist will not slip and pass the curve. ($g = 9.8\text{ m/s}^2$)
- Statement II: If the road is banked at an angle of 45° cyclist can cross the curve of 2 m radius with the speed of 18.5 kmh^{-1} without slipping.
- In the light of the above statements, choose the correct answer from the options given below.
- 1) Statement I is correct and statement II is incorrect
 - 2) Statement I is incorrect and statement II is correct
 - 3) Both statement I and statement II are true
 - 4) Both statement I and statement II are false
2. A man pushes eight identical blocks on horizontal frictionless surface with horizontal force F . The force that block-1 exerts at block-2 has magnitude F_{21} and the force that block-7 exerts on the block -8 is F_{87} . Find $\frac{F_{21}}{F_{87}}$
- \xrightarrow{F}
- | | | | | | | | |
|---|---|---|---|---|---|---|---|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 9 |
|---|---|---|---|---|---|---|---|
- 1) 3
 - 2) 5
 - 3) 7
 - 4) 9
3. A wedge of mass $M = 10\text{ kg}$, height $h = 3\text{ m}$ and angle of inclination $\alpha = 37^\circ$ is at rest on a horizontal surface. There is a small point-like object (mass $m = 0.5\text{ kg}$) next to the slope as shown in the figure. At what acceleration must wedge be moved in order that the point like object reaches its top in a time $t = 5\text{ s}$ (Neglect the friction between point like object and wedge)





- 1) $2m / s^2$ 2) $4m / s^2$ 3) $8m / s^2$ 4) $10m / s^2$

4. An object of mass M starts from rest at the bottom of a fixed incline of height H . A person decides to push the object up the incline in one of two ways with an applied force shown in the diagram. In each of the trials, the object reaches the top of the incline with speed V . How would the work done by the person on the block compare for the two trials? Assume the same constant non-zero coefficient of kinetic friction between the incline and the object for both trials




- 1) More work would be done in Trial 1.
 2) More work would be done in Trial 2
 3) It is impossible to determine for which trial there would be more work done without knowing the value of the speed V .
 4) It is impossible to determine for which trial there would be more work done Without knowing the value of the coefficient of kinetic friction






5. All the blocks are attached to an ideal pulley. If accelerations of blocks $m_1 = m$, $m_2 = 3m$, $m_3 = m$, $m_4 = 2m$ are a_1, a_2, a_3 and a_4 respectively then choose the correct option for a_1, a_2, a_3 and a_4 respectively




- 1) $a_1 \uparrow g$, $a_2 \uparrow 2g$, $a_3 \uparrow 3g$, $a_4 \downarrow g/2$ 2) $a_1 \uparrow g$, $a_2 \downarrow 2g/3$, $a_3 \uparrow 3g$, $a_4 \downarrow g/2$
- 3) $a_1 \downarrow 2g/3$, $a_2 \uparrow 2g$, $a_3 \uparrow 3g$, $a_4 \uparrow g$ 4) $a_1 \downarrow g$, $a_2 \downarrow g/2$, $a_3 \downarrow 3g$, $a_4 \downarrow 2g/3$
6. A string of mass m (can be non uniform as well) is suspended through two points which are not in same horizontal level. Tension in the string at the end points are T_1 , and T_2 and at the lowest point is T_3 . Mass of string in terms of T_1, T_2 and T_3 can be represented as (uniform gravity 'g' exists downwards)



- 1) $\frac{\sqrt{T_1^2 - T_3^2} + \sqrt{T_2^2 - T_3^2}}{g}$ 2) $\frac{(T_1 + T_2 - 2T_3)}{g}$ 3) $\frac{\sqrt{T_1^2 + T_2^2}}{g}$ 4) $\frac{\sqrt{T_1^2 + T_3^2} + \sqrt{T_2^2 + T_3^2}}{g}$



7. A man of mass 60 kg hangs himself from a massless spring balance. Which itself suspended from an ideal string-pulley system as shown in the figure. The string AB can bear maximum 900 N. Choose correct statement.




- 1) The man is never ever able to read his correct weight in the spring balance.
 - 2) If man able to manage himself at rest, the spring balance reads his true weight.
 - 3) The maximum magnitude of acceleration, man can manage is 12.5 m/s^2 while Moving upward.
 - 4) When man moves downward with acceleration 5 m/s^2 , the spring balance will read 40 kg
8. Given below are two statements:
 Statement-I: An elevator can go up or down with uniform speed when its weight is balanced with the tension of its cable.
 Statement-II: Force exerted by the floor of an elevator on the foot of a person standing on it is more than his/her weight when the elevator goes down with increasing speed. In the light of the above statements, choose the correct answer from the options given below:
- 1) Both statement I and statement II are false
 - 2) Statement I is true but Statement II is false
 - 3) Both Statement I and Statement II are true
 - 4) Statement I is false but Statement II is true






9. A particle whose velocity is given as $\vec{v} = \hat{i} + 6t\hat{j} \text{ m/s}$ is moving in $x-y$ plane. At $t = 0$, particle is at origin. Find the radius of curvature of path at point $\left(\frac{\sqrt{2}}{3}m, \frac{2}{3}m\right)$
- 1) $1.5m$ 2) $3.0m$ 3) $4.5m$ 4) $6.0m$
10. An ideal string-pulley system is shown in the figure and system is released from rest. Choose correct option.



- 1) When mass B moves with speed $4m/s$, the speed of mass A is $1m/s$.
- 2) When mass B moves with speed $1m/s$, the speed of mass A is $4m/s$.
- 3) In the time interval in which the mass A covers distance $8m$, in the same time interval, mass B travels a distance $1m$.
- 4) In the time interval in which the mass A covers distance $1m$, in the same time interval, mass B travels a distance $8m$.
11. In the given figure, a long platform of mass m is placed on a frictionless surface. Two blocks of masses $4m$ and m (where $m=10 \text{ kg}$) are placed on the platform. For both blocks, the coefficient of static friction with the platform equal to 0.16 and the coefficient of kinetic friction is equal to 0.10 . The blocks are connected by a light ideal string through a light pulley (mounted at a movable massless stand), which is acted upon by an unknown horizontal force F . If the acceleration of the platform is $2m/s^2$, find the value of unknown force F and acceleration of blocks $4m$ and m as a_1 and a_2 respectively.



$$F(N) \quad a_1(m/s^2) \quad a_2(m/s^2)$$

1) $180N, 2m/s^2, 8m/s^2$

2) $180N, 8m/s^2, 2m/s^2$

3) $90N, 2m/s^2, 8m/s^2$

4) $90N, 8m/s^2, 2m/s^2$

12. Given below are two statements:

Statement I: Area under velocity- time graph gives the displacement travelled by the body in a given time.

Statement II: Area under acceleration –time graph is equal to velocity- in the given time.

In the light of given statements, choose the correct answer from the options given below:


1) Both Statement I and Statement II are true.

2) Statement I is correct but Statement II is false.

3) Statement I is incorrect but Statement II is true.

4) Both Statement I and Statement II are false

13. Point A on the rod AB has an acceleration of $5m/s^2$ and a velocity of $6m/s$ at an instant as shown in the figure. The acceleration of the end B at the same moment is:



1) $-\frac{80}{3} \hat{i} m/s^2$

2) $\frac{80}{3} \hat{i} m/s^2$

3) $-\frac{40}{3} \hat{i} m/s^2$


4) $-\frac{75}{6} \hat{i} m/s^2$






14. The question has Statement 1 and Statement 2. Of the four choices given after the Statements, choose the one that best describes the two Statements.
- Statement 1: If you push on a cart being pulled by a horse so that it does not move, the cart pushes you back with an equal and opposite force.
- Statement 2: The cart does not move because the force described in statement 1 cancel each other.
- 1) Statement 1 is true, Statement 2 is true, Statement 2 is the correct explanation of Statement 1.
 - 2) Statement 1 is false, Statement 2 is true.
 - 3) Statement 1 is true, Statement 2 is false.
 - 4) Statement 1 is true, Statement 2 is true, Statement 2 is not the correct explanation of Statement 1.
15. Given below are two statements:
- Statement I: If heat is added to a system, its temperature must increase.
- Statement II: If positive work is done by a system in a thermodynamic process, its volume must increase.
- In the light of the above statement, choose the correct answer from the options given below:
- 1) Statement I is true but Statement II is false
 - 2) Both Statement I and Statement II are true
 - 3) Both Statement I and Statement II are false
 - 4) Statement I is false but Statement II is true
16. Two moles of an idea monoatomic gas is taken through a cyclic process as shown in the P-T diagram. In the process BC, $PT^{-2} = \text{constant}$. Then the ratio of heat absorbed and heat released by the gas during the process AB and process BC respectively is





- 1) 2 2) 3 3) 5 4) 6

17. Two ends of a rod of uniform cross sectional area are kept at temperatures $3T_0$ and T_0 as shown. The thermal conductivity of the rod varies as $k = \alpha T$, (where α is a constant and T is absolute temperature). In steady state, the temperature of the middle section of the rod is



- 1) $\sqrt{7}T_0$ 2) $\sqrt{5}T_0$ 3) $2T_0$ 4) $\sqrt{3}T_0$


18. The molar heat capacity of an ideal gas in a process varies as $C = C_v + \alpha T^2$ (where C_v is molar heat capacity at constant volume and α is a constant). Then the equation of the process is

- 1) $Ve^{-\left(\frac{\alpha T^2}{2R}\right)} = \text{constant}$ 2) $Ve^{-\left(\frac{\alpha T^2}{R}\right)} = \text{constant}$
 3) $Ve^{-\left(\frac{2\alpha T^2}{R}\right)} = \text{constant}$ 4) $Ve^{-\left(\frac{3\alpha T^2}{2R}\right)} = \text{constant}$





19. When a particle of mass m is suspended from a massless spring of natural length ℓ , the length of the spring becomes 2ℓ . When the same mass moves in conical pendulum, the length of the spring becomes L . The radius of the circle of conical pendulum is:
- 1) $\frac{L}{L-\ell}\sqrt{L(L-\ell)}$ 2) $\frac{L}{L-\ell}\sqrt{L(L-2\ell)}$
 3) $\sqrt{L(L-\ell)}$ 4) $\sqrt{L^2-2\ell^2}$
20. A point object is kept at a distance of 2 m from origin on X- axis. A quarter cylinder of radius $R=30$ cm and refractive index $\mu=1.5$ having its axis \perp to plane of paper is placed between the object and the parabolic reflector at a distance of 130 cm from origin as shown in figure. The position of the image after reflection from the parabolic reflector $y^2 = 2x$ is



- 1) 25 cm 2) 50 cm 3) 100 cm 4) 150 cm


(NUMERICAL VALUE TYPE)

Section-II contains 10 **Numerical Value Type** questions. Attempt any 5 questions only. First 5 attempted questions will be considered if more than 5 questions attempted. The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).


Marking scheme: +4 for correct answer, 0 if **not attempt** and -1 in all other cases.

21. A cylinder of mass M and radius R moves with constant speed V through a region of space that contains dust particles of mass m ($m \ll M$) which are at rest. There are n number of particles per unit volume. The cylinder moves in a direction perpendicular to its axis. Assume that the particles do not interact with each other and bounce with same speed after hitting cylinder. The drag force per unit length of the cylinder require to maintain the speed v constant for the cylinder is $\frac{K}{3}nmRv^2$ Find the value of K ?






22. A spherical aquarium of radius 10 cm filled with water of refractive index $4/3$ is kept as shown. One half of aquarium is silvered and other half is smooth. The thickness of aquarium is negligible. A fish is moving towards left with a velocity 6 m/s. An observer detects two images of the fish. Find the relative velocity (in m/s) between the two images of the fish as seen by the observer.



23. A parallel beam of monochromatic light of wavelength 4500 \AA is incident of YDSE. Let I_0 is the intensity of incident light. Now two glass slabs G_1 and G_2 are kept in front of both slits S_1 and S_2 . The refractive index of G_1 is 1.9 and for G_2 it is 1.6. The thickness of each glass slab is $15 \mu\text{m}$. It is observed that glass G_1 transmits $\frac{1}{16}$ th and G_2 transmits $\frac{1}{25}$ th of incident energy. If $I_0 = 8 \text{ Watt / m}^2$, then find the intensity (in W / m^2) observed at point 'O' on the screen




24. Suppose that one of the slits in YDSE is wider than the other, so the amplitude of the light reaching on screen from one slit, acting alone, is thrice that from the other slit, acting alone. The intensity for Central Bright Fringe is $I_0 = 5 \text{ W / m}^2$. Then find the intensity $I (\text{W / m}^2)$ on the screen for (angular position) $\theta = 37^\circ$. (Take $d / \lambda = 5 / 6$, where d is the separation between the slits and λ is the Wave length of the light)






25. A uniform ring of mass m is placed on a rough horizontal fixed surface as shown in the figure. The coefficient of friction between the left part of the ring and left part of the horizontal surface is $\mu_1 = 0.6\pi$ and between right half and the surface is $\mu_2 = 0.2\pi$. At the instant shown, now the ring has been imparted an angular velocity in clockwise sense in the figure shown. At this moment magnitude of acceleration of centre O of the ring (in m/s^2) is (take $g = 10 m/s^2$)




26. A gas bubble produced from a under water explosion oscillates with period T . The period T is found to be proportional to the a^{th} power of pressure (P), b^{th} power of density (d) and c^{th} power of the total energy (E) of explosion. Using dimensional analysis find the value of $(a + 4b + c)$. (Round off to the closest integer)
27. Light from a monochromatic point source S is incident on the two slits of an opaque wall placed at a distance L as shown in the diagram, Intensity of interference pattern is observed at the point P on the screen placed at a distance L behind the slits. When the upper slit is shifted upward then the distance between the slits is changed from $200 \mu m$ to $400 \mu m$, the intensity at P changes from I_{\max} to $\frac{3}{4}I_{\max}$. If $L=1$ m, then find the wave length of light used (in \AA)






28. A power output from a certain experimental car design to be shaped like a cube is proportional to the mass m of the car. The force of air friction on the car is proportional to Av^2 , where v is the speed of the car and A is the cross-sectional area. On a level surface the car has a maximum speed v_{\max} . Assume that all versions of this design have the same density. Then v_{\max} is proportional to $m^{1/c}$. Find C .
29. A flexible drive belt runs over a frictionless pulley as shown in figure. The pulley is rotating freely about the vertical axis passing through the centre O of the pulley. The vertical axis is fixed on the horizontal smooth surface. The mass per unit length of the drive belt is $1 \text{ kg} / m$ and the tension in the drive belt 8 N . The speed of the drive belt is 2 m/s . Find the net normal force applied by the belt on the pulley in Newton.



30. Two identical particles A & B of mass $m = 2 \text{ kg}$ is connected at the end of the spring ($k = 50 \text{ N} / m$) and placed on a smooth horizontal surface. Now, force $F = 10 \text{ N}$ and $2F$ are applied on the particles in opposite direction as shown in figure. Initially spring is in natural length. Simultaneously particle A receives an impulse towards B which imparts speed v_0 to it find v_0 in m/s for which length of elongation is two times the length of compression in the spring during subsequent motion?




**CHEMISTRY****Max Marks: 100****(SINGLE CORRECT ANSWER TYPE)**

This section contains 20 multiple choice questions. Each question has 4 options (1), (2), (3) and (4) for its answer, out of which ONLY ONE option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

31. The major product formed in following reaction is:



32. Given below are two statements one is labeled as assertion (A) and other is labeled as Reason (R)

Assertion (A): Synthesis of vinyl phenyl ether may be achieved by Williamson synthesis

Reason (R): chlorobenzene reacts with sodium vinyl oxide yields vinyl phenyl ether



In the light of the above statements choose the most appropriate answer from the options given below.

- 1) Both (A) and (R) are correct and (R) is the correct explanation of (A)
- 2) (A) is correct but (R) is not correct
- 3) Both (A) and (R) not correct
- 4) Both (A) and (R) are correct (R) is not correct explanation of (A)






33. Consider the following reaction $C_2H_5OH + H_2SO_4 \rightarrow \text{product}$ among the following which one cannot be formed as a product under any condition?
1) Ethylene 2) Acetylene 3) Diethyl ether 4) Ethyl hydrogen sulphate
34. **Assertion (A)** : Ethoxy ethane and butanol-1 have almost same miscibility with water
Reason (R) : Ethoxy ethane and butanol-1 have form Hydrogen bond with water
1) Both (A) and (R) are correct and (R) is the correct explanation of (A)
2) (A) is correct but (R) is not correct
3) Both (A) and (R) are not correct
4) Both (A) and (R) are correct (R) is not correct explanation of (A)
35. Phenol industrially prepared from
1) benzene sulphonic acid 2) Cumene
3) benzene diazonium chloride 4) chlorobenzene
36. Which of the following statements is incorrect
1) benzene diazonium fluoroborate is stable at room temperature
2) benzene diazonium fluoroborate is almost insoluble in water
3) diazonium salts stability due to '+I' effect of benzene
4) cyanobenzene easily prepared from diazonium salt






38. Identify the correct acidic strength order of following pairs





39. From which of the following compound Grignard reagent cannot be prepared directly using ether and magnesium







40. The correct order of acidity of the following hydrocarbon is



- 1) $i > ii > iii > iv$ 2) $iii > iv > ii > i$ 3) $i > iii > iv > ii$ 4) $iv > iii > i > ii$

41. Given below are two statements for the



Statement I: the value of k_2 less than K_1

Statement II: Phenoxide is a weaker base than ethoxide

In the light of the above statements, choose the most appropriate answer from the options given below


- 1) Both statement I and statement II are correct
 2) Both statement I and statement II are incorrect
 3) Statement I is correct but statement II is incorrect
 4) Statement II is correct but statement I is incorrect
42. Which of the following statement is incorrect
- 1) methylene chloride used as paint remover
 2) chloroform is used to produce Freon Refrigerant
 3) Chlorofluoro carbon compounds of methane, ethane, propane are collectively known as freons
 4) Tetrachloro methane is used to prepare chlorofluoro carbons





43. Among the following which will be the most powerful Nucleophile in water?

- 1) F^- 2) Cl^- 3) Br^- 4) I^-

44. **Assertion:**  bond angle at oxygen is less than ideal tetrahedral bond angle.

Reason: Lone pair-lone pair repulsion decreases bond angle in methanol at oxygen


- 1) Both Assertion and Reason are true and Reason is the correct explanation of Assertion
 2) Both Assertion and Reason are true but Reason is not the correct explanation of Assertion
 3) Assertion is true but Reason is false
 4) Both Assertion and Reason are false

45. **Assertion:** Aryl sulphonic acid on reacting with $NaOH$ at high temperature followed by acidification gives phenol

Reason: This reaction is electrophilic substitution reaction

- 1) Both Assertion and Reason are true and Reason is the correct explanation of Assertion
 2) Both Assertion and Reason are true but Reason is not the correct explanation of Assertion
 3) Assertion is true but Reason is false
 4) Both Assertion and Reason are false

46. Which of the following involving markonikov's addition





47. Number of moles of HI consumed by glycerol to give 3-Iodopropene

- 1) 5 2) 3 3) 4 4) 6

48. Number of resonance structures possible in tropylium cation

- 1) 7 2) 3 3) 5 4) 6





49.

Number of unhybridised orbitals in compound 'c'

- 1) 18 2) 12 3) 6 4) 24

50. In which of the following, polysubstitution may takes place?


- 1) $PhH + CH_3COCl \xrightarrow{\text{Anhyd. } AlCl_3} Ph - \overset{\overset{O}{\parallel}}{C} - CH_3$
 2) $PhH + CH_3Cl \xrightarrow{\text{Anhyd. } AlCl_3} Ph - CH_3$
 3) $PhH + HNO_3 \xrightarrow{H_2SO_4} Ph - NO_2$
 4) $PhH + H_2SO_4 \xrightarrow{90^\circ C} Ph - SO_3H$

(NUMERICAL VALUE TYPE)


Section-II contains 10 **Numerical Value Type** questions. Attempt any 5 questions only. First 5 attempted questions will be considered if more than 5 questions attempted. The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).

Marking scheme: +4 for correct answer, 0 if **not attempt** and -1 in all other cases.


51. Number of compounds giving (i) Red colouration with ceric ammonium nitrate and also (ii) Blue colouration with victor mayer test




(i)



(ii)



(iii)




(iv)



(v)



(vi)



(vii)




(viii)





52. how many of the following are more basic than phenyl methanamine

- i) Benzanamine ii) N, N-dimethyl aniline
 iii) N-Ethylethanamine iv) N, N-dimethyl methanamine
 v) N-methyl phenyl methanamine vi) 2,6-dimethylaniline
 vii) N-phenyl ethanamide




- 53.** How many of the following are white precipitates
 i) 2,4,6-tribromophenol ii) Iodoform
 iii) 2,4,6 tribromo aniline iv) Chloroform
 v) P-aminoazobenze
- 54.** Number of unshared electron pairs in electrophile of carbylamine test
- 55.** The number of substrates those can give both SN¹ and SN² reactions are



- 56.** How many of the following are having more heat of hydrogenation than hexa-2,4-diene
- i)  ii)  iii) 
- iv) 

- 57.** Number of stereo centres of the following compound



58. Number of planar atoms in propyne
59. $\text{ClCH}_2 - \text{CH}_2 - \text{Cl} \xrightarrow[2) \text{C}_2\text{H}_5\text{Cl}]{1) x \text{ mole NaNH}_2} \text{C}_2\text{H}_5 - \text{C} \equiv \text{C} - \text{C}_2\text{H}_5$
60. Number of amide isomers exist for $\text{C}_5\text{H}_{11}\text{ON}$ which on treatment with $\text{NaOH} / \text{Br}_2$ will give amines

**MATHEMATICS****Max Marks: 100****(SINGLE CORRECT ANSWER TYPE)**

This section contains 20 multiple choice questions. Each question has 4 options (1), (2), (3) and (4) for its answer, out of which ONLY ONE option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

61. Let $f : R \rightarrow [4, \infty)$ be an onto quadratic function whose leading coefficient is 1, such that $f'(x) + f'(2-x) = 0$. Then the value of $\int_1^3 \frac{dx}{f(x)}$ is equal to
- 1) $\frac{1}{4}$ 2) $\frac{1}{8}$ 3) $\frac{\pi}{4}$ 4) $\frac{\pi}{8}$
62. Let g be a continuous even function on R such that $g(x) = f^1(x)$ and $f(1) = 3$ then $\int_{-1}^1 f^2(x) g(x) dx$ is equal to
- 1) 0 2) 9 3) 18 4) 27
63. Let f be a differentiable function satisfying $\int_1^{f(x)} f^{-1}(t) dt = \frac{1}{3} \left(x^{\frac{3}{2}} - 8 \right) \forall x > 0$ and $f(1) = 0$, then the value of $f(9)$ is
- 1) 3 2) 9 3) 0 4) 2
64. Match the elements of **Column-I** with **Column-II**
- | Column-I | Column-II |
|--|--------------------------|
| (A) $f(x) = \sin x - x^2 + 1$ | (p) Has point of minima |
| (B) $f(x) = x \log_e x - x + e^{-x}$ | (q) Has point of maxima |
| (C) $f(x) = -x^3 + 2x^2 - 3x + 1$ | (r) Is always increasing |
| (D) $f(x) = \cos \pi x + 10x + 3x^2 + x^3$ | (s) Is always decreasing |
- a) A-p B-q C-r D-s b) A-r B-s C-p D-q
c) A-s B-r C-q D-p d) A-q B-p C-s D-r
65. **Statement 1:** If $27a + 9b + 3c + d = 0$, then the equation $f(x) = 4ax^3 + 3bx^2 + 2cx + d = 0$ has at least one real root lying in interval $(0, 3)$
- Statement 2:** If $f(x)$ is continuous in $[a, b]$, derivable in (a, b) such that $f(a) = f(b)$, then there exist at least one point $c \in (a, b)$ such that $f'(c) = 0$





1) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.

2) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.

3) Statement-1 is true, statement-2 is false.

4) Statement-1 is false, statement-2 is true.

66. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differential function such that

$$f\left(\frac{\pi}{4}\right) = \sqrt{2}, f\left(\frac{\pi}{2}\right) = 0, \text{ and } f'\left(\frac{\pi}{2}\right) = 1 \text{ and let}$$

$$g(x) = \int_x^{\pi/4} (f'(t)\sec(t) + \sec(t)\tan(t)f(t))dt \text{ for } x \in \left[\frac{\pi}{4}, \frac{\pi}{2}\right] \text{ Then}$$

$\lim_{x \rightarrow \left(\frac{\pi}{2}\right)^-} g(x)$ is equal to

1) 2

2) 3

3) 4

4) -3

67. If $U_n = \left(1 + \frac{1}{n^2}\right)\left(1 + \frac{2^2}{n^2}\right) \dots \left(1 + \frac{n^2}{n^2}\right)^n$, then $\lim_{n \rightarrow \infty} (U_n)^{\frac{-4}{n^2}}$ is equal to

1) $\frac{16}{e^2}$

2) $\frac{4}{e}$

3) $\frac{4}{e^2}$

4) $\frac{e^2}{16}$

68. Let $f(x)$ be a continuous function on \mathbb{R} .

$$\text{If } \int_0^1 (f(x) - f(2x))dx = 5 \text{ and } \int_0^2 (f(x) - f(4x))dx = 10$$

Then the value of $\int_0^1 (f(x) - f(8x))dx$ is equal to

1) 0

2) 5

3) 10

4) 15

69. **Statement 1:** For all $a, b \in \mathbb{R}$ the function $f(x) = 3x^4 - 4x^3 + 6x^2 + ax + b$ has exactly one extremum

Statement 2: If a cubic function is monotonic, then its graph cuts the x -axis only once





- 1) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
 2) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
 3) Statement-1 is true, statement-2 is false.
 4) Statement-1 is false, statement-2 is true.

70. **Statement-1:** Let $I_k = \int_0^{\pi} \frac{\sin kx}{\sin x} dx$ Then $\sum_{k=1}^{2022} I_k = 1011\pi$

Statement-2: $\sin(n\pi - \theta) = \begin{cases} -\sin \theta & \text{when } n \text{ is even} \\ \sin \theta & \text{when } n \text{ is odd} \end{cases}$

- 1) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
 2) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
 3) Statement-1 is true, statement-2 is false.
 4) Statement-1 is false, statement-2 is true.

71. If $f(x) = x^5 + 5x - 1$ then $\int_5^{41} \frac{dx}{(f^{-1}(x))^5 + 5 \cdot f^{-1}(x)}$ equals

- 1) 0 2) $\ln 2$ 3) $\ln 7$ 4) $\ln \left(\frac{41}{5} \right)$

72. $f(x) = \int \frac{5e^{8\log x} + 7e^{6\log x}}{(e^{2\log x} + 2e^{7\log x} + 1)^2} dx (x > 0)$

and $f(1) = 1/4$ then $f(2) =$

- 1) $\frac{128}{7}$ 2) $\frac{128}{261}$ 3) $\frac{256}{261}$ 4) $\frac{216}{261}$

73. If $f(x) = \int \frac{x^2 \sin 2x - \sin 2x(x \tan x - 1)}{(x^2 + 1)^2} dx$

and $f(0) = 0$ then $f(\pi/4) =$

- 1) $\frac{8}{\pi^2 + 16}$ 2) $\frac{4}{\pi^2 + 16}$ 3) $\frac{1}{\pi^2 + 16}$ 4) $\frac{1}{\pi^2 + 9}$





74. Let f be a continuous function on \mathbb{R} such that

$$f\left(\frac{1}{9n}\right) = (\sin n^2 + 2n + 5)e^{-n^2-n} + \frac{9n^2 + 5n + 6}{3n^2 + n + 1}. \text{ Then the value of } f(0) \text{ is}$$

- 1) 3 2) $1/2$ 3) 0 4) 2

75. **Statement 1:** The maximum value of $\left(\sqrt{-3+4x-x^2}+4\right)^2 + (x-5)^2$ (where $1 \leq x \leq 3$) is 36

Statement 2: The maximum distance between the point $(5, -4)$ and the point on the circle $(x-2)^2 + y^2 = 1$ is 6

- 1) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
 2) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
 3) Statement-1 is true, statement-2 is false.
 4) Statement-1 is false, statement-2 is true.

76. Let $f_1(x)$ and $f_2(x)$ be continuous and differentiable functions. If

$$f_1(0) = f_1(2) = f_1(4), f_1(1) + f_1(3) = f_2(0) = f_2(2) = f_2(4) = 0 \text{ and if}$$

$f_1(x) = 0$ and $f_2'(x) = 0$ do not have common root, then the minimum number of zeros of

$$f_1'(x)f_2'(x) + f_1(x)f_2^{(11)}(x) \text{ in } [0, 4], \text{ is}$$

- 1) 1 2) 2 3) 3 4) 4

77. $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos x}{1+e^x} dx$

- 1) 1 2) $\frac{\pi}{2}$ 3) π 4) 0

78. Let $f(x) = \lim_{n \rightarrow \infty} n^2 \left(x^{\frac{1}{n}} - x^{\frac{1}{n+1}} \right) : x > 0$ then $\int x \cdot f(x) dx =$

- 1) $\frac{x^2}{2} \log x + c$ 2) $\frac{x^2}{2} \log x - \frac{x^2}{2} + c$ 3) $\frac{x^2}{2} \log x + \frac{x^2}{2} + c$ 4) $\frac{x^2}{2} \log x - \frac{x^2}{4} + c$





79. Value of 'K' so that $f(k) = \int_0^4 |4x - x^2 - k| dx$ is minimum is
- 1) 3 2) 2 3) 25 4) 36

80. The value of the integral $\int_0^{\pi/2} \frac{1 + 2 \sin x}{(2 + \sin x)^2} dx$ is equal to
- 1) 2 2) 1 3) $\frac{1}{2}$ 4) 0

(NUMERICAL VALUE TYPE)

Section-II contains 10 Numerical Value Type questions. Attempt any 5 questions only. First 5 attempted questions will be considered if more than 5 questions attempted. The Answer should be within 0 to 9999. If the Answer is in Decimal then round off to the nearest Integer value (Example i.e. If answer is above 10 and less than 10.5 round off is 10 and If answer is from 10.5 and less than 11 round off is 11).

Marking scheme: +4 for correct answer, 0 if not attempt and -1 in all other cases.

81. Let $f(x)$ be periodic function with fundamental period 12. If $\int_{-6}^{12} f(x) dx = 9$ and

$$\int_0^{12} f(x) dx = 12 \text{ then the value of } \left| \int_0^6 f(x) dx \right|$$

82. The value of $\int_0^1 (1 - x^7)^{\frac{1}{4}} dx - \int_0^1 (1 - x^4)^{\frac{1}{7}} dx$

83. Number of values of x satisfying $\int_0^x t^2 \sin(x-t) dt = x^2$ in $[0, 100]$

84. If $I_1 = \int_0^1 (1 - x^{50})^{100} dx$, $I_2 = \int_0^1 (1 - x^{50})^{101} dx$ then $5050 \cdot \frac{I_1}{I_2} =$

85. Let $f(x)$ be a quadratic function such that $f(0) = 1$ and $\int \frac{f(x)}{x^2(x+1)^3} dx$ is a rational functions then $f'(0)$ is

86. Let $g(x) = |4x^3 - x| \cos \pi x$ then number of points where $g(x)$ is non differentiable in $(-\infty, \infty)$ is equal to





87. Let $P(x)$ be a real polynomial of degree 3 which vanishes at $x=3$. Let $P(x)$ have local minima at $x=1$, local maxima at $x=-1$ and $\int_{-2}^{+2} P(x)dx = 24$, then the absolute value of leading coefficient of the polynomial $P(x)$ is equal to _____
88. If f is differentiable function such that $f'(2) = \frac{1}{2}$ then $\lim_{h \rightarrow 0} \frac{f(2+h^4) - f(2-h^4)}{h^4}$ is equal to
89. If $f(2023) = \pi/2$ & $f(x) = f(2x) \forall x \in R$ & $f(x)$ is continuous for $x \in R$, then the value of $\lim_{x \rightarrow 0} \frac{5\cos(f(x)) - 3\sqrt{\cos f(x)}}{\sin^2 x + x^3}$
90. Let $f(x) = \frac{e^x - e^{-x}}{2}$ and if $f(g(x)) = x$ then the value of $g\left(\frac{e^{2022} - 1}{2e^{1011}}\right)$





Sri Chaitanya

Junior Colleges & Coaching Centres

Infinity
Learn BY SRI CHAITANYA



VAVILALA CHIDVILAS REDDY
HT.No. 236165088
Sri Chaitanya
6th - 12th Class

WITH ALL INDIA **RANK 1** IN JEE ADVANCED 2023
SRI CHAITANYA
STANDS AT THE TOP

SEIZES 5 RANKS IN TOP 10 IN ALL-INDIA OPEN CATEGORY

ANDHRA PRADESH STATE TOPPER



8th - 12th Class
Sri Chaitanya

2

RANK



AIITS Student

3

RANK



9th - 12th Class
Sri Chaitanya

4

RANK



4th - 12th Class
Sri Chaitanya

7

RANK

32 TOP RANKS BELOW 100 IN ALL-INDIA OPEN CATEGORY



RANK
12

DASARI SAKETH NAIDU
HT.No. 236060148



RANK
14

GUTHIKONDA ABHIRAM
HT.No. 236120121



RANK
17

P GNANA KOUSIK REDDY
HT.No. 236133520



RANK
18

D VENKATA YUGESH
HT.No. 236043359



RANK
19

MOULIK JINDAL
HT.No. 237021004



RANK
22

D PRATAP SINGH
HT.No. 232008139



RANK
23

SHAIK SAHIL CHANDA
HT.No. 236132870



RANK
26

SONI MAYANK
HT.No. 232097025



RANK
33

GOYAL NAMAN
HT.No. 232029345



RANK
37

SUMEDH SS
HT.No. 231028088



RANK
38

V DEERAJ SATHVIK REDDY
HT.No. 236067161



RANK
44

V VERGIYA KAUSHAL
HT.No. 231004058



RANK
46

SUTHAR HARSHAL
HT.No. 231005060



RANK
58

ABHI JAIN
HT.No. 232091038



RANK
61

RIDAM JAIN
HT.No. 232068021



RANK
62

SUTHAR HARSH
HT.No. 232100028



RANK
63

SOJITRA DIPEN
HT.No. 231017001



RANK
65

D PANENDRANADHA REDDY
HT.No. 236060298



RANK
67

GUNDA SUSHANTH
HT.No. 236129732



RANK
68

HARSH TAYA
HT.No. 237021178



RANK
69

KRISH GUPTA
HT.No. 232068189



RANK
73

KUNCHAKURTHY VIVEK
HT.No. 236134093



RANK
81

N DHARMA TEJA REDDY
HT.No. 236050138



RANK
84

S VENKAT KOUNDINYA
HT.No. 236130679



RANK
87

MRUNAL SHRIKANT
HT.No. 23104138



RANK
92

SAMOTA APURVA
HT.No. 232099210



RANK
97

N PRINCE BRANHAM REDDY
HT.No. 236064112

BELOW
20 >>
All India Open
Category Ranks

10
RANKS (50%)

BELOW
100 >>
All India Open
Category Ranks

32

BELOW
1000 >>
All India Open
Category Ranks

181

BELOW
100 >>
All India
Category Ranks
Count

89

BELOW
1000 >>
All India
Category Ranks
Count

699

NUMBER OF RANKS QUALIFIED
3,621

ADMISSIONS OPEN >> JEE ADVANCED LONG-TERM 2024

040 66 06 06 06

www.srichaitanya.net



Scan QR Code for
JEE Advanced Offers