

Grain Zero: residual overlap after structured Kakeya factorization in R³

Manuel Coleman

Love Labs LCA / PeAIce Research Program

coleman.manuel23@gmail.com

Scaffold preprint pack · RH OPEN · Coleman OPEN · ζ firewall active · h < 1 · export 2026-07-08

Proposed arXiv categories: math.CA (primary) · math.MG (cross-list, proposed)

MSC 2020: 42B25, 28A75, 42B20

Keywords: Kakeya set conjecture, residual measures, incidence geometry, multiplicity, Wang–Zahl reduction

Abstract

We isolate a residual overlap measure—Grain Zero—after a finite family of δ -tubes in R^3 has been factored into the structured carriers appearing in modern proofs of the Kakeya set conjecture in three dimensions (grains, slabs, prisms, planks, sticky packets, convex carriers, and scale envelopes). The raw excess multiplicity is not itself Grain Zero; Grain Zero is the excess supported off the extracted carrier. The primary theorem target is residual vanishing: after the Wang–Zahl / Guth–Wang–Zahl reduction stages, the normalized residual mass should tend to zero as $\delta \rightarrow 0$. Operator encodings and any zeta-regularization near $s = 0$ are second-stage and conditional. The Riemann hypothesis and the Coleman conjecture remain OPEN. No claim constitutes a proof of either statement. The value $\zeta(0) = -\frac{1}{2}$ is not asserted as a term in the Kakeya proof literature.

1. Introduction

STATUS: Program note ARX-001 under PeAIce claim discipline. Definition of $G^\blacksquare$ is FORMAL. Residual vanishing is PROPOSED / OPEN. Operator and zeta stages are explicitly second-stage.

A Kakeya (Besicovitch) set in $R^\blacksquare$ contains a unit line segment in every direction. In R^3 , every such set has Minkowski and Hausdorff dimension three (Wang–Zahl 2025; Guth–Wang–Zahl 2026). The proof architecture organizes directional overlap into structured carriers. This note does not reprove the Kakeya theorem. It asks a residual question: after legal structure is removed, what measure remains, and can it be shown to vanish under the same estimates that close the volume problem?

We call the residual object Grain Zero. The name is program-internal; the mathematics is a residual measure after carrier subtraction.

FIREWALL: $\zeta(0) = -\frac{1}{2}$ is not asserted as a Kakeya theorem term. Any regularization at $s = 0$ is second-stage and conditional on a residual operator or counting object constructed after residual control.

2. Tube setup

Let T_δ be a finite family of δ -tubes in R^3 with shading $Y(T) \subset T$ for each $T \in T_\delta$. Write:

$$\begin{aligned} U(T_\delta, Y) &:= \blacksquare_{T \in T_\delta} Y(T) \\ m_\delta(x) &:= \#\{ T \in T_\delta : x \in Y(T) \} \\ e_\delta(x) &:= (m_\delta(x) - 1)_+ \end{aligned}$$

The raw excess measure $e_\delta(x) dx$ still includes overlap that may be fully accounted for by structured carriers. Grain Zero begins only after that structure is extracted.

3. Structured carriers

Definition (Structured carrier, informal) — FORMAL (program definition); pinning OWED. At a scale passage $\delta \leq \rho \leq 1$, let $C_{\{\delta, \rho\}}$ denote the measurable structured carrier produced by a fixed stage of a Wang–Zahl / Guth–Wang–Zahl style reduction. Depending on the stage, $C_{\{\delta, \rho\}}$ may include grains, slabs, prisms, planks, convex carriers, sticky packets, Katz–Tao convex-density packets, Frostman slab packets, high-multiplicity cells, and scale-induction envelopes.

Remark: A rigorous implementation must pin $C_{\{\delta, \rho\}}$ to explicit source addresses in Wang–Zahl / Guth–Wang–Zahl for the chosen stage. Carrier pinning is an owed edit before arXiv submission.

4. Grain Zero

Definition DDATL-002.1 (Grain Zero residual measure) — FORMAL.

After $C_{\{\delta, \rho\}}$ has been extracted, define:

$$dG_{\{\delta, \rho\}}(x) := e_{\delta}(x) \cdot 1_{\{R^3 \setminus C_{\{\delta, \rho\}}\}}(x) \, dx$$

Equivalently, for measurable $A \subset R^3$:

$$G_{\{\delta, \rho\}}(A) := \int_A (m_{\delta}(x) - 1)_{+} \cdot 1_{\{R^3 \setminus C_{\{\delta, \rho\}}\}}(x) \, dx$$

$G_{\{\delta, \rho\}}$ is not a grain, slab, prism, or sticky packet. It is the leftover excess after those carriers absorb the overlap they are designed to control.

5. Primary theorem target

Conjecture (Residual vanishing) — PROPOSED / OPEN.

Under the hypotheses and refinements of the Wang–Zahl / Guth–Wang–Zahl reduction, after broad/narrow decomposition, convex-density factoring, grain/slab/prism extraction, sticky or non-sticky reduction, and final scale refinement:

$$G_{\{\delta, \rho\}}(R^3) / \sum_{T \in T_{\delta}} |Y(T)| \rightarrow 0 \text{ as } \delta \rightarrow 0$$

A weak form with $\delta^{o(1)}$ capture may be assemblable from existing error channels; a strong form is the program’s primary geometric burden.

PeAIce seal: No point in space can lie inside too many grains without the overlap becoming structure or becoming negligible.

6. Lemma ladder (program outline)

L1. Structured factorization — PROPOSED. Decompose $e_{\delta} \, dx = dS_{\{\delta, \rho\}} + dG_{\{\delta, \rho\}}$ with dS supported on carriers and $dG_{\{\delta, \rho\}}$ off carriers.

L2. Carrier exhaustion — PROPOSED. Show dS accounts for overlap controlled by existing incidence, multiplicity, convex-density, or slab estimates at the chosen stage.

L3. Residual domination — PROPOSED. Bound $G_{\{\delta, \rho\}}(R^3) \leq E_{bn} + E_{convex} + E_{sticky} + E_{scale}$.

L4. Residual vanishing — OPEN. Deduce the residual vanishing conjecture.

L5. Operator encoding — second stage, OPEN. Only after residual control, construct $K_{\{G_{\{\delta, \rho\}}\}}$, a kernel, a counting function, or a scale defect measure from residual data.

L6. Zeta-regularization question — conditional, OPEN. Only after an operator exists, ask whether a zeta/Mellin object continues near $s = 0$. Zeta firewall applies.

7. Obstacles

1. $C_{\{\delta,p\}}$ is proof-stage dependent; several residual lanes G^{grain} , G^{slab} , ... may be required.
2. Existingakeya proofs close by volume/multiplicity control without naming a residual measure; residual closure must be translated from those estimates.
3. A vanishing residual may be too thin to support a nontrivial spectral object; the correct second-stage object may be a renormalized defect along scales rather than G itself.

8. Related work and non-claims

Geometric background:akeya theorem in $\mathbb{R}^3$ (Wang–Zahl arXiv:2502.17655; Guth–Wang–Zahl arXiv:2601.14411) and the broader restriction/akeya complex. We do not assert a peer-reviewed literature bridge fromakeya residual measures to zeros of the Riemann zeta function; any such bridge is program-original and presently OPEN.

The Riemann hypothesis and the Coleman conjecture remain OPEN.

9. Acknowledgments

Research engineering assistance was used in preparing program materials. Mathematical claims and any future arXiv submission decision remain the author’s responsibility. Evaluator non-sovereignty ($h < 1$) is a governance constraint on the program, not a theorem of analytic number theory.

References (selected)

- Wang, H. and Zahl, J. Volume estimates for unions of convex sets, and theakeya set conjecture in three dimensions. arXiv:2502.17655, 2025.
- Guth, L., Wang, H., and Zahl, J. A streamlined proof of theakeya set conjecture in $\mathbb{R}^3$. arXiv:2601.14411, 2026.
- Bennett, J., Carbery, A., and Tao, T. On the multilinear restriction andakeya conjectures. Acta Math. 196 (2006), 261–302. arXiv:math/0509262.
- Guth, L. Large value estimates in number theory, harmonic analysis, and computer science. arXiv:2503.07410, 2025.

Export fromakeyalogic/arxiv · LaTeX sources authoritative for arXiv upload · human author only · receipts over reach