

KekeyaNeedleSet(Light(Basic)): a typed incidence object and two-layer decomposition

Manuel Coleman

Love Labs LCA / PeAIce Research Program

coleman.manuel23@gmail.com

Scaffold preprint pack · KNS(LB) ≠ RH proof · RH OPEN · h < 1 · export 2026-07-08

Proposed arXiv categories: math.CA / math.MG (proposed)

MSC 2020: 42B25, 28A75

Keywords: Kekeya sets, tubes, multiplicity, incidence geometry, typed objects

Abstract

We introduce a typed incidence object $\text{KekeyaNeedleSet}(\text{Light}(\text{Basic}))$ (abbreviated $\text{KNS}(\text{LB})$): a single throughput center with a full directional fan of unit segments in dimensions $d \in \{2,3\}$, together with its δ -tube discretization. We record elementary fan geometry and a two-layer separation between seen overlap statistics (multiplicity, clustering) and unseen center-action / placement declarations. Overlap functionals do not determine lawful placement and do not locate zeros of the Riemann zeta function. The Riemann hypothesis and the Coleman conjecture remain OPEN.

1. Introduction

STATUS: Program note ARX-002. Definitions and fan lemmas are FORMAL in stated scope. Any bridge to zeros is OPEN. Compute receipts from program probes are NUMERICS and not theorems of ζ .

Modern Kekeya proofs import a precise vocabulary of tubes, shadings, and multiplicity. This note isolates a measure-maximal configuration—the full fan through a center—as a typed object useful for separating incidence observables from placement declarations. It is not a proof of the Riemann hypothesis.

FIREWALL: Typed incidence objects do not locate zeros of ζ . Overlap measures are not placement measures on the critical line.

2. Light(Basic) and KekeyaNeedleSet

Definition 2.1 (Light(Basic)) — FORMAL. A $\text{Light}(\text{Basic})$ object in dimension $d \in \{2,3\}$ is a pair $L = (C, \Phi_C)$ with center $C \in \mathbb{R}^d$ and intensity $\Phi_C > 0$. The center is a declared throughput point in carrier space; it is not identified with the complex origin, nor with the value $\zeta(0)$.

Definition 2.2 (KekeyaNeedleSet) — FORMAL. For $L = (C, \Phi_C)$ and direction sphere $S^{\wedge\{d-1\}}$ (directions up to sign):

$$\text{KekeyaNeedleSet}(L) := \{ \sigma_\omega : \omega \in S^{\wedge\{d-1\}} \}, \quad \sigma_\omega = \{ C + t\omega : |t| \leq 1/2 \}$$

We write $\text{KNS}(\text{LB})$ for $\text{KekeyaNeedleSet}(\text{Light}(\text{Basic}))$.

δ -discretization: Let $\Omega_\delta \subset S^{\wedge\{d-1\}}$ be a maximal δ -separated set, so $|\Omega_\delta| \ll \delta^{-(d-1)}$, and let $T^\wedge C_\delta$ be the corresponding family of δ -tubes about the segments σ_ω with full shading $Y(T) = T$. Multiplicity μ , union U , and clustering quantities are imported from the Kekeya tube calculus (Guth–Wang–Zahl).

3. Fan geometry

Lemma 3.1 (Fan geometry) — FORMAL (stated scope). Let $T^\wedge C_\delta$ be as above with full shading.

- (a) $U(T^\wedge C_\delta) \supset B(C, 1/2)$ and $|U| \leq 1$.
- (b) Pointwise multiplicity satisfies $\mu(x) \leq (\delta + |x-C|)^{-(d-1)}$ for $|x-C| \leq 1/2$, hence $\sup_x \mu(x) \leq \delta^{-(d-1)}$ on the core $B(C, \delta)$.
- (c) Averaged multiplicity is ≤ 1 ; the configuration is the sticky, positive-measure extreme of the Kakeya family at this scale.

Proof sketch: (a) Every ray of length $1/2$ from C lies in some σ_ω up to δ ; total volume scales as $\delta^{-(d-1)} \cdot \delta^{d-1} \leq 1$. (b) $T_\omega \ni x$ forces direction within angular width $\leq \delta/|x-C|$; counting δ -caps yields the displayed decay. (c) Average the radial profile against the natural measure on annuli.

4. Two-layer separation

Proposition 4.1 (Overlap does not determine placement) — FORMAL (non-inference) / PROPOSED for full quantitative exhibits in final draft.

Incidence data $(\mu, \Delta_{\max}, U)$ generated by a tube configuration do not determine a Hilbert-space placement functional $\leq_{\text{off}}$ relative to a declared register A . In particular, high core multiplicity (glare) is compatible with total misplacement, and high average multiplicity without a common point is compatible with aligned placement.

Proof outline: (i) Incidence objects are geometric; placement requires a declared encoding space and register. (ii) Fix incidence data and vary the register so that a fixed encoding has arbitrary off-register mass. (iii) Fix encoding and rotate the register. Either freedom kills any map from incidence alone to $\leq_{\text{off}}$.

Remark (Non-inference to zeros): Any implication from multiplicity to zero placement on the critical line requires an explicit map. No such map is constructed here. Bounded square-difference coupling corridors in the companion operator program are recorded as CLOSED-NEGATIVE for determinant-level identification and do not supply the missing rung.

5. What remains open

1. Theorem-level lift of KNS(LB) beyond typed incidence geometry.
2. An explicit prime-carrying operator from placement data to zero location (the missing rung of the companion program).
3. Independent measurement obligations in the compute layer (program item CP-004) — OPEN, not a substitute for proofs.

RH OPEN · Coleman Conjecture OPEN · KNS(LB) $\neq$ RH proof.

6. Acknowledgments

Research engineering assistance was used in preparing program materials. All mathematical claims remain the author's responsibility under evaluator non-sovereignty ($h < 1$).

References (selected)

Wang, H. and Zahl, J. arXiv:2502.17655, 2025.

Guth, L., Wang, H., and Zahl, J. arXiv:2601.14411, 2026.

PeAIce program source: PEAICE-CLAUDEV6-KNS-LB-PAPER-001; claude-v6 wall registry.

