

CS492: Probabilistic Programming

Markov Chain

Monte Carlo

Hongseok Yang

KAIST

CS492: Probabilistic Programming

Markov Chain

Monte Carlo



Hongseok Yang  
KAIST

Really about: Metropolis-Hastings algorithm

```
(doquery :lmh induce-fn [ints2 outs2])
```

(doquery :lmh induce-fn [ints2 outs2])




Lightweight Metropolis Hastings algorithm\* (LMH).

\* Wingate, Stuhlmuller and Goodman's paper at AISTATS 2011

# Learning outcome

- Can explain Metropolis-Hastings algorithm.
- Can say when this algo. is correct.
- Can develop an instance of the algorithm.

# Good king Markov puzzle\*




Markov rules 10 islands.

$100i$  people live in island  $i$ .

\* Borrowed from McElreath's book "Statistical Rethinking"

# Good king Markov puzzle\*



Markov rules 10 islands.

$100i$  people live in island  $i$ .


King loves his people and wants to visit each island in proportion to its population size.

[Q] Find an algorithm. No scheduling nor bookkeeping. Can move adjacent islands only.

\* Borrowed from McElreath's book "Statistical Rethinking"

# Good king Markov puzzle\*

Bad!!



Markov rules 10 islands.

$100i$  people live in island  $i$ .


King loves his people and wants to visit each island in proportion to its population size.

[Q] Find an algorithm. **No scheduling nor bookkeeping.** Can move adjacent islands only.

\* Borrowed from McElreath's book "Statistical Rethinking"



# Good king Markov puzzle\*



Markov rules 10 islands.

$100i$  people live in island  $i$ .

$i \sim \text{discrete}(1, 2, \dots, 10)$ .  
Visit  $i$ .  
Repeat.


Bad!!

King loves his people and wants to visit each island in proportion to its population size.

[Q] Find an algorithm. No scheduling nor bookkeeping. **Can move adjacent islands only.**

\* Borrowed from McElreath's book "Statistical Rethinking"

# Good king Markov puzzle\*



Markov rules 10 islands.

$100i$  people live in island  $i$ .

King loves his people and wants to visit each island in proportion to its population size.

**[Q] Find an algorithm. No scheduling nor bookkeeping. Can move adjacent islands only.**

\* Borrowed from McElreath's book "Statistical Rethinking"

# Solution

$k_n$  — island that the king visits at step  $n$ .

Repeat the following steps:

1. Flip a coin with prob. 0.5. If head, pick next  $k'$  clockwise. If tail, use  $k'$  counterclockwise.
2.  $\alpha := \min(1, k'/k_n)$ .
3. Flip a coin with prob.  $\alpha$ . If head,  $k_{n+1} := k'$ . Otherwise,  $k_{n+1} := k_n$ .

# Solution

$k_n$  — island that the king visits at step  $n$ .

Repeat the following steps:

1. Flip a coin with prob. 0.5. If head, pick next  $k'$  clockwise. If tail, use  $k'$  counterclockwise.
2.  $\alpha := \min(1, k'/k_n)$ .
3. Flip a coin with prob.  $\alpha$ . If head,  $k_{n+1} := k'$ . Otherwise,  $k_{n+1} := k_n$ .

[Q] Why correct? What does correctness even mean?

Sequence by the algo.:  $k_1, k_2, \dots, k_n, \dots$

Corr. informally: Frequency represents probability.

Sequence by the algo.:  $k_1, k_2, \dots, k_n, \dots$

Corr. informally: Frequency represents probability.

Corr. formally: For any  $f : \{1, \dots, 10\} \rightarrow \mathbb{R}$ ,

$(\sum_{j \leq n} f(k_j))/n \longrightarrow \mathbb{E}_{p(i)}[f(i)]$  as  $n \longrightarrow \infty$  with prob. 1,  
where  $p(i)=i/55$ , target prob. for visiting island  $i$ .

Sequence by the algo.:  $k_1, k_2, \dots, k_n, \dots$

Corr. informally: Frequency represents probability.

Corr. formally: For any  $f : \{1, \dots, 10\} \rightarrow \mathbb{R}$ ,  
 $(\sum_{j \leq n} f(k_j))/n \longrightarrow \mathbb{E}_{p(i)}[f(i)]$  as  $n \longrightarrow \infty$  with prob. 1,  
where  $p(i) = i/55$ , target prob. for visiting island  $i$ .

Holds because 1) the random move of the algo.  
has  $p$  as **invariant**; 2) the algo. can **move between  
any two islands** in finitely many ( $>0$ ) steps.

Sequence by the algo.:  $k_1, k_2, \dots, k_n, \dots$

Corr. informally: Frequency represents probability.

Corr. formally: For any  $f : \{1, \dots, 10\} \rightarrow \mathbb{R}$ ,  
 $(\sum_{j \leq n} f(k_j))/n \longrightarrow \mathbb{E}_{p(i)}[f(i)]$  as  $n \longrightarrow \infty$  with prob. 1,  
where  $p(i) = i/55$ , target prob. for visiting island  $i$ .

Holds because 1) the random move of the algo. has  $p$  as invariant; 2) the algo. can move between any two islands in finitely many ( $>0$ ) steps.

[Q] Prove 1) and 2).



# Metropolis algorithm

Goal: Generate samples from target  $r(x)/Z$ , where  $Z = \int r(x)dx$ , the normalising constant.

Parameter: Conditional distribution  $q(x'|x)$ .

- Should be symmetric  $q(x'|x) = q(x|x')$ .
- Represents a random move.
- Called proposal kernel.

E.g.  $r(i)=i$ ,  $Z=55$ ,  $q(j|i)=0.5 \times [(j-i) \bmod 10 \in \{1,-1\}]$

# Metropolis algorithm

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

# Metropolis

Noisy greedy exploration.

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$   
 $\geq 1$  for better  $x'$   
 $< 1$  for worse  $x'$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0, 1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

May use  $\text{flip}(\alpha)$  instead, as in our sol. for King Markov

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q1] Does each step preserve  $r(x)/Z$  as invariant?

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q2] Posterior inference. Latent  $x \in \mathbb{R}^2$ . Observed  $y \in \mathbb{R}$

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(\mathbf{x})/Z$ . Symmetric proposal  $q(\mathbf{x}'|\mathbf{x})$ .

1. initialise  $\mathbf{x}_1$  randomly;  $n:=1$
2. repeat:
  - a)  $\mathbf{x}' \sim q(\mathbf{x}'|\mathbf{x}_n)$ ;  $\alpha := \min(1, r(\mathbf{x}')/r(\mathbf{x}_n))$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $\mathbf{x}_{n+1} := (\text{if } (u \leq \alpha) \text{ then } \mathbf{x}' \text{ else } \mathbf{x}_n)$ ;  $n:=n+1$

[Q2] Posterior inference. Latent  $\mathbf{x} \in \mathbb{R}^2$ . Observed  $y \in \mathbb{R}$



$$r(x) = p(y|x)p(x),$$

$$q(x'|x) = \text{normal}(x'_1|x_1, \varepsilon_1) \\ \times \text{normal}(x'_2|x_2, \varepsilon_2)$$

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0, 1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q2] Posterior inference. Latent  $x \in \mathbb{R}^2$ . Observed  $y \in \mathbb{R}$

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q3] How to instantiate this algo. for Anglican prog.?

# Metropolis

Noisy greedy exploration.  
No need to know  $Z$ .

Target  $r(x)/Z$ . Symmetric proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$
2. repeat:
  - a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, r(x')/r(x_n))$
  - b)  $u \sim \text{uniform}(0,1)$
  - c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q3] How to instantiate this algo. for Anglican prog.?

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}$ ,  $y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2$ ,  $y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2$ ,  $y \in \mathbb{R}^2$

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

Prob. distr. on result

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
                1  
                (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

Execute all sample exprs.  
Prob. distr. on all samples.



```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        b   (sample (normal (* x1 x1) 4))  
        x2 (if (> x1 0) 1 b)]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

Execute all sample exprs.  
Prob. distr. on all samples.

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

Prob. distr. on execution traces that record only sampled values.

```

(defquery q []
  (let [x1 (sample (normal 0 1))
        x2 (if (> x1 0)
                    1
                    (sample (normal (* x1 x1) 4)))])
    (observe (normal x2 1) 3)
    (observe (normal (+ x1 x2) 1) 5)
    x1)))

```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \textcolor{red}{\mathbb{R}} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

Prob. distr. on execution traces that record only sampled values.

```

(defquery q []
  (let [x1 (sample (normal 0 1))
        x2 (if (> x1 0)
                  1
                  (sample (normal (* x1 x1) 4)))]
    (observe (normal x2 1) 3)
    (observe (normal (+ x1 x2) 1) 5)
    x1)))

```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

Prob. distr. on **execution traces** that record only sampled values.

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

- All correct.
- (3) used for the design of MCMC for Anglican.

```
(defquery q []  
  (let [x1 (sample (normal 0 1))  
        x2 (if (> x1 0)  
              1  
              (sample (normal (* x1 x1) 4)))]  
    (observe (normal x2 1) 3)  
    (observe (normal (+ x1 x2) 1) 5)  
    x1)))
```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

- All correct.
- (3) used for the design of MCMC for Anglican.
- Difficult to find *symm. q.*



-1.8

```

(defquery q []
  (let [x1 (sample (normal 0 1))
        x2 (if (> x1 0)
              1
              (sample (normal (* x1 x1) 4)))]
    (observe (normal x2 1) 3)
    (observe (normal (+ x1 x2) 1) 5)
    x1)))

```

8.2

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

- All correct.
- (3) used for the design of MCMC for Anglican.
- Difficult to find **symm. q.**

```

(defquery q []
  (let [x1 (sample (normal 0 1))
        x2 (if (> x1 0)
                1
                (sample (normal (* x1 x1) 4)))]
    (observe (normal x2 1) 3)
    (observe (normal (+ x1 x2) 1) 5)
    x1)))

```

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

- All correct.
- (3) used for the design of MCMC for Anglican.
- Difficult to find **symm. q.**

0.9

```

(defquery q []
  (let [x1 (sample (normal 0 1))
        x2 (if (> x1 0)
                1
                (sample (normal (* x1 x1) 4)))]
    (observe (normal x2 1) 3)
    (observe (normal (+ x1 x2) 1) 5)
    x1)))

```

But q by re-execution is not symm.

[Q] Prior  $p(x)$  and likelihood  $p(y|x)$ . What are the types of  $x$  and  $y$ ?

(1)  $x \in \mathbb{R}, y \in \mathbb{R}^2$

(2)  $x \in \mathbb{R}^2, y \in \mathbb{R}^2$

(3)  $x \in \mathbb{R} \cup \mathbb{R}^2, y \in \mathbb{R}^2$

- All correct.
- (3) used for the design of MCMC for Anglican.
- Difficult to find symm. q.

# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n|x')}{r(x_n) \times q(x'|x_n)})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n); \quad n:=n+1$

# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n|x')}{r(x_n) \times q(x'|x_n)})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q1]  $q$  by re-execution for Anglican. What is  $q(x'|x)$ ?

# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n|x')}{r(x_n) \times q(x'|x_n)})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q1]  $q$  by re-execution for Anglican. What is  $q(x'|x)$ ?

[A1]  $q(x'|x) = p(x')$ .

# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n|x')}{r(x_n) \times q(x'|x_n)})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n); \quad n:=n+1$

[Q2] Noisy greedy exploration. Find a (relative) obj.

# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n|x')}{r(x_n) \times q(x'|x_n)})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n); \quad n:=n+1$

[Q2] Noisy greedy exploration. Find a (relative) obj.



# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ prop

When independent  
proposal is used.

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n)}{r(x_n) \times q(x')})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n); \quad n:=n+1$

[Q2] Noisy greedy exploration. Find a (relative) obj.

# Hastings Metropolis algorithm

Target  $r(x)/Z$ . ~~Symmetric~~ proposal  $q(x'|x)$ .

1. initialise  $x_1$  randomly;  $n:=1$

2. repeat:

a)  $x' \sim q(x'|x_n)$ ;  $\alpha := \min(1, \frac{r(x') \times q(x_n|x')}{r(x_n) \times q(x'|x_n)})$

b)  $u \sim \text{uniform}(0,1)$

c)  $x_{n+1} := (\text{if } (u \leq \alpha) \text{ then } x' \text{ else } x_n)$ ;  $n:=n+1$

[Q3] Does each step have  $r(x)/Z$  as invariant?

# Recap of the MH algo.

- Generate samples from unnormalised  $r(x)$ .  
No need to know  $Z = \int r(x)dx$ .
- Noisy greedy exploration using  $q(x'|x)$ .

# Guarantees informally

[Thm I] Each step of MH has  $r/Z$  as inv. dist.

# Guarantees informally

MH samples:  $x_1, x_2, x_3, \dots, x_n, \dots$

[Thm2] For all  $f: X \rightarrow \mathbb{R}$  with  $\mathbb{E}_{r(x)/Z}[f(x)]$  defined,

$\sum_{i \leq n} f(x_i)/n \longrightarrow \mathbb{E}_{r(x)/Z}[f(x)]$  as  $n \longrightarrow \infty$  with prob. 1,

if the MH with  $q$  is  $r/Z$ -irreducible.

# Guarantees informally

The estimate converges to the right value.

MH samples:  $x_1, x_2, x_3, \dots, x_n, \dots$

[Thm2] For all  $f: X \rightarrow \mathbb{R}$  with  $\mathbb{E}_{r(x)/Z}[f(x)]$  defined,

$\sum_{i \leq n} f(x_i)/n \longrightarrow \mathbb{E}_{r(x)/Z}[f(x)]$  as  $n \longrightarrow \infty$  with prob. 1,

if the MH with  $q$  is  $r/Z$ -irreducible.

# Guarantees informally

The estimate converges to the right value.

MH samples:  $x_1, x_2, x_3, \dots, x_n, \dots$

[Thm2] For all  $f: X \rightarrow \mathbb{R}$  with  $\mathbb{E}_{r(x)/Z}[f(x)]$  defined,

$\sum_{i \leq n} f(x_i)/n \longrightarrow \mathbb{E}_{r(x)/Z}[f(x)]$  as  $n \longrightarrow \infty$  with prob. 1,

if the MH with  $q$  is  $r/Z$ -irreducible.

MH moves well. For any  $x, x'$  with  $r(x') > 0$ , the MH can go from  $x$  to  $x'$  with non-zero prob.

# Guarantees informally

The estimate converges to the right value.

MH samples:  $x_1, x_2, x_3, \dots, x_n, \dots$

[Thm2] For all  $f: X \rightarrow \mathbb{R}$  with  $\mathbb{E}_{r(x)/Z}[f(x)]$  defined,

$\sum_{i \leq n} f(x_i)/n \longrightarrow \mathbb{E}_{r(x)/Z}[f(x)]$  as  $n \longrightarrow \infty$  with prob. 1,

if the MH with  $q$  is  $r/Z$ -irreducible.

MH moves well. For any  $x, x'$  with  $r(x') > 0$ , the MH can go from  $x$  to  $x'$  with non-zero prob.



# Guarantees informally

MH samples:  $x_1, x_2, x_3, \dots, x_n, \dots$

[Thm2] For all  $f: X \rightarrow \mathbb{R}$  with  $\mathbb{E}_{r(x)/Z}[f(x)]$  defined,

$\sum_{i \leq n} f(x_i)/n \longrightarrow \mathbb{E}_{r(x)/Z}[f(x)]$  as  $n \longrightarrow \infty$  with prob. 1,

if the MH with  $q$  is  $r/Z$ -irreducible.

# Guarantees informally

MH samples:  $x_1, x_2, x_3, \dots, x_n, \dots$

**[Thm2]** For all  $f: X \rightarrow \mathbb{R}$  with  $\mathbb{E}_{r(x)/Z}[f(x)]$  defined,

$\sum_{i \leq n} f(x_i)/n \longrightarrow \mathbb{E}_{r(x)/Z}[f(x)]$  as  $n \longrightarrow \infty$  with prob. 1,

if the MH with  $q$  is  $r/Z$ -irreducible.

Consequence of a general result in ergodic theory.  
Thm1 plays a crucial role in the proof.

# Reference

I looked at Chapters 5 and 6 of Robert & Casella's "Monte Carlo Statistical Methods".

Not recommended for general reading.

But details and pointers can be found there.