

P. Moon . D. E. Spencer

# FIELD THEORY HANDBOOK

Including Coordinate Systems, Differential  
Equations and Their Solutions

Second Edition

With 59 Figures

THE LIBRARY  
KING FAHD UNIVERSITY OF PETROLEUM & MINERALS  
DHARAN 31261, SAUDI ARABIA



Springer-Verlag Berlin Heidelberg New York 1971

PARRY MOON, Massachusetts Institute of Technology, Department of  
Electrical Engineering, Cambridge, MA 02139/USA  
DOMINA EBERLE SPENCER, University of Connecticut, Storrs, CT 06268/USA

QA  
401  
M 62

REF.

---

ISBN 3-540-02732-7 Springer-Verlag Berlin Heidelberg New York  
ISBN 0-387-02732-7 Springer-Verlag New York Heidelberg Berlin

---

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translation, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks

Under § 54 of the German Copyright Law where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher

© by Springer-Verlag, Berlin · Heidelberg 1961, 1971. Library of Congress Catalog Card Number 77-178288. Printed in Germany.  
The use of general descriptive names, trade names, trade marks, etc. in this publication, even if the former are not especially identified, is not to be taken as a sign that such names, as understood by the Trade Marks and Merchandise Marks Act, may accordingly be used freely by anyone.

## PREFACE

Let us first state exactly what this book is and what it is not. It is a compendium of equations for the physicist and the engineer working with electrostatics, magnetostatics, electric currents, electromagnetic fields, heat flow, gravitation, diffusion, optics, or acoustics. It tabulates the properties of 40 coordinate systems, states the Laplace and Helmholtz equations in each coordinate system, and gives the separation equations and their solutions. But it is not a textbook and it does not cover relativistic and quantum phenomena.

The history of classical physics may be regarded as an interplay between two ideas, the concept of *action-at-a-distance* and the concept of a *field*. Newton's equation of universal gravitation, for instance, implies action-at-a-distance. The same form of equation was employed by COULOMB to express the force between charged particles. AMPÈRE and GAUSS extended this idea to the phenomenological action between currents. In 1867, LUDVIG LORENZ formulated electrodynamics as retarded action-at-a-distance. At almost the same time, MAXWELL presented the alternative formulation in terms of fields.

In most cases, the field approach has shown itself to be the more powerful. A partial differential equation is solved, and boundary conditions are fitted to give a unique solution of the problem. The partial differential equations of classical physics, considered in this book, are the Laplace equation, Poisson equation, diffusion equation, scalar wave equation, and vector wave equation. Several methods of handling these equations are possible, but *separation of variables* is generally the most valuable. The procedure is as follows:

- (1) Transform the partial differential equation into the coordinate system that fits the geometry of the problem.
- (2) Separate this equation into three ordinary differential equations.
- (3) Obtain solutions of these ordinary differential equations.
- (4) Build up the unique solution that fits the boundary conditions, using as building blocks the particular solutions obtained in (3).

The amount of labor involved in solving a practical problem by this method, particularly when the required coordinates are unfamiliar, is rather formidable. One may well hesitate about embarking on such a program; and this hesitancy is probably responsible for the dearth of engineering solutions of field problems.

Most of the labor, however, occurs in the first three steps; and *these parts of the solution can be completed, once for all, and the results tabulated*. This is the purpose of the Handbook—to remove the routine drudgery from field solutions so that the scientist can concentrate on (4), the unique and important part of the work. No such tables have been available previously.

The range of problems that can be handled by separation of variables depends to a marked extent on the number of available coordinate systems. Accordingly, we have not limited ourselves to the eleven systems of EISENHART but have

chosen a number of others, bringing the total to 40. This by no means constitutes the totality of possible coordinates that allow simple separation or  $R$ -separation, though it includes most systems of reasonable simplicity and usefulness. Methods of obtaining further coordinates are explained in our book *Field Theory for Engineers* (D. Van Nostrand Co., Princeton, N. J., U.S.A., 1961).

For each of the 40 coordinate systems are given the relations to rectangular coordinates, the Stäckel matrix, metric coefficients, gradient, divergence, and curl, the Laplace and Helmholtz equations, the separation equations, and the solutions of these equations. Also listed are a tabulation of all the ordinary differential equations of field theory and their solutions, also a bibliography of works dealing with the mathematical functions involved.

Every equation has been checked independently by the two authors. But in a work that includes so many mathematical expressions, many of them given here for the first time, complete freedom from error would approach the miraculous. Any suggestions regarding errors or improvements will be greatly appreciated.

To a pure mathematician, our tabulations will seem ludicrously redundant. We have listed all special cases, even when they are obtainable from the basic form by a simple functional transformation. In this respect, the Handbook is similar to a table of integrals, whose practical value resides precisely in its redundancy.

We hope that the book will help the research worker in two ways:

- (a) By providing new coordinate systems, thus extending the range of engineering problems that can be handled by separation of variables, and
- (b) by freeing him from much of the annoyance and wasted effort usually associated with the routine part of the solution of partial differential equations.

The Authors

#### ACKNOWLEDGEMENT

We are grateful to B. G. TEUBNER, Leipzig, and to Dover Publications, New York, for permission to use Fig. 7.03, which was obtained from E. JAHNKE and F. EMDE, *Funktionentafeln mit Formeln und Kurven*.

## CONTENTS

| Section  | Page |
|--|------|
| I. Eleven coordinate systems . . . . .             | 1    |
| II. Transformations in the complex plane . . . . . | 49   |
| III. Cylindrical systems . . . . .                 | 77   |
| IV. Rotational systems . . . . .                   | 96   |
| V. The vector Helmholtz equation . . . . .         | 136  |
| VI. Differential equations . . . . .               | 144  |
| VII. Functions . . . . .                           | 163  |
| Bibliography . . . . .                             | 217  |
| Appendix. Symbols . . . . .                        | 226  |
| Author Index . . . . .                             | 229  |
| Subject Index . . . . .                            | 231  |



## Section I

### ELEVEN COORDINATE SYSTEMS

The book is limited to orthogonal coordinate systems in euclidean 3-space. Skew coordinate systems do not allow separation of variables and will not be considered.

One method of obtaining new coordinates—a method employed particularly by LAMÉ [1]\* and by DARBOUX [2]—is to study all the surfaces of a certain class and to determine what combinations of these surfaces will give orthogonal intersections. For surfaces of the *first* degree, we find only rectangular coordinates. Proceeding to equations of the *second* degree (and degenerate cases), we obtain the eleven coordinate systems of this section. EISENHART [3] has shown that all these systems allow simple separation of the Laplace and Helmholtz equations. Other coordinate systems, such as those built from fourth-degree surfaces [4], may have practical applications; but the eleven are undoubtedly the most important.

The use of esoteric coordinate systems may be of value, even in simple geometric considerations where field theory does not enter. With a spheroid, for instance, spheroidal coordinates eliminate the cumbersome mathematical expressions obtained with rectangular coordinates and allow the simple determination of areas and volumes.

But these unusual coordinate systems are particularly valuable in field theory. To express boundary conditions in a reasonably simple way, one must have coordinate surfaces that fit the physical boundaries of the problem [5]. In considering heat flow in a bar of elliptic cross section, for instance, one uses elliptic-cylinder coordinates; in calculating the effect of introducing a dielectric sphere into an electric field, one uses spherical coordinates; in obtaining the radiation from a slender spheroidal antenna, one uses prolate spheroid coordinates. Thus the range of field problems that can be handled effectively by an engineer or physicist will depend upon the number of coordinate systems with which he is familiar.

#### 1.01 METRIC COEFFICIENTS

An orthogonal coordinate system  $(u^1, u^2, u^3)$  may be designated by the metric coefficients  $g_{11}, g_{22}, g_{33}$ . An infinitesimal distance is written [5]

$$(ds)^2 = g_{11}(du^1)^2 + g_{22}(du^2)^2 + g_{33}(du^3)^2, \quad (1.01)$$

where

$$g_{ii} = \left( \frac{\partial x^1}{\partial u^i} \right)^2 + \left( \frac{\partial x^2}{\partial u^i} \right)^2 + \left( \frac{\partial x^3}{\partial u^i} \right)^2, \quad (1.02)$$

and  $x^i$  are rectangular coordinates.

\* See Bibliography at end of the book.

In parabolic coordinates, for instance, let  $u^1 = \mu$ ,  $u^2 = \nu$ ,  $u^3 = \psi$ . Then

$$\begin{cases} x^1 = \mu \nu \cos \psi, \\ x^2 = \mu \nu \sin \psi, \\ x^3 = \frac{1}{2}(\mu^2 - \nu^2). \end{cases}$$

The metric coefficients for parabolic coordinates are obtained by use of Eq. (1.02):

$$g_{11} = \left( \frac{\partial x^1}{\partial \mu} \right)^2 + \left( \frac{\partial x^2}{\partial \mu} \right)^2 + \left( \frac{\partial x^3}{\partial \mu} \right)^2 = (\nu \cos \psi)^2 + (\nu \sin \psi)^2 + \mu^2 = \mu^2 + \nu^2.$$

Similarly,

$$g_{22} = \mu^2 + \nu^2, \quad g_{33} = \mu^2 \nu^2.$$

Knowing the metric coefficients, one can easily write equations [5] for volume, gradient, curl, etc. Equation (1.01) shows that infinitesimal *distances* along the coordinate axes are  $(g_{11})^{\frac{1}{2}} d u^1$ ,  $(g_{22})^{\frac{1}{2}} d u^2$ ,  $(g_{33})^{\frac{1}{2}} d u^3$ . Thus an element of *area* on the  $u^1 u^2$ -surface is

$$d\mathcal{A} = [(g_{11})^{\frac{1}{2}} d u^1] [(g_{22})^{\frac{1}{2}} d u^2] = (g_{11} g_{22})^{\frac{1}{2}} d u^1 d u^2. \quad (1.03)$$

Similarly, an element of *volume* is

$$d\mathcal{V} = (g_{11} g_{22} g_{33})^{\frac{1}{2}} d u^1 d u^2 d u^3 = g^{\frac{1}{2}} d u^1 d u^2 d u^3. \quad (1.04)$$

For example, what is the area of a paraboloid of revolution ( $\mu = \mu_0$ ) from the vertex to a definite height designated by  $\nu_0$ ? Parabolic coordinates  $(\mu, \nu, \psi)$  are employed, with

$$g_{11} = g_{22} = \mu^2 + \nu^2, \quad g_{33} = \mu^2 \nu^2.$$

Thus, from Eq. (1.03),

$$d\mathcal{A} = \mu_0 \nu (\mu_0^2 + \nu^2)^{\frac{1}{2}} d\nu d\psi.$$

The total area is

$$\mathcal{A} = \int_0^{2\pi} \int_0^{\nu_0} \mu_0 \nu (\mu_0^2 + \nu^2)^{\frac{1}{2}} d\nu d\psi = \frac{2\pi \mu_0}{3} [(\mu_0^2 + \nu_0^2)^{\frac{3}{2}} - \mu_0^2].$$

*Gradient* in orthogonal curvilinear coordinates  $(u^1, u^2, u^3)$  is

$$\text{grad } \varphi = \frac{\mathbf{a}_1}{(g_{11})^{\frac{1}{2}}} \frac{\partial \varphi}{\partial u^1} + \frac{\mathbf{a}_2}{(g_{22})^{\frac{1}{2}}} \frac{\partial \varphi}{\partial u^2} + \frac{\mathbf{a}_3}{(g_{33})^{\frac{1}{2}}} \frac{\partial \varphi}{\partial u^3}, \quad (1.05)$$

where  $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$  are unit vectors.

*Divergence* is expressed as

$$\text{div } \mathbf{E} = g^{-\frac{1}{2}} \left\{ \frac{\partial}{\partial u^1} [(g/g_{11})^{\frac{1}{2}} E_1] + \frac{\partial}{\partial u^2} [(g/g_{22})^{\frac{1}{2}} E_2] + \frac{\partial}{\partial u^3} [(g/g_{33})^{\frac{1}{2}} E_3] \right\}. \quad (1.06)$$

*Curl* is

$$\begin{aligned} \text{curl } \mathbf{E} = g^{-\frac{1}{2}} & \left\{ \mathbf{a}_1 (g_{11})^{\frac{1}{2}} \left[ \frac{\partial}{\partial u^2} [(g_{33})^{\frac{1}{2}} E_3] - \frac{\partial}{\partial u^3} [(g_{22})^{\frac{1}{2}} E_2] \right] \right. \\ & + \mathbf{a}_2 (g_{22})^{\frac{1}{2}} \left[ \frac{\partial}{\partial u^3} [(g_{11})^{\frac{1}{2}} E_1] - \frac{\partial}{\partial u^1} [(g_{33})^{\frac{1}{2}} E_3] \right] \\ & \left. + \mathbf{a}_3 (g_{33})^{\frac{1}{2}} \left[ \frac{\partial}{\partial u^1} [(g_{22})^{\frac{1}{2}} E_2] - \frac{\partial}{\partial u^2} [(g_{11})^{\frac{1}{2}} E_1] \right] \right\}, \end{aligned} \quad (1.07)$$

or

$$\operatorname{curl} \mathbf{E} = \begin{vmatrix} \mathbf{a}_1(g_{11}/g)^{\frac{1}{2}} & \mathbf{a}_2(g_{22}/g)^{\frac{1}{2}} & \mathbf{a}_3(g_{33}/g)^{\frac{1}{2}} \\ \frac{\partial}{\partial u^1} & \frac{\partial}{\partial u^2} & \frac{\partial}{\partial u^3} \\ (g_{11})^{\frac{1}{2}} E_1 & (g_{22})^{\frac{1}{2}} E_2 & (g_{33})^{\frac{1}{2}} E_3 \end{vmatrix}. \quad (1.07a)$$

The *scalar Laplacian* of  $\varphi$  is defined as

$$\nabla^2 \varphi \equiv \operatorname{div} \operatorname{grad} \varphi. \quad (1.08)$$

In orthogonal curvilinear coordinates,

$$\nabla^2 \varphi = g^{-\frac{1}{2}} \sum_{i=1}^3 \frac{\partial}{\partial u^i} \left[ \frac{g^{\frac{1}{2}}}{g_{ii}} \frac{\partial \varphi}{\partial u^i} \right]. \quad (1.09)$$

As an example, write Laplace's equation in parabolic coordinates. Since

$$g_{11} = g_{22} = \mu^2 + \nu^2, \quad g_{33} = \mu^2 \nu^2, \quad g^{\frac{1}{2}} = \mu \nu (\mu^2 + \nu^2),$$

Eq. (1.09) gives

$$\nabla^2 \varphi = \frac{1}{\mu \nu (\mu^2 + \nu^2)} \left\{ \frac{\partial}{\partial \mu} \left[ \mu \nu \frac{\partial \varphi}{\partial \mu} \right] + \frac{\partial}{\partial \nu} \left[ \mu \nu \frac{\partial \varphi}{\partial \nu} \right] + \frac{\partial}{\partial \psi} \left[ \frac{(\mu^2 + \nu^2)}{\mu \nu} \frac{\partial \varphi}{\partial \psi} \right] \right\},$$

so Laplace's equation is

$$\frac{1}{\mu^2 + \nu^2} \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{1}{\mu} \frac{\partial \varphi}{\partial \mu} + \frac{\partial^2 \varphi}{\partial \nu^2} + \frac{1}{\nu} \frac{\partial \varphi}{\partial \nu} \right] + \frac{1}{\mu^2 \nu^2} \frac{\partial^2 \varphi}{\partial \psi^2} = 0.$$

The *vector Laplacian* of  $\mathbf{E}$  is defined as [6]

$$\star \star \mathbf{E} \equiv \operatorname{grad} \operatorname{div} \mathbf{E} - \operatorname{curl} \operatorname{curl} \mathbf{E}. \quad (1.10)$$

In orthogonal curvilinear coordinates  $(u^1, u^2, u^3)$ , the general expression is [5]

$$\star \star \mathbf{E} \equiv \left. \begin{aligned} \star \star \mathbf{E} &\equiv \mathbf{a}_1 \left\{ \frac{1}{(g_{11})^{\frac{1}{2}}} \frac{\partial \mathcal{T}}{\partial u^1} + (g_{11}/g)^{\frac{1}{2}} \left[ \frac{\partial \Gamma_2}{\partial u^3} - \frac{\partial \Gamma_3}{\partial u^2} \right] \right\} \\ &+ \mathbf{a}_2 \left\{ \frac{1}{(g_{22})^{\frac{1}{2}}} \frac{\partial \mathcal{T}}{\partial u^2} + (g_{22}/g)^{\frac{1}{2}} \left[ \frac{\partial \Gamma_3}{\partial u^1} - \frac{\partial \Gamma_1}{\partial u^3} \right] \right\} \\ &+ \mathbf{a}_3 \left\{ \frac{1}{(g_{33})^{\frac{1}{2}}} \frac{\partial \mathcal{T}}{\partial u^3} + (g_{33}/g)^{\frac{1}{2}} \left[ \frac{\partial \Gamma_1}{\partial u^2} - \frac{\partial \Gamma_2}{\partial u^1} \right] \right\}, \end{aligned} \right\} \quad (1.11)$$

where

$$\begin{aligned} \mathcal{T} &= g^{-\frac{1}{2}} \left\{ \frac{\partial}{\partial u^1} [(g/g_{11})^{\frac{1}{2}} E_1] + \frac{\partial}{\partial u^2} [(g/g_{22})^{\frac{1}{2}} E_2] + \frac{\partial}{\partial u^3} [(g/g_{33})^{\frac{1}{2}} E_3] \right\}, \\ \Gamma_1 &= \frac{g_{11}}{g^{\frac{1}{2}}} \left\{ \frac{\partial}{\partial u^2} [(g_{33})^{\frac{1}{2}} E_3] - \frac{\partial}{\partial u^3} [(g_{22})^{\frac{1}{2}} E_2] \right\}, \\ \Gamma_2 &= \frac{g_{22}}{g^{\frac{1}{2}}} \left\{ \frac{\partial}{\partial u^3} [(g_{11})^{\frac{1}{2}} E_1] - \frac{\partial}{\partial u^1} [(g_{33})^{\frac{1}{2}} E_3] \right\}, \\ \Gamma_3 &= \frac{g_{33}}{g^{\frac{1}{2}}} \left\{ \frac{\partial}{\partial u^1} [(g_{22})^{\frac{1}{2}} E_2] - \frac{\partial}{\partial u^2} [(g_{11})^{\frac{1}{2}} E_1] \right\}. \end{aligned}$$

## 1.02 DIFFERENTIAL EQUATIONS

The partial differential equations considered in this book are as follows:

(1) Laplace's equation,  $\nabla^2 \varphi = 0$ .

(2) Poisson's equation,  $\nabla^2 \varphi = -K(u^1, u^2, u^3)$ .

$$(3) \text{ The diffusion equation, } \nabla^2 \varphi = \frac{1}{h^2} \frac{\partial \varphi}{\partial t}.$$

$$(4) \text{ The wave equation, } \nabla^2 \varphi = \frac{1}{c^2} \frac{\partial^2 \varphi}{\partial t^2}.$$

$$(5) \text{ The damped wave equation, } \nabla^2 \varphi = \frac{1}{c^2} \frac{\partial^2 \varphi}{\partial t^2} + R \frac{\partial \varphi}{\partial t}.$$

$$(6) \text{ Transmission line equation, } \nabla^2 \varphi = \frac{1}{c^2} \frac{\partial^2 \varphi}{\partial t^2} + R \frac{\partial \varphi}{\partial t} + S \varphi.$$

$$(7) \text{ The vector wave equation, } \triangle \mathbf{E} = \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2}.$$

In electrical problems,  $\varphi$  represents the electric potential (volts); in magnetic problems,  $\varphi$  is the magnetic scalar potential (ampere-turns); in thermal problems,  $\varphi$  is the temperature (deg. cent); in gravitation,  $\varphi$  is the gravitational potential (joule kg<sup>-1</sup>); in vibration applications,  $\varphi$  is displacement (*m*); in hydrodynamics and acoustics,  $\varphi$  is the velocity potential (m<sup>2</sup> sec<sup>-1</sup>).

The solution of any of the scalar equations in the foregoing list may be reduced to a solution of the scalar *Helmholtz equation*,

$$\nabla^2 U + \kappa^2 U = 0. \quad (1.12)$$

For the *diffusion equation* (3), let

$$U(u^i) T(t),$$

where  $U$  is a function of the space coordinates and  $T$  is a function of time only. Substitution into the diffusion equation allows the separation of the time part, giving

$$\left. \begin{aligned} \nabla^2 U + \kappa^2 U &= 0, \\ \frac{dT}{dt} + \kappa^2 h^2 T &= 0, \end{aligned} \right\} \quad (1.13)$$

where  $\kappa$  is the separation constant.

Similarly, separation of the *wave equation* (4) gives

$$\left. \begin{aligned} \nabla^2 U + \kappa^2 U &= 0, \\ \frac{d^2 T}{dt^2} + \kappa^2 c^2 T &= 0. \end{aligned} \right\} \quad (1.14)$$

For the damped wave equation (5),

$$\left. \begin{aligned} \nabla^2 U + \kappa^2 U &= 0, \\ \frac{d^2 T}{dt^2} + R c^2 \frac{dT}{dt} + \kappa^2 c^2 T &= 0. \end{aligned} \right\} \quad (1.15)$$

With (6),

$$\left. \begin{aligned} \nabla^2 U + \kappa^2 U &= 0, \\ \frac{d^2 T}{dt^2} + R c^2 \frac{dT}{dt} + (S + \kappa^2) c^2 T &= 0. \end{aligned} \right\} \quad (1.16)$$

The solution of the Helmholtz equation depends on the space variables and the boundary conditions, and will be different for each problem. The equation in time, however, is independent of the coordinate system. Thus the solution of the diffusion equation (3) is always

$$\varphi = U(u^1, u^2, u^3) e^{-\kappa^2 h^2 t}, \quad (1.17)$$

and a particular solution of the wave equation (4) is

$$\varphi = U(u^1, u^2, u^3) \frac{\sin(\kappa c t)}{\cos(\kappa c t)}. \quad (1.18)$$

For the damped wave equation (5),

$$\varphi = U(u^1, u^2, u^3) e^{-\alpha t \pm [\kappa^2 - \kappa^2 c^2]^{\frac{1}{2}} t}, \quad (1.19)$$

where  $\alpha = R c^2 / 2$ . There are three cases:

- (a) *Overdamped* ( $\alpha > \kappa c$ ). The solution is given by Eq. (1.19).
- (b) *Oscillatory* ( $\alpha < \kappa c$ ). The solution may then be written

$$\varphi = U(u^1, u^2, u^3) [A e^{-\alpha t} \cos \omega t + B e^{-\alpha t} \sin \omega t], \quad (1.20)$$

where

$$\omega = [\kappa^2 c^2 - \alpha^2]^{\frac{1}{2}}.$$

- (c) *Critically damped* ( $\alpha = \kappa c$ ), where

$$\varphi = U(u^1, u^2, u^3) [A e^{-\alpha t} + B t e^{-\alpha t}]. \quad (1.21)$$

Evidently (6) is the same as (5) except that

$$\omega = [(S + \kappa^2) c^2 - \alpha^2]^{\frac{1}{2}}.$$

The Laplace equation (1) is, of course, merely a special case of the Helmholtz equation with  $\kappa = 0$ . The Poisson equation (2) can be reduced to the Laplace equation by a change of variable. Let

$$\varphi = \Omega + f(u^i),$$

where  $f(u^i)$  is so chosen that

$$\nabla^2 \Omega = 0. \quad (1.22)$$

Thus all the scalar equations mentioned in this section can be reduced to the Helmholtz equation (or its special case, the Laplace equation). In this way, the scalar fields of electrostatics, electric conduction, magnetism, heat flow, and acoustics can be based on the solutions of the Helmholtz and Laplace equations given in this book. The vector wave equation requires a somewhat different treatment, which is given in Section V.

### 1.03 SIMPLE SEPARATION

The solution of any of the partial differential equations, (1) to (6) of Section 1.02, reduces to a solution of the Helmholtz equation or the Laplace equation [7]. First consider the separation of the Helmholtz equation in 3-space.

The formulation is facilitated by the introduction of the Stäckel matrix [8]. With each coordinate system  $(u^1, u^2, u^3)$  is associated a matrix:

$$[S] = \begin{bmatrix} \Phi_{11}(u^1) & \Phi_{12}(u^1) & \Phi_{13}(u^1) \\ \Phi_{21}(u^2) & \Phi_{22}(u^2) & \Phi_{23}(u^2) \\ \Phi_{31}(u^3) & \Phi_{32}(u^3) & \Phi_{33}(u^3) \end{bmatrix}, \quad (1.23)$$

whose principal characteristic is that each row contains functions of only one variable (or constants). The Stäckel determinant is the determinant of the above

matrix:

$$S = \begin{vmatrix} \Phi_{11} & \Phi_{12} & \Phi_{13} \\ \Phi_{21} & \Phi_{22} & \Phi_{23} \\ \Phi_{31} & \Phi_{32} & \Phi_{33} \end{vmatrix}. \quad (1.24)$$

The cofactors of the elements in the first column are

$$M_{11} = \begin{vmatrix} \Phi_{22} & \Phi_{23} \\ \Phi_{32} & \Phi_{33} \end{vmatrix}, \quad M_{21} = - \begin{vmatrix} \Phi_{12} & \Phi_{13} \\ \Phi_{32} & \Phi_{33} \end{vmatrix}, \quad M_{31} = \begin{vmatrix} \Phi_{12} & \Phi_{13} \\ \Phi_{22} & \Phi_{23} \end{vmatrix}.$$

The necessary and sufficient conditions for simple separation of the scalar Helmholtz equation [9] are

$$\left. \begin{aligned} g_{ii} &= S/M_{i1}, \\ g^1/S &= f_1(u^1) \cdot f_2(u^2) \cdot f_3(u^3). \end{aligned} \right\} \quad (1.25)$$

The first requirement introduces the restriction that it be possible to form a Stäckel determinant [10] that is related to the metric coefficients as in Eq. (1.25). The second requirement states that  $g^1/S$  shall be a separable product. If these requirements are satisfied, the separation equations are

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \alpha_j \Phi_{ij} = 0. \quad (1.26)$$

Here  $i, j = 1, 2, 3$ ;  $\Phi_{ij}$  are elements of the Stäckel matrix, and  $\alpha_j$  are the separation constants with  $\alpha_1 = z^2$ .

In parabolic coordinates, for example, a few trials show that a possible form of Stäckel matrix is

$$[S] = \begin{bmatrix} \mu^2 & -1 & 1/\mu^2 \\ \nu^2 & 1 & 1/\nu^2 \\ 0 & 0 & 1 \end{bmatrix}.$$

The form is not unique: many equivalent forms would do just as well. From this matrix, we find

$$S = \mu^2 + \nu^2, \quad M_{11} = M_{21} = 1, \quad M_{31} = \frac{1}{\mu^2} + \frac{1}{\nu^2}.$$

Evidently, Eq. (1.25) is satisfied, and

$$f_1 = \mu, \quad f_2 = \nu, \quad f_3 = 1.$$

The separation equations are obtained from Eq. (1.26):

$$\left\{ \begin{aligned} \frac{1}{\mu} \frac{d}{d\mu} \left( \mu \frac{dM}{d\mu} \right) + M [\alpha_1 \mu^2 - \alpha_2 + \alpha_3/\mu^2] &= 0, \\ \frac{1}{\nu} \frac{d}{d\nu} \left( \nu \frac{dN}{d\nu} \right) + N [\alpha_1 \nu^2 + \alpha_2 + \alpha_3/\nu^2] &= 0, \\ \frac{d^2\Psi}{d\psi^2} + \Psi \alpha_3 &= 0. \end{aligned} \right.$$

These equations are solved for  $M, N, \Psi$ ; and the solution of the Helmholtz equation is

$$U(\mu, \nu, \psi) = M(\mu) \cdot N(\nu) \cdot \Psi(\psi).$$

By no means all orthogonal coordinate systems allow separation. In tangent-cylinder coordinates  $(\mu, \nu, \psi)$ , for instance,

$$g_{11} = g_{22} = (\mu^2 + \nu^2)^{-2}, \quad g_{33} = 1.$$

A few attempts will convince one that no Stäckel determinant can be devised that will satisfy Eq. (1.25).

The necessary and sufficient conditions for simple separation of the Laplace equation [9] are

$$\left. \begin{aligned} \frac{g_{ii}}{g_{jj}} &= \frac{M_{j1}}{M_{i1}}, \\ \frac{g^k}{g_{ii}} &= f_1 f_2 f_3 M_{i1}. \end{aligned} \right\} \quad (1.27)$$

The separation equations are again given by Eq. (1.26), but with  $\alpha_1 = 0$ .

For any *cylindrical* coordinate system in which the Helmholtz equation separates, the Stäckel matrix may be written [7]

$$[S] = \begin{bmatrix} 0 & \Phi_{12} & \Phi_{13} \\ 0 & \Phi_{22} & \Phi_{23} \\ 1 & 0 & 1 \end{bmatrix}. \quad (1.28)$$

For any *rotational* system (coordinate surfaces symmetrical about the  $z$ -axis),

$$[S] = \begin{bmatrix} \Phi_{11} & \Phi_{12} & \Phi_{13} \\ \Phi_{21} & \Phi_{22} & \Phi_{23} \\ 0 & 0 & 1 \end{bmatrix} \quad (1.29)$$

## 1.04 COORDINATE SYSTEMS

The eleven coordinate systems [3], formed from first and second degree surfaces, are as follows:

### Cylindrical

1. Rectangular coordinates  $(x, y, z)$ , Fig. 1.01.
2. Circular-cylinder coordinates  $(r, \psi, z)$ , Fig. 1.02.
3. Elliptic-cylinder coordinates  $(\eta, \psi, z)$ , Fig. 1.03.
4. Parabolic-cylinder coordinates  $(\mu, \nu, z)$ , Fig. 1.04.

### Rotational

5. Spherical coordinates  $(r, \theta, \psi)$ , Fig. 1.05.
6. Prolate spheroidal coordinates  $(\eta, \theta, \psi)$ , Fig. 1.06.
7. Oblate spheroidal coordinates  $(\eta, \theta, \psi)$ , Fig. 1.07.
8. Parabolic coordinates  $(\mu, \nu, \psi)$ , Fig. 1.08.

### General

9. Conical coordinates  $(r, \theta, \lambda)$ , Fig. 1.09.
10. Ellipsoidal coordinates  $(\eta, \theta, \lambda)$ , Fig. 1.10.
11. Paraboloidal coordinates  $(\mu, \nu, \lambda)$ , Fig. 1.11.

In the cylindrical systems, the cylindrical axis is always taken as the  $z$ -direction. In the rotational systems, the axis of symmetry is always taken as the  $z$ -axis and the angle about this axis is called  $\psi$ .

The Helmholtz and Laplace equations are simply separable in all these coordinate systems. The separation equations and their solutions are listed in the following tables.

As an example [11] of the use of the table, take a uniform electric field,

$$\varphi = E_0 z.$$

A metal prolate spheroid  $\eta = \eta_0$  at zero potential is now introduced into the field, the center of the spheroid being at the origin of coordinates. What is the resulting potential distribution?

Since the field has symmetry about the  $z$ -axis,  $\varphi$  is independent of  $\psi$  and the table gives as particular solutions of Laplace's equation,

$$\begin{aligned}\varphi &= P_p(\cosh \eta) P_p(\cos \theta), \\ \varphi &= Q_p(\cosh \eta) P_p(\cos \theta).\end{aligned}$$

The  $Q$ -functions of  $\cos \theta$  cannot be used because they become infinite on the  $z$ -axis. Assume a solution,

$$\varphi = A P_p(\cosh \eta) P_p(\cos \theta) + B Q_p(\cosh \eta) P_p(\cos \theta). \quad (1.30)$$

If this assumed solution does not satisfy the boundary conditions, we introduce additional terms, using infinite series if necessary.

Boundary conditions are

$$\begin{cases} \eta \gg \eta_0, & \varphi = E_0 z = E_0 a \sinh \eta \sin \theta; \\ \eta = \eta_0, & \varphi = 0. \end{cases}$$

When  $\eta \gg \eta_0$ ,  $Q_p(\cosh \eta) \rightarrow 0$  and

$$\varphi = E_0 a \sinh \eta \sin \theta = A P_p(\cosh \eta) P_p(\cos \theta).$$

Thus, to satisfy the first boundary condition,

$$A = E_0 a, \quad p = 1.$$

For the second boundary condition,

$$\varphi = 0 = [E_0 a \cosh \eta_0 + B Q_0(\cosh \eta_0)] \cos \theta$$

or

$$B = -\frac{E_0 a \cosh \eta_0}{Q_0(\cosh \eta_0)}.$$

Therefore, the unique solution of the problem is

$$\varphi = E_0 a \left[ \cosh \eta - \cosh \eta_0 \frac{Q_0(\cosh \eta)}{Q_0(\cosh \eta_0)} \right] \cos \theta. \quad (1.31)$$

The electric field strength is, according to the table,

$$\mathbf{E} = -\operatorname{grad} \varphi = -\frac{1}{a [\sinh^2 \eta + \sin^2 \theta]^{\frac{1}{2}}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} \right],$$

which is evaluated by differentiating Eq. (1.31). The field is plotted in Fig. 1.12. A similar plot for a dielectric spheroid is shown in Fig. 1.13.

**TABLE 1.01. RECTANGULAR COORDINATES  $(x, y, z)$** 

$$\begin{aligned} u^1 &= x, \quad -\infty < x < +\infty, \\ u^2 &= y, \quad -\infty < y < +\infty, \\ u^3 &= z, \quad -\infty < z < +\infty. \end{aligned}$$

Surfaces of constant  $x$ ,  $y$ , or  $z$  are mutually orthogonal planes (Fig. 1.01).

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} 0 & -1 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$S = 1,$$

$$M_{11} = M_{21} = M_{31} = 1.$$




Fig. 1.01. Rectangular coordinates  $(x, y, z)$ . Coordinate surfaces are the planes:  $x = \text{const}$ ,  $y = \text{const}$ ,  $z = \text{const}$

### Metric coefficients

$$g_{11} = g_{22} = g_{33} = 1, \quad g^1 = 1.$$

$$f_1 = f_2 = f_3 = 1.$$

### Important equations

$$(ds)^2 = (dx)^2 + (dy)^2 + (dz)^2.$$

$$\text{grad } \varphi = \mathbf{a}_x \frac{\partial \varphi}{\partial x} + \mathbf{a}_y \frac{\partial \varphi}{\partial y} + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\text{div } \mathbf{E} = \frac{\partial E_x}{\partial x} + \frac{\partial E_y}{\partial y} + \frac{\partial E_z}{\partial z}.$$

$$\text{curl } \mathbf{E} = \begin{vmatrix} \mathbf{a}_x & \mathbf{a}_y & \mathbf{a}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ E_x & E_y & E_z \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{\partial^2 \varphi}{\partial x^2} + \frac{\partial^2 \varphi}{\partial y^2} + \frac{\partial^2 \varphi}{\partial z^2}.$$

### SEPARATION OF LAPLACE'S EQUATION, $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = X(x)$ ,  $U^2 = Y(y)$ ,  $U^3 = Z(z)$ .

### General case

$$\begin{cases} \frac{d^2 X}{dx^2} - (\alpha_2 + \alpha_3) X = 0, \\ \frac{d^2 Y}{dy^2} + \alpha_2 Y = 0, \\ \frac{d^2 Z}{dz^2} + \alpha_3 Z = 0. \end{cases}$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2X}{dx^2} - (p^2 + q^2) X = 0, \quad \{04\}^* \quad X = A e^{(p^2+q^2)^{\frac{1}{2}}x} + B e^{-(p^2+q^2)^{\frac{1}{2}}x}.$$

$$\frac{d^2Y}{dy^2} + p^2 Y = 0, \quad \{04\} \quad Y = A \sin py + B \cos py.$$

$$\frac{d^2Z}{dz^2} + q^2 Z = 0, \quad \{04\} \quad Z = A \sin qz + B \cos qz.$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2X}{dx^2} - (p^2 - q^2) X = 0, \quad \{04\} \quad X = A e^{(p^2-q^2)^{\frac{1}{2}}x} + B e^{-(p^2-q^2)^{\frac{1}{2}}x}.$$

$$\frac{d^2Y}{dy^2} + p^2 Y = 0, \quad \{04\} \quad Y = A \sin py + B \cos py.$$

$$\frac{d^2Z}{dz^2} - q^2 Z = 0, \quad \{04\} \quad Z = A e^{qz} + B e^{-qz}.$$

For  $\varphi$  independent of  $z$ ,

$$\begin{cases} \frac{d^2X}{dx^2} - \alpha_2 X = 0, \\ \frac{d^2Y}{dy^2} + \alpha_2 Y = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$\frac{d^2X}{dx^2} - p^2 X = 0, \quad \{04\} \quad X = A e^{px} + B e^{-px}.$$

$$\frac{d^2Y}{dy^2} + p^2 Y = 0, \quad \{04\} \quad Y = A \sin py + B \cos py.$$

If  $\alpha_2 = -p^2$ ,

$$\frac{d^2X}{dx^2} + p^2 X = 0, \quad \{04\} \quad X = A \sin px + B \cos px.$$

$$\frac{d^2Y}{dy^2} - p^2 Y = 0, \quad \{04\} \quad Y = A e^{py} + B e^{-py}.$$

If  $\alpha_2 = 0$ ,

$$\frac{d^2X}{dx^2} = \frac{d^2Y}{dy^2} = 0, \quad \{04\} \quad X = A + Bx, \quad Y = A + By.$$

For  $\varphi$  independent of  $y$  and  $z$ ,

$$\frac{d^2\varphi}{dx^2} = 0, \quad \{04\} \quad \varphi = A + Bx.$$

SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2\varphi + \kappa^2\varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = X(x)$ ,  $U^2 = Y(y)$ ,  $U^3 = Z(z)$ .

\* The numbers in brackets designate the differential equation in terms of its singularities (see Section VI).

**General case**

$$\begin{cases} \frac{d^2X}{dx^2} - (\alpha_2 + \alpha_3) X = 0, \\ \frac{d^2Y}{dy^2} + \alpha_2 Y = 0, \\ \frac{d^2Z}{dz^2} + (\kappa^2 + \alpha_3) Z = 0. \end{cases}$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2X}{dx^2} - (p^2 + q^2) X = 0, \quad \{04\} \quad X = A e^{(p^2+q^2)^{\frac{1}{2}}x} + B e^{-(p^2+q^2)^{\frac{1}{2}}x}.$$

$$\frac{d^2Y}{dy^2} + p^2 Y = 0, \quad \{04\} \quad Y = A \sin px + B \cos px.$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 + q^2) Z = 0, \quad \{04\} \quad Z = A \sin [(\kappa^2 + q^2)^{\frac{1}{2}}z] + B \cos [(\kappa^2 + q^2)^{\frac{1}{2}}z].$$

If  $\alpha_2 = -p^2$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2X}{dx^2} + (p^2 + q^2) X = 0, \quad \{04\} \quad X = A \sin [(p^2 + q^2)^{\frac{1}{2}}x] + B \cos [(p^2 + q^2)^{\frac{1}{2}}x].$$

$$\frac{d^2Y}{dy^2} - p^2 Y = 0, \quad \{04\} \quad Y = A e^{px} + B e^{-px}.$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 - q^2) Z = 0, \quad \{04\} \quad Z = A \sin [(\kappa^2 - q^2)^{\frac{1}{2}}z] + B \cos [(\kappa^2 - q^2)^{\frac{1}{2}}z].$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2X}{dx^2} = \frac{d^2Y}{dy^2} = 0, \quad \{04\} \quad X = A + Bx, \quad Y = A + By.$$

$$\frac{d^2Z}{dz^2} + \kappa^2 Z = 0, \quad \{04\} \quad Z = A \sin \kappa z + B \cos \kappa z.$$

**For  $\varphi$  independent of  $z$** 

$$\begin{cases} \frac{d^2X}{dx^2} - \alpha_2 X = 0, \\ \frac{d^2Y}{dy^2} + (\kappa^2 + \alpha_2) Y = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$\frac{d^2X}{dx^2} - p^2 X = 0, \quad \{04\} \quad X = A e^{px} + B e^{-px}.$$

$$\frac{d^2Y}{dy^2} + (\kappa^2 + p^2) Y = 0, \quad \{04\} \quad Y = A \sin [(\kappa^2 + p^2)^{\frac{1}{2}}y] + B \cos [(\kappa^2 + p^2)^{\frac{1}{2}}y].$$

If  $\alpha_2 = -p^2$ ,

$$\frac{d^2X}{dx^2} + p^2 X = 0, \quad \{04\} \quad X = A \sin px + B \cos px.$$

$$\frac{d^2Y}{dy^2} + (\kappa^2 - p^2) Y = 0, \quad \{04\} \quad Y = A \sin [(\kappa^2 - p^2)^{\frac{1}{2}}y] + B \cos [(\kappa^2 - p^2)^{\frac{1}{2}}y].$$

If  $\alpha_2 = 0$ ,

$$\frac{d^2X}{dx^2} = 0, \quad \{04\} \quad X = A + Bx.$$

$$\frac{d^2Y}{dy^2} + \kappa^2 Y = 0, \quad \{04\} \quad Y = A \sin \kappa y + B \cos \kappa y.$$

**For  $\varphi$  independent of  $y$  and  $z$ ,**

$$\frac{d^2\varphi}{dx^2} + \kappa^2 \varphi = 0, \quad \{04\} \quad \varphi = A \sin \kappa x + B \cos \kappa x.$$


TABLE 1.02. CIRCULAR-CYLINDER COORDINATES  $(r, \psi, z)$ 

Fig. 1.02. Circular-cylinder coordinates  $(r, \psi, z)$ . Coordinate surfaces are circular cylinders ( $r = \text{const}$ ), half-planes ( $\psi = \text{const}$ ) intersecting on the  $z$ -axis, and parallel planes ( $z = \text{const}$ )

$$\begin{aligned} u^1 &= r, & 0 \leq r < \infty, \\ u^2 &= \psi, & 0 \leq \psi < 2\pi, \\ u^3 &= z. & -\infty < z < +\infty. \end{aligned}$$

$$\begin{cases} x = r \cos \psi, \\ y = r \sin \psi, \\ z = z. \end{cases}$$

Coordinate surfaces are

$$\begin{cases} x^2 + y^2 = r^2 \\ (\text{circular cylinders}, \\ r = \text{const}), \\ \tan \psi = y/x \\ (\text{half-planes}, \psi = \text{const}), \\ z = \text{const} \\ (\text{planes}). \end{cases}$$

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} 0 & -1/r^2 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

$$S = 1, \quad M_{11} = 1, \quad M_{21} = 1/r^2, \quad M_{31} = 1.$$

Metric coefficients

$$g_{11} = 1, \quad g_{22} = r^2, \quad g_{33} = 1, \quad g^{\frac{1}{2}} = r.$$

$$f_1 = r, \quad f_2 = 1, \quad f_3 = 1.$$

Important equations

$$(ds)^2 = (dr)^2 + r^2(d\psi)^2 + (dz)^2.$$

$$\text{grad } \varphi = \mathbf{a}_r \frac{\partial \varphi}{\partial r} + \frac{\mathbf{a}_\psi}{r} \frac{\partial \varphi}{\partial \psi} + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\text{div } \mathbf{E} = \frac{\partial E_r}{\partial r} + \frac{E_r}{r} + \frac{1}{r} \frac{\partial E_\psi}{\partial \psi} + \frac{\partial E_z}{\partial z}.$$

$$\text{curl } \mathbf{E} = \frac{1}{r} \begin{vmatrix} \mathbf{a}_r & \mathbf{a}_\psi r & \mathbf{a}_z \\ \frac{\partial}{\partial r} & \frac{\partial}{\partial \psi} & \frac{\partial}{\partial z} \\ E_r & E_\psi r & E_z \end{vmatrix}.$$

$$V^2 \varphi = \frac{\partial^2 \varphi}{\partial r^2} + \frac{1}{r} \frac{\partial \varphi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \varphi}{\partial \psi^2} + \frac{\partial^2 \varphi}{\partial z^2}.$$

ALTERNATIVE CIRCULAR-CYLINDER SYSTEM  $(\xi, \psi, z)$ 

Let  $r = e^\xi$ . Then

$$\begin{aligned} u^1 &= \xi, & -\infty < \xi < +\infty, \\ u^2 &= \psi, & 0 \leq \psi < 2\pi, \\ u^3 &= z. & -\infty < z < +\infty. \end{aligned}$$

$$\begin{cases} x = e^\xi \cos \psi, \\ y = e^\xi \sin \psi, \\ z = z. \end{cases}$$

The Stäckel matrix is

$$[S] = \begin{bmatrix} 0 & -1 & -e^{-2\xi} \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

$$S = e^{2\xi}, \quad M_{11} = M_{21} = 1, \quad M_{31} = e^{2\xi}.$$

## Metric coefficients

$$g_{11} = g_{22} = e^{2\xi}, \quad g_{33} = 1, \quad g^1 = e^{2\xi}.$$

$$f_1 = f_2 = f_3 = 1.$$

## Important equations

$$(ds)^2 = e^{2\xi} [(d\xi)^2 + (d\psi)^2] + (dz)^2.$$

$$\text{grad } \varphi = e^{-\xi} \left[ \mathbf{a}_\xi \frac{\partial \varphi}{\partial \xi} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} + \mathbf{a}_z \frac{\partial \varphi}{\partial z} \right].$$

$$\text{div } \mathbf{E} = e^{-\xi} \left[ \frac{\partial E_\xi}{\partial \xi} + E_\xi + \frac{\partial E_\psi}{\partial \psi} \right] + \frac{\partial E_z}{\partial z}.$$

$$\text{curl } \mathbf{E} = e^{-2\xi} \begin{vmatrix} \mathbf{a}_\xi e^\xi & \mathbf{a}_\psi e^\xi & \mathbf{a}_z \\ \frac{\partial}{\partial \xi} & \frac{\partial}{\partial \psi} & \frac{\partial}{\partial z} \\ E_\xi e^\xi & E_\psi e^\xi & E_z \end{vmatrix}.$$

$$\nabla^2 \varphi = e^{-2\xi} \left\{ \frac{\partial^2 \varphi}{\partial \xi^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right\} + \frac{\partial^2 \varphi}{\partial z^2}.$$

SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = R(r)$ ,  $U^2 = \Psi(\psi)$ ,  $U^3 = Z(z)$ .

## General case

$$\begin{cases} \frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - \left( \frac{\alpha_2}{r^2} + \alpha_3 \right) R = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_2 \Psi = 0, \\ \frac{d^2 Z}{dz^2} + \alpha_3 Z = 0. \end{cases}$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - (q^2 + p^2/r^2) R = 0, \quad \{24\} \quad R = A\mathcal{J}_p(iqr) + B\mathcal{I}_{-p}(iqr)$$

or \*  $R = A\mathcal{J}_n(iqr) + B\mathcal{Y}_n(iqr).$

$$\frac{d^2\Psi}{d\psi^2} + p^2\Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

$$\frac{d^2Z}{dz^2} + q^2Z = 0, \quad \{04\} \quad Z = A \sin qz + B \cos qz.$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + (q^2 - p^2/r^2) R = 0, \quad \{24\} \quad R = A\mathcal{J}_p(qr) + B\mathcal{I}_{-p}(qr)$$

or  $R = A\mathcal{J}_n(qr) + B\mathcal{Y}_n(qr).$

$$\frac{d^2\Psi}{d\psi^2} + p^2\Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

$$\frac{d^2Z}{dz^2} - q^2Z = 0, \quad \{04\} \quad Z = A e^{qz} + B e^{-qz}.$$

If  $\alpha_2 = 0$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - q^2R = 0, \quad \{14\} \quad R = A\mathcal{J}_0(iqr) + B\mathcal{Y}_0(iqr).$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B\psi.$$

$$\frac{d^2Z}{dz^2} + q^2Z = 0, \quad \{04\} \quad Z = A \sin qz + B \cos qz.$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} = 0, \quad \{04\} \quad R = A + B \ln r.$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B\psi.$$

$$\frac{d^2Z}{dz^2} = 0, \quad \{04\} \quad Z = A + Bz.$$

For  $\varphi$  independent of  $z$ ,

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - \frac{\alpha_2}{r^2} R = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_2 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - \frac{p^2}{r^2} R = 0, \quad \{04\} \quad R = A r^p + B r^{-p}.$$

$$\frac{d^2\Psi}{d\psi^2} + p^2\Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

If  $\alpha_2 = 0$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} = 0, \quad \{04\} \quad R = A + B \ln r.$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B\psi.$$

\* If  $p$  is an integer, the second form must be employed.

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - \alpha_3 R = 0, \\ \frac{d^2Z}{dz^2} + \alpha_3 Z = 0. \end{cases}$$

If  $\alpha_3 = q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - q^2 R = 0, \quad \{14\} \quad R = A J_0(iqr) + B Y_0(iqr).$$

$$\frac{d^2Z}{dz^2} + q^2 Z = 0, \quad \{04\} \quad Z = A \sin qz + B \cos qz.$$

If  $\alpha_3 = -q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + q^2 R = 0, \quad \{14\} \quad R = A J_0(qr) + B Y_0(qr).$$

$$\frac{d^2Z}{dz^2} - q^2 Z = 0, \quad \{04\} \quad Z = A e^{qz} + B e^{-qz}.$$

If  $\alpha_3 = 0$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} = 0, \quad \{04\} \quad R = A + B \ln r.$$

$$\frac{d^2Z}{dz^2} = 0, \quad \{04\} \quad Z = A + Bz.$$

For  $\varphi$  independent of  $r$  and  $z$ ,

$$\frac{d^2\varphi}{d\psi^2} = 0, \quad \{04\} \quad \varphi = A + B\psi.$$

For  $\varphi$  independent of  $\psi$  and  $z$ ,

$$\frac{d^2\varphi}{dr^2} + \frac{1}{r} \frac{d\varphi}{dr} = 0, \quad \{04\} \quad \varphi = A + B \ln r.$$

**SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2\varphi + \kappa^2\varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = R(r)$ ,  $U^2 = \Psi(\psi)$ ,  $U^3 = Z(z)$ .

General case

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - \left( \frac{\alpha_2}{r^2} + \alpha_3 \right) R = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_2 \Psi = 0, \\ \frac{d^2Z}{dz^2} + (\kappa^2 + \alpha_3) Z = 0. \end{cases}$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - (q^2 + p^2/r^2) R = 0, \quad \{24\} \quad R = A J_p(iqr) + B J_{-p}(iqr) \\ \text{or} \quad R = A J_n(iqr) + B Y_n(iqr).$$

$$\frac{d^2\Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 + q^2) Z = 0, \quad \{04\} \quad Z = A \sin [(\kappa^2 + q^2)^{1/2} z] \\ + B \cos [(\kappa^2 + q^2)^{1/2} z].$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + (q^2 - p^2/r^2) R = 0, \quad \begin{cases} 24 \end{cases} \quad R = A\mathcal{I}_p(qr) + B\mathcal{I}_{-p}(qr) \\ \text{or} \quad R = A\mathcal{I}_n(qr) + B\mathcal{Y}_n(qr). \end{math>$$

$$\frac{d^2\Psi}{d\psi^2} + p^2\Psi = 0, \quad \begin{cases} 04 \end{cases} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 - q^2) Z = 0, \quad \begin{cases} 04 \end{cases} \quad Z = A \sin [(\kappa^2 - q^2)^{\frac{1}{2}}z] \\ + B \cos [(\kappa^2 - q^2)^{\frac{1}{2}}z]. \end{math>$$

If  $\alpha_2 = 0$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - q^2 R = 0, \quad \begin{cases} 14 \end{cases} \quad R = A\mathcal{I}_0(iqr) + B\mathcal{Y}_0(iqr). \end{math>$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \begin{cases} 04 \end{cases} \quad \Psi = A + B\psi.$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 + q^2) Z = 0, \quad \begin{cases} 04 \end{cases} \quad Z = A \sin [(\kappa^2 + q^2)^{\frac{1}{2}}z] \\ + B \cos [(\kappa^2 + q^2)^{\frac{1}{2}}z]. \end{math>$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} = 0, \quad \begin{cases} 04 \end{cases} \quad R = A + B \ln r.$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \begin{cases} 04 \end{cases} \quad \Psi = A + B\psi.$$

$$\frac{d^2Z}{dz^2} + \kappa^2 Z = 0, \quad \begin{cases} 04 \end{cases} \quad Z = A \sin \kappa z + B \cos \kappa z.$$

For  $\varphi$  independent of  $z$ ,

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + (\kappa^2 - \alpha_2/r^2) R = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_2 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + (\kappa^2 - p^2/r^2) R = 0, \quad \begin{cases} 24 \end{cases} \quad R = A\mathcal{I}_p(\kappa r) + B\mathcal{I}_{-p}(\kappa r) \\ \text{or} \quad R = A\mathcal{I}_n(\kappa r) + B\mathcal{Y}_n(\kappa r). \end{math>$$

$$\frac{d^2\Psi}{d\psi^2} + p^2\Psi = 0, \quad \begin{cases} 04 \end{cases} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

If  $\alpha_2 = 0$ ,

$$\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \kappa^2 R = 0, \quad \begin{cases} 14 \end{cases} \quad R = A\mathcal{I}_0(\kappa r) + B\mathcal{Y}_0(\kappa r).$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \begin{cases} 04 \end{cases} \quad \Psi = A + B\psi.$$

For  $\varphi$  independent of  $\Psi$ ,

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - \alpha_3 R = 0, \\ \frac{d^2Z}{dz^2} + (\kappa^2 + \alpha_3) Z = 0. \end{cases}$$

$$\begin{aligned}
 & \text{If } \alpha_3 = q^2, \\
 & \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} - q^2 R = 0, \quad \{14\} \quad R = A \mathcal{J}_0(iqr) + B \mathcal{Y}_0(iqr). \\
 & \frac{d^2Z}{dz^2} + (\kappa^2 + q^2) Z = 0, \quad \{04\} \quad Z = A \sin[(\kappa^2 + q^2)^{\frac{1}{2}} z] \\
 & \quad + B \cos[(\kappa^2 + q^2)^{\frac{1}{2}} z]. \\
 & \text{If } \alpha_3 = -q^2, \\
 & \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + q^2 R = 0, \quad \{14\} \quad R = A \mathcal{J}_0(qr) + B \mathcal{Y}_0(qr). \\
 & \frac{d^2Z}{dz^2} + (\kappa^2 - q^2) Z = 0, \quad \{04\} \quad Z = A \sin[(\kappa^2 - q^2)^{\frac{1}{2}} z] \\
 & \quad + B \cos[(\kappa^2 - q^2)^{\frac{1}{2}} z]. \\
 & \text{If } \alpha_3 = 0, \\
 & \frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} = 0, \quad \{04\} \quad R = A + B \ln r. \\
 & \frac{d^2Z}{dz^2} + \kappa^2 Z = 0, \quad \{04\} \quad Z = A \sin \kappa z + B \cos \kappa z.
 \end{aligned}$$

For  $\varphi$  independent of  $\psi$  and  $z$ ,

$$\frac{d^2\varphi}{dr^2} + \frac{1}{r} \frac{d\varphi}{dr} + \kappa^2 \varphi = 0, \quad \{14\} \quad \varphi = A \mathcal{J}_0(\kappa r) + B \mathcal{Y}_0(\kappa r).$$

TABLE 1.03. ELLIPTIC-CYLINDER COORDINATES  $(\eta, \psi, z)$

$$\begin{aligned}
 u^1 &= \eta, \quad 0 \leq \eta < \infty, \\
 u^2 &= \psi, \quad 0 \leq \psi < 2\pi, \\
 u^3 &= z, \quad -\infty < z < +\infty. \\
 &\left\{ \begin{array}{l} x = a \cosh \eta \cos \psi, \\ y = a \sinh \eta \sin \psi, \\ z = z. \end{array} \right.
 \end{aligned}$$

Coordinate surfaces are confocal cylinders and planes:

$$\left\{ \begin{array}{l} \left( \frac{x}{a \cosh \eta} \right)^2 + \left( \frac{y}{a \sinh \eta} \right)^2 = 1 \\ \quad (\text{elliptic cylinders, } \eta = \text{const}), \\ \left( \frac{x}{a \cos \psi} \right)^2 - \left( \frac{y}{a \sin \psi} \right)^2 = 1 \\ \quad (\text{hyperbolic cylinders, } \psi = \text{const}), \\ z = \text{const} \\ \quad (\text{planes}). \end{array} \right.$$




Fig. 1.03. Elliptic-cylinder coordinates  $(\eta, \psi, z)$ . Coordinate surfaces are elliptic cylinders ( $\eta = \text{const}$ ), hyperbolic cylinders ( $\psi = \text{const}$ ), and planes ( $z = \text{const}$ )

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} 0 & -1 & -a^2 \cosh^2 \eta \\ 0 & 1 & a^2 \cos^2 \psi \\ 1 & 0 & 1 \end{bmatrix}.$$

$$S = a^2 (\cosh^2 \eta - \cos^2 \psi), \quad M_{11} = M_{21} = 1, \quad M_{31} = a^2 (\cosh^2 \eta - \cos^2 \psi).$$

**Metric coefficients**

$$g_{11} = g_{22} = a^2(\cosh^2 \eta - \cos^2 \psi), \quad g_{33} = 1,$$

$$g^{\frac{1}{2}} = a^2(\cosh^2 \eta - \cos^2 \psi).$$

$$f_1 = f_2 = f_3 = 1.$$

**Important equations**

$$(ds)^2 = a^2(\cosh^2 \eta - \cos^2 \psi) [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\text{grad } \varphi = \frac{1}{a[\cosh^2 \eta - \cos^2 \psi]^{\frac{1}{2}}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{1}{a[\cosh^2 \eta - \cos^2 \psi]^{\frac{1}{2}}} \left\{ \frac{\partial}{\partial \eta} [(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}} E_\eta] \right. \\ &\quad \left. + \frac{\partial}{\partial \psi} [(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}} E_\psi] \right\} + \frac{\partial E_z}{\partial z}. \end{aligned}$$

$$\begin{aligned} \text{curl } \mathbf{E} &= \frac{1}{(\cosh^2 \eta - \cos^2 \psi)} \\ &\times \begin{vmatrix} \mathbf{a}_\eta (\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}} & \mathbf{a}_\psi (\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}} & \mathbf{a}_z/a \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \psi} & \frac{\partial}{\partial z} \\ E_\eta (\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}} & E_\psi (\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}} & E_z/a \end{vmatrix}. \end{aligned}$$

$$\nabla^2 \varphi = \frac{1}{a^2 [\cosh^2 \eta - \cos^2 \psi]} \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = H(\eta)$ ,  $U^2 = \Psi(\psi)$ ,  $U^3 = Z(z)$ .

**General case**

$$\begin{cases} \frac{d^2 H}{d\eta^2} - (\alpha_2 + \alpha_3 a^2 \cosh^2 \eta) H = 0, \\ \frac{d^2 \Psi}{d\psi^2} + (\alpha_2 + \alpha_3 a^2 \cos^2 \psi) \Psi = 0, \\ \frac{d^2 Z}{dz^2} + \alpha_3 Z = 0. \end{cases}$$

If  $q = \alpha_3 a^2 / 4$  and  $\lambda = \alpha_2 + \alpha_3 a^2 / 2$ ,

$$\frac{d^2 H}{d\eta^2} - (\lambda + 2q \cosh 2\eta) H = 0, \quad \{113\} \quad H = A \text{ce}_m(i\eta, -q) + B \text{fe}_m(i\eta, -q)$$

or  $H = A \text{se}_m(i\eta, -q) + B \text{ge}_m(i\eta, -q)$ .

$$\frac{d^2 \Psi}{d\psi^2} + (\lambda + 2q \cos 2\psi) \Psi = 0, \quad \{113\} \quad \Psi = A \text{ce}_m(\psi, -q) + B \text{fe}_m(\psi, -q)$$

or  $\Psi = A \text{se}_m(\psi, -q) + B \text{ge}_m(\psi, -q)$ .

$$\frac{d^2 Z}{dz^2} + \frac{4q}{a^2} Z = 0, \quad \{04\} \quad Z = A \sin(2q^{\frac{1}{2}} z/a) + B \cos(2q^{\frac{1}{2}} z/a),$$

where  $m = 0, 1, 2, 3, \dots$

If  $q = -\alpha_3 a^2/4$  and  $\lambda = \alpha_2 + \alpha_3 a^2/2$ ,

$$\frac{d^2H}{d\eta^2} - (\lambda - 2q \cosh 2\eta) H = 0, \quad \{113\} \quad H = A \operatorname{ce}_m(i\eta, q) + B \operatorname{fe}_m(i\eta, q)$$

or  $H = A \operatorname{se}_m(i\eta, q) + B \operatorname{ge}_m(i\eta, q).$

$$\frac{d^2\Psi}{d\psi^2} + (\lambda - 2q \cos 2\psi) \Psi = 0, \quad \{113\} \quad \Psi = A \operatorname{ce}_m(\psi, q) + B \operatorname{fe}_m(\psi, q)$$

or  $\Psi = A \operatorname{se}_m(\psi, q) + B \operatorname{ge}_m(\psi, q).$

$$\frac{d^2Z}{dz^2} - \frac{4q}{a^2} Z = 0, \quad \{04\} \quad Z = A e^{2q^{\frac{1}{2}}z/a} + B e^{-2q^{\frac{1}{2}}z/a}.$$

For  $\varphi$  independent of  $z$ ,

$$\begin{cases} \frac{d^2H}{d\eta^2} - \alpha_2 H = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_2 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$\frac{d^2H}{d\eta^2} - p^2 H = 0, \quad \{04\} \quad H = A e^{p\eta} + B e^{-p\eta}.$$

$$\frac{d^2\Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

For  $\varphi$  independent of  $\eta$  and  $z$ ,

$$\frac{d^2\varphi}{d\psi^2} = 0, \quad \{04\} \quad \varphi = A + B\psi.$$

For  $\varphi$  independent of  $\psi$  and  $z$ ,

$$\frac{d^2\varphi}{d\eta^2} = 0, \quad \{04\} \quad \varphi = A + B\eta.$$

SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = H(\eta)$ ,  $U^2 = \Psi(\psi)$ ,  $U^3 = Z(z)$ .

General case

$$\begin{cases} \frac{d^2H}{d\eta^2} - (\alpha_2 + \alpha_3 a^2 \cosh^2 \eta) H = 0, \\ \frac{d^2\Psi}{d\psi^2} + (\alpha_2 + \alpha_3 a^2 \cos^2 \psi) \Psi = 0, \\ \frac{d^2Z}{dz^2} + (\kappa^2 + \alpha_3) Z = 0. \end{cases}$$

If  $q = \alpha_3 a^2/4$  and  $\lambda = \alpha_2 + \alpha_3 a^2/2$ ,

$$\frac{d^2H}{d\eta^2} - (\lambda + 2q \cosh 2\eta) H = 0, \quad \{113\} \quad H = A \text{ce}_m(i\eta, -q) + B \text{fe}_m(i\eta, -q) \\ \text{or} \quad H = A \text{se}_m(i\eta, -q) + B \text{ge}_m(i\eta, -q).$$

$$\frac{d^2\Psi}{d\psi^2} + (\lambda + 2q \cos 2\psi) \Psi = 0, \quad \{113\} \quad \Psi = A \text{ce}_m(\psi, -q) + B \text{fe}_m(\psi, -q) \\ \text{or} \quad \Psi = A \text{se}_m(\psi, -q) + B \text{ge}_m(\psi, -q).$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 + 4q/a^2) Z = 0, \quad \{04\} \quad Z = A \sin[(\kappa^2 + 4q/a^2)^{\frac{1}{2}}z] \\ + B \cos[(\kappa^2 + 4q/a^2)^{\frac{1}{2}}z].$$

If  $q = -\alpha_3 a^2/4$  and  $\lambda = \alpha_2 + \alpha_3 a^2/2$ ,

$$\frac{d^2H}{d\eta^2} - (\lambda - 2q \cosh 2\eta) H = 0, \quad \{113\} \quad H = A \text{ce}_m(i\eta, q) + B \text{fe}_m(i\eta, q) \\ \text{or} \quad H = A \text{se}_m(i\eta, q) + B \text{ge}_m(i\eta, q).$$

$$\frac{d^2\Psi}{d\psi^2} + (\lambda - 2q \cos 2\psi) \Psi = 0, \quad \{113\} \quad \Psi = A \text{ce}_m(\psi, q) + B \text{fe}_m(\psi, q) \\ \text{or} \quad \Psi = A \text{se}_m(\psi, q) + B \text{ge}_m(\psi, q).$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 - 4q/a^2) Z = 0, \quad \{04\} \quad Z = A \sin[(\kappa^2 - 4q/a^2)^{\frac{1}{2}}z] \\ + B \cos[(\kappa^2 - 4q/a^2)^{\frac{1}{2}}z].$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2H}{d\eta^2} = 0, \quad \{04\} \quad H = A + B\eta.$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B\psi.$$

$$\frac{d^2Z}{dz^2} + \kappa^2 Z = 0, \quad \{04\} \quad Z = A \sin \kappa z + B \cos \kappa z.$$

For  $\varphi$  independent of  $z$ ,

$$\begin{cases} \frac{d^2H}{d\eta^2} - (\alpha_2 - \kappa^2 a^2 \cosh^2 \eta) H = 0, \\ \frac{d^2\Psi}{d\psi^2} + (\alpha_2 - \kappa^2 a^2 \cos^2 \psi) \Psi = 0. \end{cases}$$

If  $q = \kappa^2 a^2/4$  and  $\lambda = \alpha_2 - \kappa^2 a^2/2$ ,

$$\frac{d^2H}{d\eta^2} - (\lambda - 2q \cosh 2\eta) H = 0, \quad \{113\} \quad H = A \text{ce}_m(i\eta, q) + B \text{fe}_m(i\eta, q) \\ \text{or} \quad H = A \text{se}_m(i\eta, q) + B \text{ge}_m(i\eta, q).$$

$$\frac{d^2\Psi}{d\psi^2} + (\lambda - 2q \cos 2\psi) \Psi = 0, \quad \{113\} \quad \Psi = A \text{ce}_m(\psi, q) + B \text{fe}_m(\psi, q) \\ \text{or} \quad \Psi = A \text{se}_m(\psi, q) + B \text{ge}_m(\psi, q).$$

If  $q = -\kappa^2 a^2/4$  and  $\lambda = \alpha_2 - \kappa^2 a^2/2$ ,

$$\frac{d^2H}{d\eta^2} - (\lambda + 2q \cosh 2\eta) H = 0, \quad \{113\} \quad H = A \text{ce}_m(i\eta, -q) + B \text{fe}_m(i\eta, -q) \\ \text{or} \quad H = A \text{se}_m(i\eta, -q) + B \text{ge}_m(i\eta, -q).$$

$$\frac{d^2\Psi}{d\psi^2} + (\lambda + 2q \cos 2\psi) \Psi = 0, \quad \{113\} \quad \Psi = A \text{ce}_m(\psi, -q) + B \text{fe}_m(\psi, -q) \\ \text{or} \quad \Psi = A \text{se}_m(\psi, -q) + B \text{ge}_m(\psi, -q).$$

TABLE 1.04. PARABOLIC-CYLINDER COORDINATES  $(\mu, \nu, z)$ 

$$u^1 = \mu, \quad 0 \leq \mu < +\infty,$$

$$u^2 = \nu, \quad -\infty < \nu < +\infty,$$

$$u^3 = z, \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{1}{2}(\mu^2 - \nu^2), \\ y = \mu\nu, \\ z = z. \end{cases}$$

The coordinate surfaces are confocal parabolic cylinders and planes:

$$\begin{cases} y^2 = \mu^2(\mu^2 - 2x) \\ \text{(parabolic cylinders, } \mu = \text{const}), \\ y^2 = \nu^2(\nu^2 + 2x) \\ \text{(parabolic cylinders, } \nu = \text{const}), \\ z = \text{const} \\ \text{(planes).} \end{cases}$$




Fig. 1.04. Parabolic-cylinder coordinates  $(\mu, \nu, z)$ . Coordinate surfaces are parabolic cylinders ( $\mu = \text{const}$ ,  $\nu = \text{const}$ ) and planes ( $z = \text{const}$ )

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} 0 & -1 & -\mu^2 \\ 0 & 1 & -\nu^2 \\ 1 & 0 & 1 \end{bmatrix}.$$

$$S = \mu^2 + \nu^2, \quad M_{11} = M_{21} = 1, \quad M_{31} = \mu^2 + \nu^2.$$

#### Metric coefficients

$$g_{11} = g_{22} = \mu^2 + \nu^2, \quad g_{33} = 1, \quad g^{\frac{1}{2}} = \mu^2 + \nu^2.$$

$$f_1 = f_2 = f_3 = 1.$$

#### Important equations

$$(ds)^2 = (\mu^2 + \nu^2) [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\text{grad } \varphi = (\mu^2 + \nu^2)^{-\frac{1}{2}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\text{div } \mathbf{E} = (\mu^2 + \nu^2)^{-1} \left\{ \frac{\partial}{\partial \mu} [(\mu^2 + \nu^2)^{\frac{1}{2}} E_\mu] + \frac{\partial}{\partial \nu} [(\mu^2 + \nu^2)^{\frac{1}{2}} E_\nu] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\text{curl } \mathbf{E} = (\mu^2 + \nu^2)^{-1} \begin{vmatrix} \mathbf{a}_\mu (\mu^2 + \nu^2)^{\frac{1}{2}} & \mathbf{a}_\nu (\mu^2 + \nu^2)^{\frac{1}{2}} & \mathbf{a}_z \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial z} \\ E_\mu (\mu^2 + \nu^2)^{\frac{1}{2}} & E_\nu (\mu^2 + \nu^2)^{\frac{1}{2}} & E_z \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{1}{\mu^2 + \nu^2} \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = Z(z)$ .

**General case**

$$\begin{cases} \frac{d^2M}{d\mu^2} - (\alpha_2 + \alpha_3 \mu^2) M = 0, \\ \frac{d^2N}{d\nu^2} + (\alpha_2 - \alpha_3 \nu^2) N = 0, \\ \frac{d^2Z}{dz^2} + \alpha_3 Z = 0. \end{cases}$$

If  $\alpha_2 = q^2(\phi + \frac{1}{2})$  and  $\alpha_3 = q^4/4$ ,

$$\frac{d^2M}{d\mu^2} - [q^2(\phi + \frac{1}{2}) + q^4\mu^2/4] M = 0, \quad \{06\} \quad M = A\mathcal{W}_e(\phi, iq\mu) + B\mathcal{W}_o(\phi, iq\mu).$$

$$\frac{d^2N}{d\nu^2} + [q^2(\phi + \frac{1}{2}) - q^4\nu^2/4] N = 0, \quad \{06\} \quad N = A\mathcal{W}_e(\phi, q\nu) + B\mathcal{W}_o(\phi, q\nu).$$

$$\frac{d^2Z}{dz^2} + \frac{q^4}{4} Z = 0, \quad \{04\} \quad Z = A \sin(q^2z/2) + B \cos(q^2z/2).$$

If  $\alpha_2 = -q^2(\phi + \frac{1}{2})$  and  $\alpha_3 = q^4/4$ ,

$$\frac{d^2M}{d\mu^2} + [q^2(\phi + \frac{1}{2}) - q^4\mu^2/4] M = 0, \quad \{06\} \quad M = A\mathcal{W}_e(\phi, q\mu) + B\mathcal{W}_o(\phi, q\mu).$$

$$\frac{d^2N}{d\nu^2} - [q^2(\phi + \frac{1}{2}) + q^4\nu^2/4] N = 0, \quad \{06\} \quad N = A\mathcal{W}_e(\phi, iq\nu) + B\mathcal{W}_o(\phi, iq\nu).$$

$$\frac{d^2Z}{dz^2} + \frac{q^4}{4} Z = 0, \quad \{04\} \quad Z = A \sin(q^2z/2) + B \cos(q^2z/2).$$

If  $\alpha_2 = 0$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2M}{d\mu^2} + q^2\mu^2 M = 0, \quad \{06\} \quad M = \mu^{\frac{1}{2}} [A\mathcal{J}_{\frac{1}{4}}(q\mu^2/2) + B\mathcal{J}_{-\frac{1}{4}}(q\mu^2/2)].$$

$$\frac{d^2N}{d\nu^2} + q^2\nu^2 N = 0, \quad \{06\} \quad N = \nu^{\frac{1}{2}} [A\mathcal{J}_{\frac{1}{4}}(q\nu^2/2) + B\mathcal{J}_{-\frac{1}{4}}(q\nu^2/2)].$$

$$\frac{d^2Z}{dz^2} - q^2 Z = 0, \quad \{04\} \quad Z = A e^{qz} + B e^{-qz}.$$

For  $\varphi$  independent of  $z$ ,

$$\begin{cases} \frac{d^2M}{d\mu^2} - \alpha_2 M = 0, \\ \frac{d^2N}{d\nu^2} + \alpha_2 N = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$\frac{d^2M}{d\mu^2} - p^2 M = 0, \quad \{04\} \quad M = A e^{p\mu} + B e^{-p\mu}.$$

$$\frac{d^2N}{d\nu^2} + p^2 N = 0, \quad \{04\} \quad N = A \sin p\nu + B \cos p\nu.$$

If  $\alpha_2 = -p^2$ ,

$$\frac{d^2M}{d\mu^2} + p^2 M = 0, \quad \{04\} \quad M = A \sin p\mu + B \cos p\mu.$$

$$\frac{d^2N}{d\nu^2} - p^2 N = 0, \quad \{04\} \quad N = A e^{p\nu} + B e^{-p\nu}.$$

**For  $\varphi$  independent of  $\mu$  and  $z$ ,**

$$\frac{d^2\varphi}{d\nu^2} = 0, \quad \{01\} \quad \varphi = A + B\nu.$$

**For  $\varphi$  independent of  $\nu$  and  $z$ ,**

$$\frac{d^2\varphi}{d\mu^2} = 0, \quad \{01\} \quad \varphi = A + B\mu.$$

**SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{d u^i} \left( f_i \frac{d U^i}{d u^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = Z(z)$ .

**General case**

$$\begin{cases} \frac{d^2M}{d\mu^2} - (\alpha_2 + \alpha_3\mu^2) M = 0, \\ \frac{d^2N}{d\nu^2} + (\alpha_2 - \alpha_3\nu^2) N = 0, \\ \frac{d^2Z}{dz^2} + (\kappa^2 + \alpha_3) Z = 0. \end{cases}$$

If  $\alpha_2 = q^2(\phi + \frac{1}{2})$  and  $\alpha_3 = q^4/4$ ,

$$\frac{d^2M}{d\mu^2} - [q^2(\phi + \frac{1}{2}) + q^4\mu^2/4] M = 0, \quad \{06\} \quad M = A\mathcal{W}_e(\phi, iq\mu) + B\mathcal{W}_o(\phi, iq\mu).$$

$$\frac{d^2N}{d\nu^2} + [q^2(\phi + \frac{1}{2}) - q^4\nu^2/4] N = 0, \quad \{06\} \quad N = A\mathcal{W}_e(\phi, q\nu) + B\mathcal{W}_o(\phi, q\nu).$$

$$\begin{aligned} \frac{d^2Z}{dz^2} + (\kappa^2 + q^4/4) Z = 0, \quad \{04\} \quad Z &= A \sin[(\kappa^2 + q^4/4)^{\frac{1}{2}}z] \\ &\quad + B \cos[(\kappa^2 + q^4/4)^{\frac{1}{2}}z]. \end{aligned}$$

If  $\alpha_2 = -q^2(\phi + \frac{1}{2})$  and  $\alpha_3 = q^4/4$ ,

$$\frac{d^2M}{d\mu^2} + [q^2(\phi + \frac{1}{2}) - q^4\mu^2/4] M = 0, \quad \{06\} \quad M = A\mathcal{W}_e(\phi, q\mu) + B\mathcal{W}_o(\phi, q\mu).$$

$$\frac{d^2N}{d\nu^2} - [q^2(\phi + \frac{1}{2}) + q^4\nu^2/4] N = 0, \quad \{06\} \quad N = A\mathcal{W}_e(\phi, iq\nu) + B\mathcal{W}_o(\phi, iq\nu).$$

$$\begin{aligned} \frac{d^2Z}{dz^2} + (\kappa^2 + q^4/4) Z = 0, \quad \{04\} \quad Z &= A \sin[(\kappa^2 + q^4/4)^{\frac{1}{2}}z] \\ &\quad + B \cos[(\kappa^2 + q^4/4)^{\frac{1}{2}}z]. \end{aligned}$$

If  $\alpha_2 = 0$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2M}{d\mu^2} + q^2\mu^2 M = 0, \quad \{06\} \quad M = \mu^{\frac{1}{2}}[A\mathcal{J}_{\frac{1}{4}}(q\mu^2/2) + B\mathcal{J}_{-\frac{1}{4}}(q\mu^2/2)].$$

$$\frac{d^2N}{d\nu^2} + q^2\nu^2 N = 0, \quad \{06\} \quad N = \nu^{\frac{1}{2}}[A\mathcal{J}_{\frac{1}{4}}(q\nu^2/2) + B\mathcal{J}_{-\frac{1}{4}}(q\nu^2/2)].$$

$$\frac{d^2Z}{dz^2} + (\kappa^2 - q^2) Z = 0, \quad \{04\} \quad Z = A \sin[(\kappa^2 - q^2)^{\frac{1}{2}}z] + B \cos[(\kappa^2 - q^2)^{\frac{1}{2}}z].$$

**For  $\varphi$  independent of  $z$ ,**

$$\begin{cases} \frac{d^2M}{d\mu^2} - (\alpha_2 - \kappa^2\mu^2) M = 0, \\ \frac{d^2N}{d\nu^2} + (\alpha_2 + \kappa^2\nu^2) N = 0. \end{cases}$$

$$\text{If } \alpha_2 = q^2,$$

$$\frac{d^2M}{d\mu^2} - (q^2 - \kappa^2\mu^2) M = 0, \quad \{06\} \quad M = \sqrt{i\mu} [A\mathcal{J}_{\frac{1}{2}}(\kappa, q, i\mu) + B\mathcal{J}_{-\frac{1}{2}}(\kappa, q, i\mu)].$$

$$\frac{d^2N}{d\nu^2} + (q^2 + \kappa^2\nu^2) N = 0, \quad \{06\} \quad N = \nu^{\frac{1}{2}} [A\mathcal{J}_{\frac{1}{2}}(\kappa, q, \nu) + B\mathcal{J}_{-\frac{1}{2}}(\kappa, q, \nu)].$$

$$\text{If } \alpha_2 = -q^2,$$

$$\frac{d^2M}{d\mu^2} + (q^2 + \kappa^2\mu^2) M = 0, \quad \{06\} \quad M = \mu^{\frac{1}{2}} [A\mathcal{J}_{\frac{1}{2}}(\kappa, q, \mu) + B\mathcal{J}_{-\frac{1}{2}}(\kappa, q, \mu)].$$

$$\frac{d^2N}{d\nu^2} - (q^2 - \kappa^2\nu^2) N = 0, \quad \{06\} \quad N = \sqrt{i\nu} [A\mathcal{J}_{\frac{1}{2}}(\kappa, q, i\nu) + B\mathcal{J}_{-\frac{1}{2}}(\kappa, q, i\nu)].$$

$$\text{If } \alpha_2 = 0,$$

$$\frac{d^2M}{d\mu^2} + \kappa^2\mu^2 M = 0, \quad \{06\} \quad M = \mu^{\frac{1}{2}} [A\mathcal{J}_{\frac{1}{4}}(\kappa\mu^2/2) + B\mathcal{J}_{-\frac{1}{4}}(\kappa\mu^2/2)].$$

$$\frac{d^2N}{d\nu^2} + \kappa^2\nu^2 N = 0, \quad \{06\} \quad N = \nu^{\frac{1}{2}} [A\mathcal{J}_{\frac{1}{4}}(\kappa\nu^2/2) + B\mathcal{J}_{-\frac{1}{4}}(\kappa\nu^2/2)].$$

TABLE 1.05. SPHERICAL COORDINATES  $(r, \theta, \psi)$




Fig. 1.05. Spherical coordinates  $(r, \theta, \psi)$ . Coordinate surfaces are spheres ( $r = \text{const}$ ), circular cones ( $\theta = \text{const}$ ), and half planes ( $\psi = \text{const}$ )

$$u^1 = r, \quad 0 \leq r < \infty,$$

$$u^2 = \theta, \quad 0 \leq \theta \leq \pi,$$

$$u^3 = \psi, \quad 0 \leq \psi < 2\pi.$$

$$\begin{cases} x = r \sin \theta \cos \psi, \\ y = r \sin \theta \sin \psi, \\ z = r \cos \theta. \end{cases}$$

Coordinate surfaces are

$$\begin{cases} x^2 + y^2 + z^2 = r^2 \\ (\text{spheres}, r = \text{const}), \end{cases}$$

$$\begin{cases} \tan \theta = (x^2 + y^2)^{\frac{1}{2}}/z \\ (\text{circular cones}, \theta = \text{const}), \end{cases}$$

$$\begin{cases} \tan \psi = y/x \\ (\text{half planes}, \psi = \text{const}). \end{cases}$$

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} 1 & -1/r^2 & 0 \\ 0 & 1 & -1/\sin^2 \theta \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = 1, \quad M_{11} = 1,$$

$$M_{21} = 1/r^2, \quad M_{31} = 1/(r^2 \sin^2 \theta).$$

#### Metric coefficients

$$g_{11} = 1, \quad g_{22} = r^2, \quad g_{33} = r^2 \sin^2 \theta, \quad g^{\frac{1}{2}} = r^2 \sin \theta.$$

$$f_1 = r^2, \quad f_2 = \sin \theta, \quad f_3 = 1.$$

**Important equations**

$$(ds)^2 = (dr)^2 + r^2(d\theta)^2 + r^2 \sin^2 \theta (d\psi)^2.$$

$$\text{grad } \varphi = \mathbf{a}_r \frac{\partial \varphi}{\partial r} + \frac{\mathbf{a}_\theta}{r} \frac{\partial \varphi}{\partial \theta} + \frac{\mathbf{a}_\psi}{r \sin \theta} \frac{\partial \varphi}{\partial \psi}.$$

$$\text{div } \mathbf{E} = \frac{\partial E_r}{\partial r} + \frac{2}{r} E_r + \frac{1}{r} \frac{\partial E_\theta}{\partial \theta} + \frac{\cot \theta}{r} E_\theta + \frac{1}{r \sin \theta} \frac{\partial E_\psi}{\partial \psi}.$$

$$\text{curl } \mathbf{E} = \frac{1}{r^2 \sin \theta} \begin{vmatrix} \mathbf{a}_r & \mathbf{a}_\theta r & \mathbf{a}_\psi r \sin \theta \\ \frac{\partial}{\partial r} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_r & E_\theta r & E_\psi r \sin \theta \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{\partial^2 \varphi}{\partial r^2} + \frac{2}{r} \frac{\partial \varphi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \varphi}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial \varphi}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

**ALTERNATIVE SPHERICAL SYSTEM  $(\xi, \theta, \psi)$** 

An alternative coordinate system is obtained by changing the radial scale:

$$r = e^\xi,$$

which makes  $g_{11} = g_{22}$ . Then

$$\begin{cases} x = e^\xi \sin \theta \cos \psi, \\ y = e^\xi \sin \theta \sin \psi, \\ z = e^\xi \cos \theta. \end{cases}$$

The Stäckel matrix is

$$[S] = \begin{bmatrix} e^{2\xi} - 1 & 0 & 0 \\ 0 & 1 - 1/\sin^2 \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = e^{2\xi}, \quad M_{11} = M_{21} = 1, \quad M_{31} = 1/\sin^2 \theta.$$

**Metric coefficients**

$$g_{11} = g_{22} = e^{2\xi}, \quad g_{33} = e^{2\xi} \sin^2 \theta, \quad g^b = e^{3\xi} \sin \theta.$$

$$f_1 = e^\xi, \quad f_2 = \sin \theta, \quad f_3 = 1.$$

**Important equations**

$$(ds)^2 = e^{2\xi} [(d\xi)^2 + (d\theta)^2 + \sin^2 \theta (d\psi)^2].$$

$$\text{grad } \varphi = e^{-\xi} \left[ \mathbf{a}_\xi \frac{\partial \varphi}{\partial \xi} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} + \frac{\mathbf{a}_\psi}{\sin \theta} \frac{\partial \varphi}{\partial \psi} \right].$$

$$\text{div } \mathbf{E} = e^{-\xi} \left\{ \frac{\partial E_\xi}{\partial \xi} + 2E_\xi + \frac{\partial E_\theta}{\partial \theta} + E_\theta \cot \theta + \frac{1}{\sin \theta} \frac{\partial E_\psi}{\partial \psi} \right\}.$$

$$\text{curl } \mathbf{E} = \frac{e^{-2\xi}}{\sin \theta} \begin{vmatrix} \mathbf{a}_\xi & \mathbf{a}_\theta & \mathbf{a}_\psi \sin \theta \\ \frac{\partial}{\partial \xi} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_\xi e^\xi & E_\theta e^\xi & E_\psi e^\xi \sin \theta \end{vmatrix}.$$

$$\nabla^2 \varphi = e^{-2\xi} \left\{ \frac{\partial^2 \varphi}{\partial \xi^2} + \frac{\partial \varphi}{\partial \xi} + \frac{\partial^2 \varphi}{\partial \theta^2} + \cot \theta \frac{\partial \varphi}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2} \right\}.$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = R(r)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ .

**General case**

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{\alpha_2}{r^2} R = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} \frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{p(p+1)}{r^2} R = 0, & \quad \{22\} \quad R = A r^p + B r^{-(p+1)}. \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left[ p(p+1) - \frac{q^2}{\sin^2 \theta} \right] \Theta = 0, & \quad \{222\} \quad \Theta = A \mathcal{P}_p^q(\cos \theta) \\ & \quad + B \mathcal{Q}_p^q(\cos \theta). \end{aligned}$$

$$\frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{\alpha_2}{r^2} R = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \alpha_2 \Theta = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$ ,

$$\begin{aligned} \frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{p(p+1)}{r^2} R = 0, & \quad \{22\} \quad R = A r^p + B r^{-(p+1)}. \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + p(p+1) \Theta = 0, & \quad \{112\} \quad \Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta). \end{aligned}$$

For  $\varphi$  independent of  $r$ ,

$$\begin{cases} \frac{d^2\Theta}{d\theta^2} + \cos \theta \frac{d\Theta}{d\theta} - \frac{\alpha_3}{\sin^2 \theta} \Theta = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_3 = q^2$ ,

$$\begin{aligned} \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} - \frac{q^2}{\sin^2 \theta} \Theta = 0, & \quad \{220\} \quad \Theta = A \mathcal{P}_0^q(\cos \theta) + B \mathcal{Q}_0^q(\cos \theta). \\ \frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0, & \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi. \end{aligned}$$

For  $\varphi$  independent of  $\theta$  and  $\psi$ ,

$$\frac{d^2\varphi}{dr^2} + \frac{2}{r} \frac{d\varphi}{dr} = 0, \quad \{01\} \quad \varphi = A + Br^{-1}.$$

For  $\varphi$  independent of  $r$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\theta^2} + \cot\theta \frac{d\varphi}{d\theta} = 0, \quad \{01\} \quad \varphi = A + B \ln \cot(\theta/2).$$

**SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = R(r)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ .

General case

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + (\kappa^2 - \alpha_2/r^2) R = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2\theta} \right) \Theta = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \left[ \kappa^2 - \frac{p(p+1)}{r^2} \right] R = 0, \quad \{24\} \quad R = r^{-\frac{1}{2}} [A J_{p+\frac{1}{2}}(\kappa r) + B J_{-(p+\frac{1}{2})}(\kappa r)].$$

$$\frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} + \left[ p(p+1) - \frac{q^2}{\sin^2\theta} \right] \Theta = 0, \quad \{222\} \quad \Theta = A P_p^q(\cos\theta) + B Q_p^q(\cos\theta).$$

$$\frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + (\kappa^2 - \alpha_2/r^2) R = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} + \alpha_2 \Theta = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$ ,

$$\frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \left[ \kappa^2 - \frac{p(p+1)}{r^2} \right] R = 0, \quad \{24\} \quad R = r^{-\frac{1}{2}} [A J_{p+\frac{1}{2}}(\kappa r) + B J_{-(p+\frac{1}{2})}(\kappa r)].$$

$$\frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} + p(p+1) \Theta = 0, \quad \{112\} \quad \Theta = A P_p(\cos\theta) + B Q_p(\cos\theta).$$

For  $\varphi$  independent of  $\theta$  and  $\psi$ ,

$$\frac{d^2\varphi}{dr^2} + \frac{2}{r} \frac{d\varphi}{dr} + \kappa^2 \varphi = 0, \quad \{04\} \quad \varphi = \frac{1}{r} [A \sin \kappa r + B \cos \kappa r].$$


TABLE 1.06. PROLATE SPHEROIDAL COORDINATES  $(\eta, \theta, \psi)$ 

Fig. 1.06. Prolate spheroidal coordinates  $(\eta, \theta, \psi)$ . Coordinate surfaces are prolate spheroids ( $\eta = \text{const}$ ), hyperboloids of revolution ( $\theta = \text{const}$ ), and half-planes ( $\psi = \text{const}$ )

$$\begin{aligned} u^1 &= \eta, & 0 \leq \eta < \infty, \\ u^2 &= \theta, & 0 \leq \theta \leq \pi, \\ u^3 &= \psi, & 0 \leq \psi < 2\pi. \end{aligned}$$

$$\left\{ \begin{array}{l} x = a \sinh \eta \sin \theta \cos \psi, \\ y = a \sinh \eta \sin \theta \sin \psi, \\ z = a \cosh \eta \cos \theta. \end{array} \right.$$

Coordinate surfaces are

$$\left\{ \begin{array}{l} \frac{x^2}{a^2 \sinh^2 \eta} + \frac{y^2}{a^2 \sinh^2 \eta} + \frac{z^2}{a^2 \cosh^2 \eta} = 1 \\ \quad (\text{prolate spheroids, } \eta = \text{const}), \\ \frac{x^2}{a^2 \sin^2 \theta} - \frac{y^2}{a^2 \sin^2 \theta} + \frac{z^2}{a^2 \cos^2 \theta} = 1 \\ \quad (\text{hyperboloids of two sheets, } \theta = \text{const}), \\ \tan \psi = y/x \\ \quad (\text{half planes, } \psi = \text{const}). \end{array} \right.$$

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} a^2 \sinh^2 \eta & -1 & -1/\sinh^2 \eta \\ a^2 \sin^2 \theta & 1 & -1/\sin^2 \theta \\ 0 & 0 & 1 \end{bmatrix}$$

$$S = a^2 (\sinh^2 \eta + \sin^2 \theta) = a^2 (\cosh^2 \eta - \cos^2 \theta),$$

$$M_{11} = M_{21} = 1, \quad M_{31} = \frac{\sinh^2 \eta + \sin^2 \theta}{\sinh^2 \eta \sin^2 \theta}.$$

#### Metric coefficients

$$g_{11} = g_{22} = a^2 (\sinh^2 \eta + \sin^2 \theta), \quad g_{33} = a^2 \sinh^2 \eta \sin^2 \theta,$$

$$g^1 = a^2 (\sinh^2 \eta + \sin^2 \theta) \sinh \eta \sin \theta.$$

$$f_1 = \sinh \eta, \quad f_2 = \sin \theta, \quad f_3 = a.$$

#### Important equations

$$(ds)^2 = a^2 (\sinh^2 \eta + \sin^2 \theta) [(d\eta)^2 + (d\theta)^2] + a^2 \sinh^2 \eta \sin^2 \theta (d\psi)^2.$$

$$\text{grad } \varphi = \frac{1}{a(\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} \right] + \frac{\mathbf{a}_\psi}{a \sinh \eta \sin \theta} \frac{\partial \varphi}{\partial \psi}.$$

$$\operatorname{div} \mathbf{E} = \frac{1}{a(\sinh^2 \eta + \sin^2 \theta)} \left\{ \frac{1}{\sinh \eta} \frac{\partial}{\partial \eta} [(\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}} \sinh \eta E_\eta] + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} [(\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}} \sin \theta E_\theta] \right\} + \frac{1}{a \sinh \eta \sin \theta} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{1}{a(\sinh^2 \eta + \sin^2 \theta) \sinh \eta \sin \theta} \times \begin{vmatrix} \mathbf{a}_\eta (\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}} & \mathbf{a}_\theta (\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}} & \mathbf{a}_\psi \sinh \eta \sin \theta \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_\eta (\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}} & E_\theta (\sinh^2 \eta + \sin^2 \theta)^{\frac{1}{2}} & E_\psi \sinh \eta \sin \theta \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{1}{a^2(\sinh^2 \eta + \sin^2 \theta)} \left\{ \frac{\partial^2 \varphi}{\partial \eta^2} + \coth \eta \frac{\partial \varphi}{\partial \eta} + \frac{\partial^2 \varphi}{\partial \theta^2} + \cot \theta \frac{\partial \varphi}{\partial \theta} \right\} + \frac{1}{a^2 \sinh^2 \eta \sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ .

General case

$$\begin{cases} \frac{d^2 H}{d \eta^2} + \coth \eta \frac{dH}{d \eta} - \left( \alpha_2 + \frac{\alpha_3}{\sinh^2 \eta} \right) H = 0, \\ \frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2 \Psi}{d \psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} \frac{d^2 H}{d \eta^2} + \coth \eta \frac{dH}{d \eta} - \left[ p(p+1) + \frac{q^2}{\sinh^2 \eta} \right] H &= 0, \\ \{222\} \quad H &= A \mathcal{P}_p^q(\cosh \eta) + B \mathcal{Q}_p^q(\cosh \eta). \\ \frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + \left[ p(p+1) - \frac{q^2}{\sin^2 \theta} \right] \Theta &= 0, \\ \{222\} \quad \Theta &= A \mathcal{P}_p^q(\cos \theta) + B \mathcal{Q}_p^q(\cos \theta). \\ \frac{d^2 \Psi}{d \psi^2} + q^2 \Psi &= 0, \\ \{04\} \quad \Psi &= A \sin q\psi + B \cos q\psi. \end{aligned}$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2 H}{d \eta^2} + \coth \eta \frac{dH}{d \eta} - \alpha_2 H = 0, \\ \frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + \alpha_2 \Theta = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$ ,

$$\begin{aligned} \frac{d^2 H}{d \eta^2} + \coth \eta \frac{dH}{d \eta} - p(p+1) H &= 0, \quad \{112\} \quad H = A \mathcal{P}_p(\cosh \eta) + B \mathcal{Q}_p(\cosh \eta). \\ \frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + p(p+1) \Theta &= 0, \quad \{112\} \quad \Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta). \end{aligned}$$

For  $\varphi$  independent of  $\theta$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\eta^2} + \coth \eta \frac{d\varphi}{d\eta} = 0, \quad \begin{cases} 01 \end{cases} \quad \varphi = A + B \ln \coth(\eta/2) \\ = A + C \ln \tanh(\eta/2).$$

For  $\varphi$  independent of  $\eta$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\theta^2} + \cot \theta \frac{d\varphi}{d\theta} = 0, \quad \begin{cases} 01 \end{cases} \quad \varphi = A + B \ln \cot(\theta/2) \\ = A + C \ln \tan(\theta/2).$$

**SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ .

**General case**

$$\begin{cases} \frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + \left( \kappa^2 a^2 \sinh^2 \eta - \alpha_2 - \frac{\alpha_3}{\sinh^2 \eta} \right) H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left( \kappa^2 a^2 \sin^2 \theta + \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} \frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + \left[ \kappa^2 a^2 \sinh^2 \eta - p(p+1) - \frac{q^2}{\sinh^2 \eta} \right] H &= 0, \\ \{224\} \quad H &= A \mathcal{P}_p^q(\kappa a, \cosh \eta) + B \mathcal{Q}_p^q(\kappa a, \cosh \eta), \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left[ \kappa^2 a^2 \sin^2 \theta + p(p+1) - \frac{q^2}{\sin^2 \theta} \right] \Theta &= 0, \\ \{224\} \quad \Theta &= A \mathcal{P}_p^q(\kappa a, \cos \theta) + B \mathcal{Q}_p^q(\kappa a, \cos \theta). \end{aligned}$$

$$\frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + (\kappa^2 a^2 \sinh^2 \eta - \alpha_2) H = 0, \\ \frac{d^2\Theta}{d\theta^2} - \cot \theta \frac{d\Theta}{d\theta} + (\kappa^2 a^2 \sin^2 \theta + \alpha_2) \Theta = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$ ,

$$\begin{aligned} \frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + [\kappa^2 a^2 \sinh^2 \eta - p(p+1)] H &= 0, \\ \{114\} \quad H &= A \mathcal{P}_p(\kappa a, \cosh \eta) + B \mathcal{Q}_p(\kappa a, \cosh \eta). \\ \frac{d^2\Theta}{d\theta^2} - \cot \theta \frac{d\Theta}{d\theta} + [\kappa^2 a^2 \sin^2 \theta + p(p+1)] \Theta &= 0, \\ \{114\} \quad \Theta &= A \mathcal{P}_p(\kappa a, \cos \theta) + B \mathcal{Q}_p(\kappa a, \cos \theta). \end{aligned}$$

**TABLE 1.07. OBLATE SPHEROIDAL COORDINATES  $(\eta, \theta, \psi)$** 

$$u^1 = \eta, \quad 0 \leq \eta < \infty,$$

$$u^2 = \theta, \quad 0 \leq \theta \leq \pi,$$

$$u^3 = \psi, \quad 0 \leq \psi < 2\pi.$$

$$\begin{cases} x = a \cosh \eta \sin \theta \cos \psi, \\ y = a \cosh \eta \sin \theta \sin \psi, \\ z = a \sinh \eta \cos \theta. \end{cases}$$

Coordinate surfaces are

$$\left\{ \begin{array}{l} \frac{x^2}{a^2 \cosh^2 \eta} + \frac{y^2}{a^2 \cosh^2 \eta} \\ \quad + \frac{z^2}{a^2 \sinh^2 \eta} = 1 \\ \text{(oblate spheroids, } \eta = \text{const}), \end{array} \right.$$

$$\left\{ \begin{array}{l} \frac{x^2}{a^2 \sin^2 \theta} + \frac{y^2}{a^2 \sin^2 \theta} \\ \quad - \frac{z^2}{a^2 \cos^2 \theta} = 1 \end{array} \right.$$

(hyperboloids of one sheet,  
 $\theta = \text{const}$ ),

$$\tan \psi = y/x$$

(half planes,  $\psi = \text{const}$ ).




Fig. 1.07. Oblate spheroidal coordinates  $(\eta, \theta, \psi)$ . Coordinate surfaces are oblate spheroids ( $\eta = \text{const}$ ), hyperboloids of revolution ( $\theta = \text{const}$ ), and half-planes ( $\psi = \text{const}$ )

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} a^2 \cosh^2 \eta & -1 & 1/\cosh^2 \eta \\ -a^2 \sin^2 \theta & 1 & -1/\sin^2 \theta \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = a^2 (\cosh^2 \eta - \sin^2 \theta) = a^2 (\sinh^2 \eta + \cos^2 \theta),$$

$$M_{11} = M_{21} = 1, \quad M_{31} = \frac{\cosh^2 \eta - \sin^2 \theta}{\cosh^2 \eta \sin^2 \theta}.$$

### Metric coefficients

$$g_{11} = g_{22} = a^2 (\cosh^2 \eta - \sin^2 \theta), \quad g_{33} = a^2 \cosh^2 \eta \sin^2 \theta,$$

$$g^{\frac{1}{2}} = a^3 (\cosh^2 \eta - \sin^2 \theta) \cosh \eta \sin \theta.$$

$$f_1 = \cosh \eta, \quad f_2 = \sin \theta, \quad f_3 = a.$$

### Important equations

$$(ds)^2 = a^2 (\cosh^2 \eta - \sin^2 \theta) [(d\eta)^2 + (d\theta)^2] + a^2 \cosh^2 \eta \sin^2 \theta (d\psi)^2.$$

$$\text{grad } \varphi = \frac{1}{a(\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} \right] + \mathbf{a}_\psi \frac{1}{a \cosh \eta \sin \theta} \frac{\partial \varphi}{\partial \psi}.$$

$$\begin{aligned}\operatorname{div} \mathbf{E} &= \frac{1}{a(\cosh^2 \eta - \sin^2 \theta)} \left\{ \frac{1}{\cosh \eta} \frac{\partial}{\partial \eta} [(\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} \cosh \eta E_\eta] \right. \\ &\quad \left. + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} [(\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} \sin \theta E_\theta] \right\} + \frac{1}{a \cosh \eta \sin \theta} \frac{\partial E_\psi}{\partial \psi}. \\ \operatorname{curl} \mathbf{E} &= \frac{1}{a(\cosh^2 \eta - \sin^2 \theta) \cosh \eta \sin \theta} \\ &\quad \times \begin{vmatrix} \mathbf{a}_\eta (\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} & \mathbf{a}_\theta (\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} & \mathbf{a}_\psi \cosh \eta \sin \theta \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_\eta (\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} & E_\theta (\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} & E_\psi \cosh \eta \sin \theta \end{vmatrix}. \\ \nabla^2 \varphi &= \frac{1}{a^2(\cosh^2 \eta - \sin^2 \theta)} \left\{ \frac{\partial^2 \varphi}{\partial \eta^2} + \tanh \eta \frac{\partial \varphi}{\partial \eta} + \frac{\partial^2 \varphi}{\partial \theta^2} + \cot \theta \frac{\partial \varphi}{\partial \theta} \right\} \\ &\quad + \frac{1}{a^2 \cosh^2 \eta \sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2}.\end{aligned}$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ .

**General case**

$$\begin{cases} \frac{d^2 H}{d \eta^2} + \tanh \eta \frac{dH}{d \eta} + \left( -\alpha_2 + \frac{\alpha_3}{\cosh^2 \eta} \right) H = 0, \\ \frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2 \Psi}{d \psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2 H}{d \eta^2} + \tanh \eta \frac{dH}{d \eta} + \left[ -p(p+1) + \frac{q^2}{\cosh^2 \eta} \right] H = 0, \quad \{222\} \quad H = A \mathcal{P}_p^q(i \sinh \eta) + B \mathcal{Q}_p^q(i \sinh \eta).$$

$$\frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + \left[ p(p+1) - \frac{q^2}{\sin^2 \theta} \right] \Theta = 0, \quad \{222\} \quad \Theta = A \mathcal{P}_p^q(\cos \theta) + B \mathcal{Q}_p^q(\cos \theta).$$

$$\frac{d^2 \Psi}{d \psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q \psi + B \cos q \psi.$$

**For  $\varphi$  independent of  $\psi$ ,**

$$\begin{cases} \frac{d^2 H}{d \eta^2} + \tanh \eta \frac{dH}{d \eta} - \alpha_2 H = 0, \\ \frac{d^2 \Theta}{d \theta^2} + \cot \theta \frac{d\Theta}{d \theta} + \alpha_2 \Theta = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$ ,

$$\frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} - p(p+1)H = 0, \quad \{112\} \quad H = A\mathcal{P}_p(i \sinh \eta) + B\mathcal{Q}_p(i \sinh \eta).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + p(p+1)\Theta = 0, \quad \{112\} \quad \Theta = A\mathcal{P}_p(\cos \theta) + B\mathcal{Q}_p(\cos \theta).$$

For  $\varphi$  independent of  $\theta$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\eta^2} + \tanh \eta \frac{d\varphi}{d\eta} = 0, \quad \{01\} \quad \varphi = A + B \cot^{-1}(\sinh \eta)$$

or  $\varphi = C + D \tan^{-1}(\sinh \eta).$

For  $\varphi$  independent of  $\eta$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\theta^2} + \cot \theta \frac{d\varphi}{d\theta} = 0, \quad \{01\} \quad \varphi = A + B \ln \cot(\theta/2).$$

**SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ .

**General case**

$$\begin{cases} \frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} + \left( \kappa^2 a^2 \cosh^2 \eta - \alpha_2 + \frac{\alpha_3}{\cosh^2 \eta} \right) H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left( -\kappa^2 a^2 \sin^2 \theta + \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} + \left[ \kappa^2 a^2 \cosh^2 \eta - p(p+1) + \frac{q^2}{\cosh^2 \eta} \right] H = 0,$$

$$\{224\} \quad H = A\mathcal{P}_p^q(i \kappa a, i \sinh \eta) + B\mathcal{Q}_p^q(i \kappa a, i \sinh \eta).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left[ -\kappa^2 a^2 \sin^2 \theta + p(p+1) - \frac{q^2}{\sin^2 \theta} \right] \Theta = 0,$$

$$\{224\} \quad \Theta = A\mathcal{P}_p^q(i \kappa a, \cos \theta) + B\mathcal{Q}_p^q(i \kappa a, \cos \theta).$$

$$\frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} + (\kappa^2 a^2 \cosh^2 \eta - \alpha_2) H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + (-\kappa^2 a^2 \sin^2 \theta + \alpha_2) \Theta = 0. \end{cases}$$

$$\text{If } \alpha_2 = p(p+1),$$

$$\frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} + [\kappa^2 a^2 \cosh^2 \eta - p(p+1)] H = 0,$$

$$\{114\} \quad H = A \mathcal{P}_p(i \kappa a, i \sinh \eta) + B \mathcal{Q}_p(i \kappa a, i \sinh \eta).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + [-\kappa^2 a^2 \sin^2 \theta + p(p+1)] \Theta = 0,$$

$$\{114\} \quad \Theta = A \mathcal{P}_p(i \kappa a, \cos \theta) + B \mathcal{Q}_p(i \kappa a, \cos \theta).$$

TABLE 1.08. PARABOLIC COORDINATES  $(\mu, \nu, \psi)$




Fig. 1.08. Parabolic coordinates  $(\mu, \nu, \psi)$ . Coordinate surfaces are paraboloids of revolution ( $\mu = \text{const}$ ,  $\nu = \text{const}$ ) and half-planes ( $\psi = \text{const}$ )

$$\begin{aligned} u^1 &= \mu, & 0 \leq \mu < \infty, \\ u^2 &= \nu, & 0 \leq \nu < \infty, \\ u^3 &= \psi, & 0 \leq \psi < 2\pi. \end{aligned}$$

$$\begin{cases} x = \mu \nu \cos \psi, \\ y = \mu \nu \sin \psi, \\ z = \frac{1}{2} (\mu^2 - \nu^2). \end{cases}$$

The coordinate surfaces are

$$\left. \begin{aligned} x^2 + y^2 &= \mu^2 (\mu^2 - 2z) \\ &\quad (\text{paraboloids of revolution, } \mu = \text{const}), \\ x^2 + y^2 &= \nu^2 (\nu^2 + 2z) \\ &\quad (\text{paraboloids of revolution, } \nu = \text{const}), \\ \tan \psi &= y/x \\ &\quad (\text{half planes, } \psi = \text{const}). \end{aligned} \right\}$$

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} \mu^2 & -1 & -1/\mu^2 \\ \nu^2 & 1 & -1/\nu^2 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = \mu^2 + \nu^2, \quad M_{11} = M_{21} = 1, \quad M_{31} = \frac{\mu^2 + \nu^2}{\mu^2 \nu^2}.$$

#### Metric coefficients

$$g_{11} = g_{22} = \mu^2 + \nu^2, \quad g_{33} = \mu^2 \nu^2, \quad g^1 = \mu \nu (\mu^2 + \nu^2).$$

$$f_1 = \mu, \quad f_2 = \nu, \quad f_3 = 1.$$

**Important equations**

$$(ds)^2 = (\mu^2 + \nu^2) [(d\mu)^2 + (d\nu)^2] + \mu^2 \nu^2 (d\psi)^2.$$

$$\text{grad } \varphi = \frac{1}{(\mu^2 + \nu^2)^{\frac{1}{2}}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \frac{\mathbf{a}_\psi}{\mu \nu} \frac{\partial \varphi}{\partial \psi}.$$

$$\text{div } \mathbf{E} = \frac{1}{\mu^2 + \nu^2} \left\{ \frac{1}{\mu} \frac{\partial}{\partial \mu} [\mu (\mu^2 + \nu^2)^{\frac{1}{2}} E_\mu] + \frac{1}{\nu} \frac{\partial}{\partial \nu} [\nu (\mu^2 + \nu^2)^{\frac{1}{2}} E_\nu] \right\} + \frac{1}{\mu \nu} \frac{\partial E_\psi}{\partial \psi}.$$

$$\text{curl } \mathbf{E} = \frac{1}{\mu \nu (\mu^2 + \nu^2)} \times \begin{vmatrix} \mathbf{a}_\mu (\mu^2 + \nu^2)^{\frac{1}{2}} & \mathbf{a}_\nu (\mu^2 + \nu^2)^{\frac{1}{2}} & \mathbf{a}_\psi \mu \nu \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ E_\mu (\mu^2 + \nu^2)^{\frac{1}{2}} & E_\nu (\mu^2 + \nu^2)^{\frac{1}{2}} & E_\psi \mu \nu \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{1}{\mu^2 + \nu^2} \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{1}{\mu} \frac{\partial \varphi}{\partial \mu} + \frac{\partial^2 \varphi}{\partial \nu^2} + \frac{1}{\nu} \frac{\partial \varphi}{\partial \nu} \right] + \frac{1}{\mu^2 \nu^2} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

**SEPARATION OF LAPLACE'S EQUATION,**  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ .

**General case**

$$\begin{cases} \frac{d^2 M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \left( \alpha_2 + \frac{\alpha_3}{\mu^2} \right) M = 0, \\ \frac{d^2 N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + \left( \alpha_2 - \frac{\alpha_3}{\nu^2} \right) N = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

$$\text{If } \alpha_2 = q^2 \text{ and } \alpha_3 = p^2,$$

$$\frac{d^2 M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - (q^2 + p^2/\mu^2) M = 0, \quad \{24\} \quad M = A \mathcal{I}_p(i q \mu) + B \mathcal{J}_{-p}(i q \mu) \\ \text{or} \quad M = A \mathcal{I}_n(i q \mu) + B \mathcal{Y}_n(i q \mu).$$

$$\frac{d^2 N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + (q^2 - p^2/\nu^2) N = 0, \quad \{24\} \quad N = A \mathcal{I}_p(q \nu) + B \mathcal{J}_{-p}(q \nu) \\ \text{or} \quad N = A \mathcal{I}_n(q \nu) + B \mathcal{Y}_n(q \nu).$$

$$\frac{d^2 \Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p \psi + B \cos p \psi.$$

**For  $\varphi$  independent of  $\Psi$ ,**

$$\begin{cases} \frac{d^2 M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \alpha_2 M = 0, \\ \frac{d^2 N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + \alpha_2 N = 0. \end{cases}$$

$$\begin{aligned} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - q^2 M &= 0, & \{14\} \quad M &= A \mathcal{J}_0(i q \mu) + B \mathcal{Y}_0(i q \mu). \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + q^2 N &= 0, & \{14\} \quad N &= A \mathcal{J}_0(q \nu) + B \mathcal{Y}_0(q \nu). \end{aligned}$$

For  $\varphi$  independent of  $\nu$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\mu^2} + \frac{1}{\mu} \frac{d\varphi}{d\mu} = 0, \quad \{01\} \quad \varphi = A + B \ln \mu.$$

For  $\varphi$  independent of  $\mu$  and  $\psi$ ,

$$\frac{d^2\varphi}{d\nu^2} + \frac{1}{\nu} \frac{d\varphi}{d\nu} = 0, \quad \{01\} \quad \varphi = A + B \ln \nu.$$

**SEPARATION OF HELMHOLTZ EQUATION,**  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ .

General case

$$\begin{cases} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + (\kappa^2 \mu^2 - \alpha_2 - \alpha_3/\mu^2) M = 0, \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + (\kappa^2 \nu^2 + \alpha_2 - \alpha_3/\nu^2) N = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = q^2$  and  $\alpha_3 = p^2$ ,

$$\begin{aligned} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + (\kappa^2 \mu^2 - q^2 - p^2/\mu^2) M &= 0, \\ \{26\} \quad M &= A \mathcal{J}_p(\kappa, i q \mu) + B \mathcal{J}_{-p}(\kappa, i q \mu) \\ &\quad \text{or} \quad M = A \mathcal{J}_n(\kappa, i q \mu) + B \mathcal{Y}_n(\kappa, i q \mu). \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + (\kappa^2 \nu^2 + q^2 - p^2/\nu^2) N &= 0, \\ \{26\} \quad N &= A \mathcal{J}_p(\kappa, q \nu) + B \mathcal{J}_{-p}(\kappa, q \nu) \\ &\quad \text{or} \quad N = A \mathcal{J}_n(\kappa, q \nu) + B \mathcal{Y}_n(\kappa, q \nu). \end{aligned}$$

$$\frac{d^2\Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p \psi + B \cos p \psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + (\kappa^2 \mu^2 - \alpha_2) M = 0, \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + (\kappa^2 \nu^2 + \alpha_2) N = 0. \end{cases}$$

If  $\alpha_2 = q^2$ ,

$$\begin{aligned} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + (\kappa^2 \mu^2 - q^2) M &= 0, \quad \{16\} \quad M = A \mathcal{J}_0(\kappa, i q \mu) + B \mathcal{Y}_0(\kappa, i q \mu). \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + (\kappa^2 \nu^2 + q^2) N &= 0, \quad \{16\} \quad N = A \mathcal{J}_0(\kappa, q \nu) + B \mathcal{Y}_0(\kappa, q \nu). \end{aligned}$$

TABLE 1.09. CONICAL COORDINATES  $(r, \theta, \lambda)$ 

$$\begin{aligned} u^1 &= r, & 0 \leqq r < \infty, \\ u^2 &= \theta, & b^2 < \theta^2 < c^2, \\ u^3 &= \lambda, & 0 < \lambda^2 < b^2. \end{aligned}$$

$$\left\{ \begin{array}{l} (x)^2 = \left( \frac{r \theta \lambda}{b c} \right)^2, \\ (y)^2 = \frac{r^2 (\theta^2 - b^2) (b^2 - \lambda^2)}{b^2 (c^2 - b^2)}, \\ (z)^2 = \frac{r^2 (c^2 - \theta^2) (c^2 - \lambda^2)}{c^2 (c^2 - b^2)}, \end{array} \right.$$

with  $c^2 > \theta^2 > b^2 > \lambda^2 > 0$ .

Coordinate surfaces are

$$\left\{ \begin{array}{l} x^2 + y^2 + z^2 = r^2 \\ \quad (\text{spheres, } r = \text{const}), \\ \frac{x^2}{\theta^2} + \frac{y^2}{\theta^2 - b^2} - \frac{z^2}{c^2 - \theta^2} = 0 \\ \quad (\text{elliptic cones, } \theta = \text{const}), \\ \frac{x^2}{\lambda^2} - \frac{y^2}{b^2 - \lambda^2} - \frac{z^2}{c^2 - \lambda^2} = 0 \\ \quad (\text{elliptic cones, } \lambda = \text{const}). \end{array} \right.$$




Fig. 1.09. Conical coordinates  $(r, \theta, \lambda)$ . Coordinate surfaces are spheres ( $r = \text{const}$ ), and elliptic cones ( $\theta = \text{const}$ ,  $\lambda = \text{const}$ )

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} 1 & -1/r^2 & 0 \\ 0 & \frac{\theta^2}{(\theta^2 - b^2)(c^2 - \theta^2)} & \frac{-1}{(\theta^2 - b^2)(c^2 - \theta^2)} \\ 0 & \frac{-\lambda^2}{(b^2 - \lambda^2)(c^2 - \lambda^2)} & \frac{1}{(b^2 - \lambda^2)(c^2 - \lambda^2)} \end{bmatrix},$$

$$S = \frac{(\theta^2 - \lambda^2)}{(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$M_{11} = S,$$

$$M_{21} = \frac{1}{r^2(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$M_{31} = \frac{1}{r^2(\theta^2 - b^2)(c^2 - \theta^2)}.$$

#### Metric coefficients

$$g_{11} = 1,$$

$$g_{22} = \frac{r^2(\theta^2 - \lambda^2)}{(\theta^2 - b^2)(c^2 - \theta^2)},$$

$$g_{33} = \frac{r^2(\theta^2 - \lambda^2)}{(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$g^{\frac{1}{2}} = \frac{r^2(\theta^2 - \lambda^2)}{[(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)]^{\frac{1}{2}}}.$$

$$f_1 = r^2,$$

$$f_2 = [(\theta^2 - b^2)(c^2 - \theta^2)]^{\frac{1}{2}},$$

$$f_3 = [(b^2 - \lambda^2)(c^2 - \lambda^2)]^{\frac{1}{2}}.$$

**Important equations**

$$\begin{aligned}
(ds)^2 &= (dr)^2 + r^2(\theta^2 - \lambda^2) \left[ \frac{(d\theta)^2}{(\theta^2 - b^2)(c^2 - \theta^2)} + \frac{(d\lambda)^2}{(b^2 - \lambda^2)(c^2 - \lambda^2)} \right]. \\
\text{grad } \varphi &= \mathbf{a}_r \frac{\partial \varphi}{\partial r} + \frac{1}{r(\theta^2 - \lambda^2)^{\frac{1}{2}}} \\
&\quad \times \left\{ \mathbf{a}_\theta (\theta^2 - b^2)^{\frac{1}{2}} (c^2 - \theta^2)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \theta} + \mathbf{a}_\lambda (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \lambda} \right\}. \\
\text{div } \mathbf{E} &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 E_r) + \frac{1}{r(\theta^2 - \lambda^2)} \left\{ (\theta^2 - b^2)^{\frac{1}{2}} (c^2 - \theta^2)^{\frac{1}{2}} \right. \\
&\quad \left. \times \frac{\partial}{\partial \theta} [(\theta^2 - \lambda^2)^{\frac{1}{2}} E_\theta] + (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{\partial}{\partial \lambda} [(\theta^2 - \lambda^2)^{\frac{1}{2}} E_\lambda] \right\}. \\
\text{curl } \mathbf{E} &= \frac{[(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)]^{\frac{1}{2}}}{(\theta^2 - \lambda^2)} \\
&\quad \times \left| \begin{array}{ccc} \mathbf{a}_r & \mathbf{a}_\theta \frac{(\theta^2 - \lambda^2)^{\frac{1}{2}}}{(\theta^2 - b^2)^{\frac{1}{2}}(c^2 - \theta^2)^{\frac{1}{2}}} & \mathbf{a}_\lambda \frac{(\theta^2 - \lambda^2)^{\frac{1}{2}}}{(b^2 - \lambda^2)^{\frac{1}{2}}(c^2 - \lambda^2)^{\frac{1}{2}}} \\ \frac{\partial}{\partial r} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \lambda} \\ E_r & E_\theta \frac{(\theta^2 - \lambda^2)^{\frac{1}{2}}}{(\theta^2 - b^2)^{\frac{1}{2}}(c^2 - \theta^2)^{\frac{1}{2}}} & E_\lambda \frac{(\theta^2 - \lambda^2)^{\frac{1}{2}}}{(b^2 - \lambda^2)^{\frac{1}{2}}(c^2 - \lambda^2)^{\frac{1}{2}}} \end{array} \right|. \\
\nabla^2 \varphi &= \frac{\partial^2 \varphi}{\partial r^2} + \frac{2}{r} \frac{\partial \varphi}{\partial r} + \frac{1}{r^2(\theta^2 - \lambda^2)} \left\{ (\theta^2 - b^2)(c^2 - \theta^2) \frac{\partial^2 \varphi}{\partial \theta^2} \right. \\
&\quad - \theta [2\theta^2 - (b^2 + c^2)] \frac{\partial \varphi}{\partial \theta} + (b^2 - \lambda^2)(c^2 - \lambda^2) \frac{\partial^2 \varphi}{\partial \lambda^2} \\
&\quad \left. + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{\partial \varphi}{\partial \lambda} \right\}.
\end{aligned}$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = R(r)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Lambda(\lambda)$ .

**General case**

$$\begin{cases} \frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{\alpha_2}{r^2} R = 0, \\ (\theta^2 - b^2)(c^2 - \theta^2) \frac{d^2 \Theta}{d\theta^2} - \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} + [\alpha_2 \theta^2 - \alpha_3] \Theta = 0, \\ (b^2 - \lambda^2)(c^2 - \lambda^2) \frac{d^2 \Lambda}{d\lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d\Lambda}{d\lambda} - [\alpha_2 \lambda^2 - \alpha_3] \Lambda = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = (b^2 + c^2)q$ ,

$$\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{p(p+1)}{r^2} R = 0, \quad \{22\} \quad R = A r^p + B r^{-(p+1)}.$$

$$\begin{aligned} (\theta^2 - b^2)(\theta^2 - c^2) \frac{d^2 \Theta}{d\theta^2} + \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} + [(b^2 + c^2)q - p(p+1)\theta^2] \Theta &= 0, \\ \{1112\} \quad \Theta = A \mathcal{E}_p^q(\theta) + B \mathcal{F}_p^q(\theta). \end{aligned}$$

$$(\lambda^2 - b^2)(\lambda^2 - c^2) \frac{d^2 A}{d \lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d A}{d \lambda} + [(b^2 + c^2) q - p(p+1)\lambda^2] A = 0,$$

$$\{1112\} \quad A = A \mathcal{E}_p^q(\lambda) + B \mathcal{F}_p^q(\lambda).$$

$$If \alpha_2 = \alpha_3 = 0,$$

$$\frac{d^2 R}{d r^2} + \frac{2}{r} \frac{d R}{d r} = 0, \quad \{01\} \quad R = A + B/r.$$

$$\frac{d}{d \theta} \left[ (\theta^2 - b^2)^{\frac{1}{2}} (c^2 - \theta^2)^{\frac{1}{2}} \frac{d \Theta}{d \theta} \right] = 0, \quad \{01\} \quad \Theta = A + B \operatorname{sn}^{-1} \left( \frac{\theta}{b}, \frac{b}{c} \right).$$

$$\frac{d}{d \lambda} \left[ (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{d \Lambda}{d \lambda} \right] = 0, \quad \{01\} \quad \Lambda = A + B \operatorname{sn}^{-1} \left( \frac{\lambda}{b}, \frac{b}{c} \right).$$

For  $\varphi$  independent of  $r$ ,

$$\begin{cases} (\theta^2 - b^2)(\theta^2 - c^2) \frac{d^2 \Theta}{d \theta^2} + \theta [2\theta^2 - (b^2 + c^2)] \frac{d \Theta}{d \theta} + \alpha_3 \Theta = 0, \\ (\lambda^2 - b^2)(\lambda^2 - c^2) \frac{d^2 \Lambda}{d \lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d \Lambda}{d \lambda} + \alpha_3 \Lambda = 0. \end{cases}$$

$$If \alpha_3 = q(b^2 + c^2),$$

$$(\theta^2 - b^2)(\theta^2 - c^2) \frac{d^2 \Theta}{d \theta^2} + \theta [2\theta^2 - (b^2 + c^2)] \frac{d \Theta}{d \theta} + q(b^2 + c^2) \Theta = 0,$$

$$\{1111\} \quad \Theta = A \mathcal{E}_0^q(\theta) + B \mathcal{F}_0^q(\theta).$$

$$(\lambda^2 - b^2)(\lambda^2 - c^2) \frac{d^2 \Lambda}{d \lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d \Lambda}{d \lambda} + q(b^2 + c^2) \Lambda = 0,$$

$$\{1111\} \quad \Lambda = A \mathcal{E}_0^q(\lambda) + B \mathcal{F}_0^q(\lambda).$$

For  $\varphi$  independent of  $r$  and  $\lambda$ ,

$$\frac{d}{d \theta} \left[ (\theta^2 - b^2)^{\frac{1}{2}} (c^2 - \theta^2)^{\frac{1}{2}} \frac{d \varphi}{d \theta} \right] = 0, \quad \{01\} \quad \varphi = A + B \operatorname{sn}^{-1} \left( \frac{\theta}{b}, \frac{b}{c} \right).$$

For  $\varphi$  independent of  $r$  and  $\theta$ ,

$$\frac{d}{d \lambda} \left[ (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{d \varphi}{d \lambda} \right] = 0, \quad \{01\} \quad \varphi = A + B \operatorname{sn}^{-1} \left( \frac{\lambda}{b}, \frac{b}{c} \right).$$

SEPARATION OF HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{d u^i} \left( f_i \frac{d U^i}{d u^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = R(r)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Lambda(\lambda)$ .

General case

$$\begin{cases} \frac{d^2 R}{d r^2} + \frac{2}{r} \frac{d R}{d r} + (\kappa^2 - \alpha_2/r^2) R = 0, \\ (\theta^2 - b^2)(c^2 - \theta^2) \frac{d^2 \Theta}{d \theta^2} - \theta [2\theta^2 - (b^2 + c^2)] \frac{d \Theta}{d \theta} + [\alpha_2 \theta^2 - \alpha_3] \Theta = 0, \\ (b^2 - \lambda^2)(c^2 - \lambda^2) \frac{d^2 \Lambda}{d \lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d \Lambda}{d \lambda} - [\alpha_2 \lambda^2 - \alpha_3] \Lambda = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q(b^2 + c^2)$ ,

$$\frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + (\kappa^2 - p(p+1)/r^2) R = 0, \quad \{24\} \quad R = r^{-\frac{1}{2}} [A \mathcal{J}_{p+\frac{1}{2}}(\kappa r) + B \mathcal{J}_{-(p+\frac{1}{2})}(\kappa r)].$$

$$(\theta^2 - b^2)(\theta^2 - c^2) \frac{d^2\Theta}{dt^2} + \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} - [p(p+1)\theta^2 - q(b^2 + c^2)] \Theta = 0, \quad \{1112\} \quad \Theta = A \mathcal{E}_p^q(\theta) + B \mathcal{F}_p^q(\theta).$$

$$(\lambda^2 - b^2)(\lambda^2 - c^2) \frac{d^2\Lambda}{d\lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d\Lambda}{d\lambda} - [p(p+1)\lambda^2 - q(b^2 + c^2)] \Lambda = 0, \quad \{1112\} \quad \Lambda = A \mathcal{E}_p^q(\lambda) + B \mathcal{F}_p^q(\lambda).$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \kappa^2 R = 0, \quad \{04\} \quad R = \frac{1}{r} [A \sin \kappa r + B \cos \kappa r].$$

$$\frac{d}{d\theta} \left[ (\theta^2 - b^2)^{\frac{1}{2}} (c^2 - \theta^2)^{\frac{1}{2}} \frac{d\Theta}{d\theta} \right] = 0, \quad \{01\} \quad \Theta = A + B \operatorname{sn}^{-1} \left( \frac{\theta}{b}, \frac{b}{c} \right).$$

$$\frac{d}{d\lambda} \left[ (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{d\Lambda}{d\lambda} \right] = 0, \quad \{01\} \quad \Lambda = A + B \operatorname{sn}^{-1} \left( \frac{\lambda}{b}, \frac{b}{c} \right).$$

For  $\varphi$  independent of  $\theta$  and  $\lambda$ ,

$$\frac{d^2\varphi}{dr^2} + \frac{2}{r} \frac{d\varphi}{dr} + \kappa^2 \varphi = 0, \quad \{04\} \quad R = \frac{1}{r} [A \sin \kappa r + B \cos \kappa r].$$

TABLE 1.10. ELLIPSOIDAL COORDINATES  $(\eta, \theta, \lambda)$

$$u^1 = \eta, \quad c^2 < \eta^2 < \infty^2,$$

$$u^2 = \theta, \quad b^2 < \theta^2 < c^2,$$

$$u^3 = \lambda, \quad 0 \leqq \lambda^2 < b^2.$$

$$\begin{cases} (x)^2 = \left( \frac{\eta \theta \lambda}{b c} \right)^2, \\ (y^2) = \frac{(\eta^2 - b^2)(\theta^2 - b^2)(b^2 - \lambda^2)}{b^2(c^2 - b^2)}, \\ (z)^2 = \frac{(\eta^2 - c^2)(c^2 - \theta^2)(c^2 - \lambda^2)}{c^2(c^2 - b^2)}, \end{cases}$$

with  $\eta^2 > c^2 > \theta^2 > b^2 > \lambda^2 > 0$ .

Coordinate surfaces are

$$\begin{cases} \frac{x^2}{\eta^2} + \frac{y^2}{\eta^2 - b^2} + \frac{z^2}{\eta^2 - c^2} = 1 & \text{(ellipsoids, } \eta = \text{const}), \\ \frac{x^2}{\theta^2} + \frac{y^2}{\theta^2 - b^2} - \frac{z^2}{c^2 - \theta^2} = 1 & \text{(hyperboloids of one sheet, } \theta = \text{const}), \\ \frac{x^2}{\lambda^2} - \frac{y^2}{b^2 - \lambda^2} - \frac{z^2}{c^2 - \lambda^2} = 1 & \text{(hyperboloids of two sheets, } \lambda = \text{const}). \end{cases}$$

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} \frac{\eta^4}{(\eta^2 - b^2)(\eta^2 - c^2)} & \frac{1}{(\eta^2 - b^2)(\eta^2 - c^2)} & \frac{\eta^2}{(\eta^2 - b^2)(\eta^2 - c^2)} \\ \frac{-\theta^4}{(\theta^2 - b^2)(c^2 - \theta^2)} & \frac{-1}{(\theta^2 - b^2)(c^2 - \theta^2)} & \frac{-\theta^2}{(\theta^2 - b^2)(c^2 - \theta^2)} \\ \frac{\lambda^4}{(b^2 - \lambda^2)(c^2 - \lambda^2)} & \frac{1}{(b^2 - \lambda^2)(c^2 - \lambda^2)} & \frac{\lambda^2}{(b^2 - \lambda^2)(c^2 - \lambda^2)} \end{bmatrix},$$

$$S = \frac{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)}{(\eta^2 - b^2)(\eta^2 - c^2)(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$M_{11} = \frac{(\theta^2 - \lambda^2)}{(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$M_{21} = \frac{(\eta^2 - \lambda^2)}{(\eta^2 - b^2)(\eta^2 - c^2)(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$M_{31} = \frac{(\eta^2 - \theta^2)}{(\eta^2 - b^2)(\eta^2 - c^2)(\theta^2 - b^2)(c^2 - \theta^2)}.$$




Fig. 1.10. Ellipsoidal coordinates  $(\eta, \theta, \lambda)$ . Coordinate surfaces are ellipsoids ( $\eta = \text{const}$ ), and hyperboloids ( $\theta = \text{const}$ ,  $\lambda = \text{const}$ )

### Metric coefficients

$$g_{11} = \frac{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)}{(\eta^2 - b^2)(\eta^2 - c^2)},$$

$$g_{22} = \frac{(\theta^2 - \lambda^2)(\eta^2 - \theta^2)}{(\theta^2 - b^2)(c^2 - \theta^2)},$$

$$g_{33} = \frac{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)}{(b^2 - \lambda^2)(c^2 - \lambda^2)},$$

$$g^{\frac{1}{2}} = \frac{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)}{[(\eta^2 - b^2)(\eta^2 - c^2)(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)]^{\frac{1}{2}}}.$$

$$f_1 = [(\eta^2 - b^2)(\eta^2 - c^2)]^{\frac{1}{2}},$$

$$f_2 = [(\theta^2 - b^2)(c^2 - \theta^2)]^{\frac{1}{2}},$$

$$f_3 = [(b^2 - \lambda^2)(c^2 - \lambda^2)]^{\frac{1}{2}}.$$

### Important equations

$$(ds)^2 = \left[ \frac{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)}{(\eta^2 - b^2)(\eta^2 - c^2)} \right] (d\eta)^2$$

$$+ \left[ \frac{(\theta^2 - \lambda^2)(\eta^2 - \theta^2)}{(\theta^2 - b^2)(c^2 - \theta^2)} \right] (d\theta)^2 + \left[ \frac{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)}{(b^2 - \lambda^2)(c^2 - \lambda^2)} \right] (d\lambda)^2.$$

$$\text{grad } \varphi = \mathbf{a}_\eta \left[ \frac{(\eta^2 - b^2)(\eta^2 - c^2)}{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)} \right]^{\frac{1}{2}} \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \left[ \frac{(\theta^2 - b^2)(c^2 - \theta^2)}{(\theta^2 - \lambda^2)(\eta^2 - \theta^2)} \right]^{\frac{1}{2}} \frac{\partial \varphi}{\partial \theta} + \mathbf{a}_\lambda \left[ \frac{(b^2 - \lambda^2)(c^2 - \lambda^2)}{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)} \right]^{\frac{1}{2}} \frac{\partial \varphi}{\partial \lambda}.$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{(\eta^2 - b^2)^{\frac{1}{2}}(\eta^2 - c^2)^{\frac{1}{2}}}{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)} \frac{\partial}{\partial \eta} [(\eta^2 - \theta^2)^{\frac{1}{2}}(\eta^2 - \lambda^2)^{\frac{1}{2}} E_\eta] \\ &+ \frac{(\theta^2 - b^2)^{\frac{1}{2}}(c^2 - \theta^2)^{\frac{1}{2}}}{(\eta^2 - \theta^2)(\theta^2 - \lambda^2)} \frac{\partial}{\partial \theta} [(\eta^2 - \theta^2)^{\frac{1}{2}}(\theta^2 - \lambda^2)^{\frac{1}{2}} E_\theta] \\ &+ \frac{(b^2 - \lambda^2)^{\frac{1}{2}}(c^2 - \lambda^2)^{\frac{1}{2}}}{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)} \frac{\partial}{\partial \lambda} [(\eta^2 - \lambda^2)^{\frac{1}{2}}(\theta^2 - \lambda^2)^{\frac{1}{2}} E_\lambda]. \end{aligned}$$

$$\begin{aligned} \text{curl } \mathbf{E} &= \frac{[(\eta^2 - b^2)(\eta^2 - c^2)(\theta^2 - b^2)(c^2 - \theta^2)(b^2 - \lambda^2)(c^2 - \lambda^2)]^{\frac{1}{2}}}{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)} \\ &\times \left| \begin{array}{ccc} \mathbf{a}_\eta \left[ \frac{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)}{(\eta^2 - b^2)(\eta^2 - c^2)} \right]^{\frac{1}{2}} & \mathbf{a}_\theta \left[ \frac{(\theta^2 - \lambda^2)(\eta^2 - \theta^2)}{(\theta^2 - b^2)(c^2 - \theta^2)} \right]^{\frac{1}{2}} & \mathbf{a}_\lambda \left[ \frac{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)}{(b^2 - \lambda^2)(c^2 - \lambda^2)} \right]^{\frac{1}{2}} \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \lambda} \\ E_\eta \left[ \frac{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)}{(\eta^2 - b^2)(\eta^2 - c^2)} \right]^{\frac{1}{2}} & E_\theta \left[ \frac{(\theta^2 - \lambda^2)(\eta^2 - \theta^2)}{(\theta^2 - b^2)(c^2 - \theta^2)} \right]^{\frac{1}{2}} & E_\lambda \left[ \frac{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)}{(b^2 - \lambda^2)(c^2 - \lambda^2)} \right]^{\frac{1}{2}} \end{array} \right|. \end{aligned}$$

$$\begin{aligned} V^2 \varphi &= \frac{(\eta^2 - b^2)^{\frac{1}{2}}(\eta^2 - c^2)^{\frac{1}{2}}}{(\eta^2 - \theta^2)(\eta^2 - \lambda^2)} \frac{\partial}{\partial \eta} \left[ (\eta^2 - b^2)^{\frac{1}{2}}(\eta^2 - c^2)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \eta} \right] \\ &+ \frac{(\theta^2 - b^2)^{\frac{1}{2}}(c^2 - \theta^2)^{\frac{1}{2}}}{(\eta^2 - \theta^2)(\theta^2 - \lambda^2)} \frac{\partial}{\partial \theta} \left[ (\theta^2 - b^2)^{\frac{1}{2}}(c^2 - \theta^2)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \theta} \right] \\ &+ \frac{(b^2 - \lambda^2)^{\frac{1}{2}}(c^2 - \lambda^2)^{\frac{1}{2}}}{(\eta^2 - \lambda^2)(\theta^2 - \lambda^2)} \frac{\partial}{\partial \lambda} \left[ (b^2 - \lambda^2)^{\frac{1}{2}}(c^2 - \lambda^2)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \lambda} \right]. \end{aligned}$$

SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_i, \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = A(\lambda)$ .

### General case

$$\left\{ \begin{array}{l} (\eta^2 - b^2)(\eta^2 - c^2) \frac{d^2 H}{d\eta^2} + \eta [2\eta^2 - (b^2 + c^2)] \frac{dH}{d\eta} + (\alpha_2 + \alpha_3 \eta^2) H = 0, \\ (\theta^2 - b^2)(c^2 - \theta^2) \frac{d^2 \Theta}{d\theta^2} - \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} - (\alpha_2 + \alpha_3 \theta^2) \Theta = 0, \\ (b^2 - \lambda^2)(c^2 - \lambda^2) \frac{d^2 A}{d\lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{dA}{d\lambda} + (\alpha_2 + \alpha_3 \lambda^2) A = 0. \end{array} \right.$$

If  $\alpha_2 = (b^2 + c^2) q$  and  $\alpha_3 = -p(p+1)$ ,

$$(\eta^2 - b^2)(\eta^2 - c^2) \frac{d^2 H}{d\eta^2} + \eta [2\eta^2 - (b^2 + c^2)] \frac{dH}{d\eta} + [(b^2 + c^2)q - p(p+1)\eta^2] H = 0,$$

$$\{1112\} \quad H = A \mathcal{E}_p^q(\eta) + B \mathcal{F}_p^q(\eta).$$

$$(\theta^2 - b^2)(\theta^2 - c^2) \frac{d^2 \Theta}{d\theta^2} + \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} + [(b^2 + c^2)q - p(p+1)\theta^2] \Theta = 0,$$

$$\{1112\} \quad \Theta = A \mathcal{E}_p^q(\theta) + B \mathcal{F}_p^q(\theta).$$

$$(\lambda^2 - b^2)(\lambda^2 - c^2) \frac{d^2 A}{d\lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{dA}{d\lambda} + [(b^2 + c^2)q - p(p+1)\lambda^2] A = 0,$$

$$\{1112\} \quad A = A \mathcal{E}_p^q(\lambda) + B \mathcal{F}_p^q(\lambda).$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d}{d\eta} \left[ (\eta^2 - b^2)^{\frac{1}{2}} (\eta^2 - c^2)^{\frac{1}{2}} \frac{dH}{d\lambda} \right] = 0, \quad \{01\} \quad H = A + B \operatorname{sn}^{-1}\left(\frac{\eta}{b}, \frac{b}{c}\right).$$

$$\frac{d}{d\theta} \left[ (\theta^2 - b^2)^{\frac{1}{2}} (\theta^2 - c^2)^{\frac{1}{2}} \frac{d\Theta}{d\theta} \right] = 0, \quad \{01\} \quad \Theta = A + B \operatorname{sn}^{-1}\left(\frac{\theta}{b}, \frac{b}{c}\right).$$

$$\frac{d}{d\lambda} \left[ (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{dA}{d\lambda} \right] = 0, \quad \{01\} \quad A = A + B \operatorname{sn}^{-1}\left(\frac{\lambda}{b}, \frac{b}{c}\right).$$

For  $\varphi$  independent of  $\theta$  und  $\lambda$ ,

$$\frac{d}{d\eta} \left[ (\eta^2 - b^2)^{\frac{1}{2}} (\eta^2 - c^2)^{\frac{1}{2}} \frac{d\varphi}{d\eta} \right] = 0, \quad \{01\} \quad \varphi = A + B \operatorname{sn}^{-1}\left(\frac{\eta}{b}, \frac{b}{c}\right)$$

For  $\varphi$  independent of  $\eta$  and  $\lambda$ ,

$$\frac{d}{d\theta} \left[ (\theta^2 - b^2)^{\frac{1}{2}} (\theta^2 - c^2)^{\frac{1}{2}} \frac{d\varphi}{d\theta} \right] = 0, \quad \{01\} \quad \varphi = A + B \operatorname{sn}^{-1}\left(\frac{\theta}{b}, \frac{b}{c}\right).$$

For  $\varphi$  independent of  $\eta$  and  $\theta$ ,

$$\frac{d}{d\lambda} \left[ (b^2 - \lambda^2)^{\frac{1}{2}} (c^2 - \lambda^2)^{\frac{1}{2}} \frac{d\varphi}{d\lambda} \right] = 0, \quad \{01\} \quad \varphi = A + B \operatorname{sn}^{-1}\left(\frac{\lambda}{b}, \frac{b}{c}\right).$$

**SEPARATION OF THE HELMHOLTZ EQUATION,  $\nabla^2 \varphi + \kappa^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = A(\lambda)$ .

**General case**

$$\begin{cases} (\eta^2 - b^2)(\eta^2 - c^2) \frac{d^2 H}{d\eta^2} + \eta [2\eta^2 - (b^2 + c^2)] \frac{dH}{d\eta} + [\kappa^2 \eta^4 + \alpha_3 \eta^2 + \alpha_2] H = 0, \\ (\theta^2 - b^2)(c^2 - \theta^2) \frac{d^2 \Theta}{d\theta^2} - \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} - [\kappa^2 \theta^4 + \alpha_3 \theta^2 + \alpha_2] \Theta = 0, \\ (b^2 - \lambda^2)(c^2 - \lambda^2) \frac{d^2 A}{d\lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{dA}{d\lambda} + [\kappa^2 \lambda^4 + \alpha_3 \lambda^2 + \alpha_2] A = 0. \end{cases}$$

$$\begin{aligned}
& \text{If } \alpha_2 = q(b^2 + c^2) \text{ and } \alpha_3 = -p(p+1), \\
& (\eta^2 - b^2)(\eta^2 - c^2) \frac{d^2 H}{d\eta^2} + \eta [2\eta^2 - (b^2 + c^2)] \frac{dH}{d\eta} \\
& + [\kappa^2 \eta^4 - p(p+1)\eta^2 + q(b^2 + c^2)] H = 0, \quad \{1113\} \quad H = A\mathcal{E}_p^q(\kappa, \eta) + B\mathcal{F}_p^q(\kappa, \eta). \\
& (\theta^2 - b^2)(\theta^2 - c^2) \frac{d^2 \Theta}{d\theta^2} + \theta [2\theta^2 - (b^2 + c^2)] \frac{d\Theta}{d\theta} \\
& + [\kappa^2 \theta^4 - p(p+1)\theta^2 + q(b^2 + c^2)] \Theta = 0, \quad \{1113\} \quad \Theta = A\mathcal{E}_p^q(\kappa, \theta) + B\mathcal{F}_p^q(\kappa, \theta). \\
& (\lambda^2 - b^2)(\lambda^2 - c^2) \frac{d^2 \Lambda}{d\lambda^2} + \lambda [2\lambda^2 - (b^2 + c^2)] \frac{d\Lambda}{d\lambda} \\
& + [\kappa^2 \lambda^4 - p(p+1)\lambda^2 + q(b^2 + c^2)] \Lambda = 0, \quad \{1113\} \quad \Lambda = A\mathcal{E}_p^q(\kappa, \lambda) + B\mathcal{F}_p^q(\kappa, \lambda).
\end{aligned}$$


TABLE 1.11. PARABOLOIDAL COORDINATES  $(\mu, v, \lambda)$ 

Fig. 1.11. Paraboloidal coordinates  $(\mu, v, \lambda)$ . Coordinate surfaces are elliptic paraboloids ( $\mu = \text{const}$ ,  $v = \text{const}$ ), and hyperbolic paraboloids ( $\lambda = \text{const}$ )

$$u^1 = \mu, \quad b < \mu < \infty,$$

$$u^2 = v, \quad 0 < v < c,$$

$$u^3 = \lambda, \quad c < \lambda < b.$$

$$\left\{
\begin{array}{l}
(x)^2 = \frac{4}{(b-c)} (\mu - b) \\
\quad \times (b - v)(b - \lambda), \\
(y)^2 = \frac{4}{(b-c)} (\mu - c) \\
\quad \times (c - v)(\lambda - c),
\end{array}
\right.$$

where  $\mu > b > \lambda > c > v > 0$ .

Coordinate surfaces are

$$\frac{x^2}{\mu - b} + \frac{y^2}{\mu - c} = -4(z - \mu)$$

(elliptic paraboloid opening downward,  $\mu = \text{const}$ ),

$$\frac{x^2}{b - v} + \frac{y^2}{c - v} = 4(z - v)$$

(elliptic paraboloid opening upward,  $v = \text{const}$ ),

$$\frac{x^2}{b - \lambda} - \frac{y^2}{\lambda - c} = 4(z - \lambda)$$

(hyperbolic paraboloid,  $\lambda = \text{const}$ ).

The Stäckel matrix may be written

$$[S] = \begin{bmatrix} \frac{\mu^2}{(\mu - b)(\mu - c)} & \frac{-1}{(\mu - b)(\mu - c)} & \frac{\mu}{(\mu - b)(\mu - c)} \\ \frac{v^2}{(b - v)(c - v)} & \frac{-1}{(b - v)(c - v)} & \frac{v}{(b - v)(c - v)} \\ \frac{-\lambda^2}{(b - \lambda)(\lambda - c)} & \frac{1}{(b - \lambda)(\lambda - c)} & \frac{-\lambda}{(b - \lambda)(\lambda - c)} \end{bmatrix}.$$

$$S = \frac{(\mu - v)(\mu - \lambda)(\lambda - v)}{(\mu - b)(\mu - c)(b - v)(c - v)(b - \lambda)(\lambda - c)},$$

$$M_{11} = \frac{(\lambda - \nu)}{(b - \nu)(c - \nu)(b - \lambda)(\lambda - c)},$$

$$M_{21} = \frac{(\mu - \lambda)}{(\mu - b)(\mu - c)(b - \lambda)(\lambda - c)},$$

$$M_{31} = \frac{(\mu - \nu)}{(\mu - b)(\mu - c)(b - \nu)(c - \nu)}.$$

**Metric coefficients**

$$g_{11} = \frac{(\mu - \nu)(\mu - \lambda)}{(\mu - b)(\mu - c)},$$

$$g_{22} = \frac{(\mu - \nu)(\lambda - \nu)}{(b - \nu)(c - \nu)},$$

$$g_{33} = \frac{(\lambda - \nu)(\mu - \lambda)}{(b - \lambda)(\lambda - c)},$$

$$g^1 = \frac{(\mu - \nu)(\mu - \lambda)(\lambda - \nu)}{[(\mu - b)(\mu - c)(b - \nu)(c - \nu)(b - \lambda)(\lambda - c)]^{\frac{1}{2}}}.$$

$$f_1 = (\mu - b)^{\frac{1}{2}}(\mu - c)^{\frac{1}{2}},$$

$$f_2 = (b - \nu)^{\frac{1}{2}}(c - \nu)^{\frac{1}{2}},$$

$$f_3 = (b - \lambda)^{\frac{1}{2}}(\lambda - c)^{\frac{1}{2}}.$$

**Important equations**

$$(ds)^2 = \left[ \frac{(\mu - \nu)(\mu - \lambda)}{(\mu - b)(\mu - c)} \right] (d\mu)^2 + \left[ \frac{(\mu - \nu)(\lambda - \nu)}{(b - \nu)(c - \nu)} \right] (d\nu)^2 + \left[ \frac{(\lambda - \nu)(\mu - \lambda)}{(b - \lambda)(\lambda - c)} \right] (d\lambda)^2.$$

$$\begin{aligned} \text{grad } \varphi &= \mathbf{a}_\mu \left[ \frac{(\mu - b)(\mu - c)}{(\mu - \nu)(\mu - \lambda)} \right]^{\frac{1}{2}} \frac{\partial \varphi}{\partial \mu} \\ &\quad + \mathbf{a}_\nu \left[ \frac{(b - \nu)(c - \nu)}{(\mu - \nu)(\lambda - \nu)} \right]^{\frac{1}{2}} \frac{\partial \varphi}{\partial \nu} + \mathbf{a}_\lambda \left[ \frac{(b - \lambda)(\lambda - c)}{(\lambda - \nu)(\mu - \lambda)} \right]^{\frac{1}{2}} \frac{\partial \varphi}{\partial \lambda}. \end{aligned}$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{(\mu - b)^{\frac{1}{2}}(\mu - c)^{\frac{1}{2}}}{(\mu - \nu)(\mu - \lambda)} \frac{\partial}{\partial \mu} [(\mu - \nu)^{\frac{1}{2}}(\mu - \lambda)^{\frac{1}{2}} E_\mu] \\ &\quad + \frac{(b - \nu)^{\frac{1}{2}}(c - \nu)^{\frac{1}{2}}}{(\mu - \nu)(\lambda - \nu)} \frac{\partial}{\partial \nu} [(\mu - \nu)^{\frac{1}{2}}(\lambda - \nu)^{\frac{1}{2}} E_\nu] \\ &\quad + \frac{(b - \lambda)^{\frac{1}{2}}(\lambda - c)^{\frac{1}{2}}}{(\mu - \lambda)(\lambda - \nu)} \frac{\partial}{\partial \lambda} [(\mu - \lambda)^{\frac{1}{2}}(\lambda - \nu)^{\frac{1}{2}} E_\lambda]. \end{aligned}$$

$$\begin{aligned} \text{curl } \mathbf{E} &= \frac{[(\mu - b)(\mu - c)(b - \nu)(c - \nu)(b - \lambda)(\lambda - c)]^{\frac{1}{2}}}{(\mu - \nu)(\mu - \lambda)(\lambda - \nu)} \\ &\quad \times \begin{vmatrix} \mathbf{a}_\mu \left[ \frac{(\mu - \nu)(\mu - \lambda)}{(\mu - b)(\mu - c)} \right]^{\frac{1}{2}} & \mathbf{a}_\nu \left[ \frac{(\mu - \nu)(\lambda - \nu)}{(b - \nu)(c - \nu)} \right]^{\frac{1}{2}} & \mathbf{a}_\lambda \left[ \frac{(\lambda - \nu)(\mu - \lambda)}{(b - \lambda)(\lambda - c)} \right]^{\frac{1}{2}} \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \lambda} \\ E_\mu \left[ \frac{(\mu - \nu)(\mu - \lambda)}{(\mu - b)(\mu - c)} \right]^{\frac{1}{2}} & E_\nu \left[ \frac{(\mu - \nu)(\lambda - \nu)}{(b - \nu)(c - \nu)} \right]^{\frac{1}{2}} & E_\lambda \left[ \frac{(\lambda - \nu)(\mu - \lambda)}{(b - \lambda)(\lambda - c)} \right]^{\frac{1}{2}} \end{vmatrix}. \end{aligned}$$

$$\begin{aligned} \nabla^2 \varphi &= \left[ \frac{(\mu - b)(\mu - c)}{(\mu - \nu)(\mu - \lambda)} \right]^{\frac{1}{2}} \frac{\partial}{\partial \mu} \left[ (\mu - b)^{\frac{1}{2}}(\mu - c)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \mu} \right] \\ &\quad + \left[ \frac{(b - \nu)(c - \nu)}{(\mu - \nu)(\lambda - \nu)} \right]^{\frac{1}{2}} \frac{\partial}{\partial \nu} \left[ (b - \nu)^{\frac{1}{2}}(c - \nu)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \nu} \right] \\ &\quad + \left[ \frac{(b - \lambda)(\lambda - c)}{(\mu - \lambda)(\lambda - \nu)} \right]^{\frac{1}{2}} \frac{\partial}{\partial \lambda} \left[ (b - \lambda)^{\frac{1}{2}}(\lambda - c)^{\frac{1}{2}} \frac{\partial \varphi}{\partial \lambda} \right]. \end{aligned}$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = A(\lambda)$ .

**General case**

$$\begin{cases} (\mu - b)(\mu - c) \frac{d^2 M}{d\mu^2} + \frac{1}{2} [2\mu - (b+c)] \frac{dM}{d\mu} - [\alpha_2 - \alpha_3 \mu] M = 0, \\ (b - \nu)(c - \nu) \frac{d^2 N}{d\nu^2} + \frac{1}{2} [2\nu - (b+c)] \frac{dN}{d\nu} - [\alpha_2 - \alpha_3 \nu] N = 0, \\ (b - \lambda)(\lambda - c) \frac{d^2 A}{d\lambda^2} - \frac{1}{2} [2\lambda - (b+c)] \frac{dA}{d\lambda} + [\alpha_2 - \alpha_3 \lambda] A = 0. \end{cases}$$

If  $\alpha_2 = (b+c)q$  and  $\alpha_3 = -p(p+1)$ ,

$$(\mu - b)(\mu - c) \frac{d^2 M}{d\mu^2} + \frac{1}{2} [2\mu - (b+c)] \frac{dM}{d\mu} - [p(p+1)\mu + q(b+c)] M = 0, \\ \{113\} \quad M = A \mathcal{B}_p^q(\mu) + B \mathcal{C}_p^q(\mu).$$

$$(\nu - b)(\nu - c) \frac{d^2 N}{d\nu^2} + \frac{1}{2} [2\nu - (b+c)] \frac{dN}{d\nu} - [p(p+1)\nu + q(b+c)] N = 0, \\ \{113\} \quad N = A \mathcal{B}_p^q(\nu) + B \mathcal{C}_p^q(\nu).$$

$$(\lambda - b)(\lambda - c) \frac{d^2 A}{d\lambda^2} + \frac{1}{2} [2\lambda - (b+c)] \frac{dA}{d\lambda} - [p(p+1)\lambda + q(b+c)] A = 0, \\ \{113\} \quad A = A \mathcal{B}_p^q(\lambda) + B \mathcal{C}_p^q(\lambda).$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d}{d\mu} \left[ (\mu - b)^{\frac{1}{2}} (\mu - c)^{\frac{1}{2}} \frac{dM}{d\mu} \right] = 0, \quad \{01\} \quad M = A + B \sin^{-1} \left[ \frac{2\mu - (b+c)}{b-c} \right].$$

$$\frac{d}{d\nu} \left[ (b - \nu)^{\frac{1}{2}} (c - \nu)^{\frac{1}{2}} \frac{dN}{d\nu} \right] = 0, \quad \{01\} \quad N = A + B \sin^{-1} \left[ \frac{2\nu - (b+c)}{b-c} \right].$$

$$\frac{d}{d\lambda} \left[ (b - \lambda)^{\frac{1}{2}} (\lambda - c)^{\frac{1}{2}} \frac{dA}{d\lambda} \right] = 0, \quad \{01\} \quad A = A + B \sin^{-1} \left[ \frac{2\lambda - (b+c)}{b-c} \right].$$

**For  $\varphi$  independent of  $\nu$  and  $\lambda$ ,**

$$\frac{d}{d\mu} \left[ (\mu - b)^{\frac{1}{2}} (\mu - c)^{\frac{1}{2}} \frac{d\varphi}{d\mu} \right] = 0, \quad \{01\} \quad \varphi = A + B \sin^{-1} \left[ \frac{2\mu - (b+c)}{b-c} \right].$$

**For  $\varphi$  independent of  $\mu$  and  $\lambda$ ,**

$$\frac{d}{d\nu} \left[ (b - \nu)^{\frac{1}{2}} (c - \nu)^{\frac{1}{2}} \frac{d\varphi}{d\nu} \right] = 0, \quad \{01\} \quad \varphi = A + B \sin^{-1} \left[ \frac{2\nu - (b+c)}{b-c} \right].$$

**For  $\varphi$  independent of  $\mu$  and  $\nu$ ,**

$$\frac{d}{d\lambda} \left[ (b - \lambda)^{\frac{1}{2}} (\lambda - c)^{\frac{1}{2}} \frac{d\varphi}{d\lambda} \right] = 0, \quad \{01\} \quad \varphi = A + B \sin^{-1} \left[ \frac{2\lambda - (b+c)}{b-c} \right].$$

SEPARATION OF THE HELMHOLTZ EQUATION,  $V^2 \varphi + \kappa^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \kappa^2$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = A(\lambda)$ .




Fig. 1.12. The distorted electric field produced by the introduction of a metal spheroid ( $\eta_0 = 0.444$ ) into a uniform field

### General case

$$\begin{cases} (\mu - b)(\mu - c) \frac{d^2M}{d\mu^2} + \frac{1}{2} [2\mu - (b+c)] \frac{dM}{d\mu} + [\kappa^2 \mu^2 + \alpha_3 \mu - \alpha_2] M = 0, \\ (b - \nu)(c - \nu) \frac{d^2N}{d\nu^2} + \frac{1}{2} [2\nu - (b+c)] \frac{dN}{d\nu} + [\kappa^2 \nu^2 + \alpha_3 \nu - \alpha_2] N = 0, \\ (b - \lambda)(\lambda - c) \frac{d^2A}{d\lambda^2} - \frac{1}{2} [2\lambda - (b+c)] \frac{dA}{d\lambda} - [\kappa^2 \lambda^2 + \alpha_3 \lambda - \alpha_2] A = 0. \end{cases}$$




Fig. 1.13. A dielectric spheroid introduced into a uniform electric field ( $\epsilon_2/\epsilon_1 = 10$ ,  $\eta_0 = 0.444$ )

If  $\alpha_2 = (b + c)q$  and  $\alpha_3 = -p(p + 1)$ ,

$$(\mu - b)(\mu - c) \frac{d^2M}{d\mu^2} + \frac{1}{2} [2\mu - (b + c)] \frac{dM}{d\mu} + [\kappa^2 \mu^2 - p(p + 1)\mu - q(b + c)] M = 0, \\ \{114\} \quad M = A \mathcal{B}_p^q(\kappa, \mu) + B \mathcal{C}_p^q(\kappa, \mu).$$

$$(\nu - b)(\nu - c) \frac{d^2N}{d\nu^2} + \frac{1}{2} [2\nu - (b + c)] \frac{dN}{d\nu} + [\kappa^2 \nu^2 - p(p + 1)\nu - q(b + c)] N = 0, \\ \{114\} \quad N = A \mathcal{B}_p^q(\kappa, \nu) + B \mathcal{C}_p^q(\kappa, \nu).$$

$$(\lambda - b)(\lambda - c) \frac{d^2A}{d\lambda^2} + \frac{1}{2} [2\lambda - (b + c)] \frac{dA}{d\lambda} + [\kappa^2 \lambda^2 - p(p + 1)\lambda - q(b + c)] A = 0, \\ \{114\} \quad A = A \mathcal{B}_p^q(\kappa, \lambda) + B \mathcal{C}_p^q(\kappa, \lambda).$$

## Section II

### TRANSFORMATIONS IN THE COMPLEX PLANE

The most promising way of extending the engineering applications of field theory is to develop new coordinate systems. Section I listed the eleven systems whose coordinate surfaces are of the first or second degree. KLEIN [12] and BÔCHER [4] extended this list to include a class of fourth-degree surfaces known as cyclides [13]. All possible systems of this class are treated by BÔCHER. They include the eleven coordinates of Section I, as well as more complicated coordinates; but all are either simply separable or  $R$ -separable (Section IV).

Another procedure for obtaining new coordinate systems is *inversion* [14]. The coordinate surfaces are reflected in a sphere, the new surfaces forming an orthogonal coordinate system which is generally more complicated than the original. The mathematics are simple but will not be considered here.

A third procedure is to form new coordinate systems [15] by complex-plane transformations. A rectangular map in the  $w$ -plane is transformed into a curvilinear but orthogonal map in the  $z$ -plane. The map is then translated to form a new *cylindrical coordinate system* in 3-space, or it is twirled about an axis of symmetry to form a *rotational coordinate system*. Infinitely many systems can be obtained in this way. There is no guarantee, however, that the Helmholtz equation or the Laplace equation will separate in the new coordinates; and this question must be investigated by the methods of Sections I and IV.

The complex-plane transformation is a very fruitful way of extending the totality of coordinate systems. Of course it does not produce the asymmetric systems such as Nos. 9, 10, and 11 of Section I or the general cyclidal systems of BÔCHER [4]. But asymmetric systems are of little practical value, so nothing of importance is lost by restriction to the symmetric case.

#### 2.01 CONFORMAL TRANSFORMATIONS [16]

Take an arbitrary relation between the  $w$ -plane and the  $z$ -plane:

$$z = \mathcal{F}(w). \quad (2.01)$$

The function  $\mathcal{F}$  may be chosen at random or may be found by the Schwarz-Christoffel method [16]. The Cauchy-Riemann equations apply:

$$\frac{\partial x}{\partial u} = \frac{\partial y}{\partial v}, \quad \frac{\partial x}{\partial v} = -\frac{\partial y}{\partial u}, \quad (2.02)$$

where  $w = u + i v$ ,  $z = x + i y$ . Thus angles are preserved by the transformation, and squares in the  $w$ -plane always map into curvilinear squares in the  $z$ -plane. Separation of Eq. (2.02) into real and imaginary parts gives the two equations,

$$\left. \begin{aligned} x &= \xi_1(u, v), \\ y &= \xi_2(u, v). \end{aligned} \right\} \quad (2.03)$$

These equations can be used in plotting the two families of curves,  $u = \text{const}$  and  $v = \text{const}$ , which form the orthogonal map in the  $z$ -plane. Twenty-one transformations of this kind are listed in Tables 2.01 and 2.02, and accurate maps are included. Note that all intersections are at right angles and all subdivisions are curvilinear squares.

We now generate coordinate systems in 3-space. *Cylindrical coordinates* are formed by translating the  $z$ -plane map perpendicular to itself, thus forming families of cylinders. The resulting coordinate system  $(u^1, u^2, u^3)$  is specified by

$$\left. \begin{aligned} x &= \xi_1(u^1, u^2), \\ y &= \xi_2(u^1, u^2), \\ z &= u^3, \end{aligned} \right\} \quad (2.04)$$

where  $\xi_1$  and  $\xi_2$  are the same as in the plane case, Eq. (2.03). The  $z$ -axis is always taken parallel to the generators of the cylinders. Metric coefficients are

$$g_{11} = g_{22} = \left( \frac{\partial \xi_1}{\partial u^1} \right)^2 + \left( \frac{\partial \xi_2}{\partial u^1} \right)^2, \quad g_{33} = 1. \quad (2.05)$$

Data on cylindrical systems are listed in Section III.

If the plane map is rotated about what was originally the  $y$ -axis, the *rotational system*  $(u^1, u^2, \psi)$  is specified by

$$\left. \begin{aligned} x &= \xi_1(u^1, u^2) \cdot \cos \psi, \\ y &= \xi_1(u^1, u^2) \cdot \sin \psi, \\ z &= \xi_2(u^1, u^2). \end{aligned} \right\} \quad (2.06)$$

Metric coefficients are

$$g_{11} = g_{22} = \left( \frac{\partial \xi_1}{\partial u^1} \right)^2 + \left( \frac{\partial \xi_2}{\partial u^1} \right)^2, \quad g_{33} = [\xi_1(u^1, u^2)]^2. \quad (2.07)$$

If the plane map is rotated about what was originally the  $x$ -axis,

$$\left. \begin{aligned} x &= \xi_2(u^1, u^2) \cdot \cos \psi, \\ y &= \xi_2(u^1, u^2) \cdot \sin \psi, \\ z &= \xi_1(u^1, u^2), \end{aligned} \right\} \quad (2.08)$$

and the metric coefficients are

$$g_{11} = g_{22} = \left( \frac{\partial \xi_2}{\partial u^1} \right)^2 + \left( \frac{\partial \xi_1}{\partial u^1} \right)^2, \quad g_{33} = [\xi_2(u^1, u^2)]^2. \quad (2.09)$$

In all rotational coordinate systems, the axis of rotation is called the  $z$ -axis and the angle about this axis is called  $\psi$ . Data on a number of rotational systems are given in Section IV.

Conformal transformations have been widely used in *two-dimensional* field problems [17], particularly in electrostatics. Practical applications have stimulated the study of a great number of such transformations [18], and these transformations can be employed in the development of new coordinate systems. Note, however, that the previous applications have been valid only for two-dimensional fields and usually only where the potential is constant on the lines  $v = \text{const}$ . In the present treatment, on the other hand, we are using the transformation as a foundation for coordinate systems in 3-space; and this method allows a wide variety of applications with arbitrary potential distributions on the boundaries.

**TABLE 2.01. TRANSFORMATIONS**

in the complex plane,

$$z = \mathcal{F}(w),$$

where  $z = x + i y$ ,  $w = u + i v$ ;  $\bar{z} = x - i y$ ,  $\bar{w} = u - i v$ .

| No.                          | Equation  | Designation                 | Fig. No. |
|------------------------------|---|-----------------------------|----------|
| <b>Power Functions</b>       |   |                             |          |
| P 1                          | $\bar{z} = 1/w$   | Tangent circles             | 2.01     |
| P 2                          | $z = \frac{1}{2} w^2$   | Parabolas                   | 2.02     |
| P 3                          | $\bar{z} = \frac{1}{2} w^{-2}$  | Cardioids                   | 2.03     |
| P 4                          | $z = \sqrt{2} w^{\frac{1}{2}}$  | Hyperbolas                  | 2.04     |
| P 5                          | $\bar{z} = \sqrt{2} w^{-\frac{1}{2}}$   | 4-leaf Roses                | 2.05     |
| <b>Exponential Functions</b> |   |                             |          |
| E 1                          | $z = e^w$   | Circles                     | 2.06     |
| E 2                          | $z = a(e^w + 1)^{\frac{1}{2}}$  | Cassinian ovals             | 2.07     |
| E 3                          | $\bar{z} = a(e^w + 1)^{-\frac{1}{2}}$   | Inverse Cassinian ovals     | 2.08     |
| → E 4                        | $\bar{z} = \frac{a(e^w + 1)}{e^w - 1}$  | Bipolar circles             | 2.09     |
| E 5                          | $z = \frac{a}{\pi} (w + 1 + e^w)$   | Maxwell curves              | 2.10     |
| <b>Logarithmic Functions</b> |   |                             |          |
| L 1                          | $z = \frac{2a}{\pi} \ln w$  | Logarithmic curves          | 2.11     |
| L 2                          | $z = \frac{2a}{\pi} \ln \tan w - i a$   | $\ln \tan$                  | 2.12     |
| L 3                          | $z = \frac{2a}{\pi} \ln \cosh w$  | $\ln \cosh$                 | 2.13     |
| <b>Hyperbolic Functions</b>  |   |                             |          |
| H 1                          | $z = a \cosh w$   | Ellipses                    | 2.14     |
| H 2                          | $z = a \operatorname{sech} w$   | Inverse ellipses            | 2.15     |
| <b>Elliptic Functions</b>    |   |                             |          |
| J 1                          | $z = a \operatorname{sn} w$   | $\operatorname{sn}$         | 2.16     |
| J 2                          | $\bar{z} = a \operatorname{cn} w$   | $\operatorname{cn}$         | 2.17     |
| J 3                          | $\bar{z} = \frac{k^{\frac{1}{2}}}{i 2a} \left( \frac{1 + i k^{\frac{1}{2}} \operatorname{sn} w}{1 - i k^{\frac{1}{2}} \operatorname{sn} w} \right)$ | Inverse $\operatorname{sn}$ | 2.18     |
| J 4                          | $\bar{z} = \frac{a}{\pi} \ln \left( \frac{1}{k \operatorname{sn}^2 w} \right)$  | $\ln \operatorname{sn}$     | 2.19     |
| J 5                          | $\bar{z} = \frac{2a}{\pi} \ln \operatorname{cn} w$  | $\ln \operatorname{cn}$     | 2.20     |
| J 6                          | $\bar{z} = \frac{2K a}{\pi} Z(w + i K') + i a$  | Zeta function               | 2.21     |

TABLE 2.02. TRANSFORMATIONS

| No.               | Transformation                         | Fig. No. | Inver-sion of | $x$  | $y$  | $g_{11}$                              |
|-------------------|--|----------|---------------|--|--|---------------------------------------|
| Power Functions   |  |          |               |  |  |                                       |
| Rect-ang. coords. |  |          |               |  |  |                                       |
| P 1               | $\bar{z} = 1/w$                        | 2.01     |               | $\frac{u}{u^2+v^2}$  | $\frac{v}{u^2+v^2}$  | $\frac{1}{(u^2+v^2)^2}$               |
| P 2               | $z = \frac{1}{2}w^2$                   | 2.02     | P 3           | $\frac{1}{2}(u^2-v^2)$   | $uv$   | $u^2+v^2$                             |
| P 3               | $\bar{z} = \frac{1}{2}w^{-2}$          | 2.03     | P 2           | $\frac{1}{2}\frac{u^2-v^2}{(u^2+v^2)^2}$                         | $\frac{uv}{(u^2+v^2)^2}$   | $\frac{1}{(u^2+v^2)^3}$               |
| P 4               | $z = \sqrt[3]{w^{\frac{1}{2}}}$        | 2.04     | P 5           | where<br>$\varrho = + (u^2+v^2)^{\frac{1}{2}}$                   | $(\varrho - u)^{\frac{1}{3}}$                                    | $\frac{1}{2}(u^2+v^2)^{-\frac{1}{3}}$ |
| P 5               | $\bar{z} = \sqrt[3]{w^{-\frac{1}{2}}}$ | 2.05     | P 4           | where<br>$\varrho = + (u^2+v^2)^{\frac{1}{2}}$                   | $\frac{1}{\varrho}(\varrho - u)^{\frac{1}{3}}$                   | $\frac{1}{2}(u^2+v^2)^{-\frac{1}{3}}$ |
| Exponentials      |  |          |               |  |  |                                       |
| E 1               | $z = e^w$                              | 2.06     | E 1           | $e^w \cos v$   | $e^w \sin v$   | $e^{2w}$                              |
| E 2               | $z = a(e^w+1)^{\frac{1}{2}}$           | 2.07     | E 3           | $\frac{a}{\sqrt{2}}[\varrho_1 + (e^w \cos v + 1)]^{\frac{1}{2}}$ | $\frac{a}{\sqrt{2}}[\varrho_1 - (e^w \cos v + 1)]^{\frac{1}{2}}$ | $\frac{a^2 e^{2w}}{4 \varrho_1}$      |

Table 2.02. Transformations

|                              |   |      |     |   |  |
|------------------------------|---|------|-----|---|--|
| <b>E 3</b>                   | $\bar{z} = a(e^w + 1)^{-\frac{1}{2}}$     | 2.08 | E 2 | $\frac{a}{\sqrt{2}\varrho_1} [\varrho_1 + (e^u \cos v + 1)]^{\frac{1}{2}}$<br>where<br>$\varrho_1 = + (e^{2u} + 2e^u \cos v + 1)^{\frac{1}{2}}$ | $\frac{a}{\sqrt{2}\varrho_1} [\varrho_1 - (e^u \cos v + 1)]^{\frac{1}{2}}$<br>$\frac{a^2 e^{2u}}{4\varrho_1^3}$  |
|                              | $\bar{z} = \frac{a(e^w + 1)}{e^w - 1}$    | 2.09 | E 1 | $\frac{a \sinh u}{\cosh u - \cos v}$  | $\frac{a \sin v}{\cosh u - \cos v}$<br>$\frac{(cosh u - cos v)^2}{(cosh u - cos v)^2}$   |
| <b>E 5</b>                   | $z = \frac{a}{\pi} (w + 1 + e^w)$         | 2.10 | —   | $\frac{a}{\pi} (u + 1 + e^u \cos v)$  | $\frac{a}{\pi} (v + e^u \sin v)$<br>$\left(\frac{a}{\pi}\right)^2 (e^{2u} + 2e^u \cos v + 1)$  |
| <b>Logarithmic Functions</b> |   |      |     |   |  |
| <b>L 1</b>                   | $z = \frac{2a}{\pi} \ln w$                | 2.11 | —   | $\frac{a}{\pi} \ln (w^2 + v^2)$   | $\frac{2a}{\pi} \tan^{-1}(v/w)$<br>$\left(\frac{2a}{\pi}\right)^2 \frac{1}{w^2 + v^2}$   |
| <b>L 2</b>                   | $z = \frac{2a}{\pi} \ln \tan w - i\alpha$ | 2.12 | —   | $\frac{a}{\pi} \ln \left[ \frac{\sin^2 u + \sinh^2 v}{\cos^2 u + \sinh^2 v} \right]$  | $\frac{2a}{\pi} \tan^{-1} \left( \frac{\sinh 2v}{\sin 2u} \right)$<br>$\left(\frac{4a}{\pi}\right)^2 \frac{1}{\sin^2 2u + \sinh^2 2v}$   |
| <b>L 3</b>                   | $z = \frac{2a}{\pi} \ln \cosh w$          | 2.13 | —   | $\frac{a}{\pi} \ln (\cosh^2 u - \sin^2 v)$  | $\frac{2a}{\pi} \tan^{-1}(\tanh u \tan v)$<br>$\left(\frac{2a}{\pi}\right)^2 \left\{ \frac{\cosh^2 u \sinh^2 u}{(\cosh^2 u + \sinh^2 u)^2} \right. \right. \\ \left. \left. + \frac{[\sinh u \cosh u + \sin v \cos v]^2}{\sin^2 v} \right\}$ |
| <b>Hyperbolic Functions</b>  |   |      |     |   |  |
| <b>H 1</b>                   | $z = a \cosh w$                           | 2.14 | H 2 | $a \cosh u \cos v$  | $a \sinh u \sin v$<br>$a^2 (\cosh^2 u - \cos^2 v)$   |
| <b>H 2</b>                   | $\bar{z} = a \operatorname{sech} w$       | 2.15 | H 1 | $\frac{a \cosh u \cos v}{\cosh^2 u - \sin^2 v}$   | $\frac{a \sinh u \sin v}{\cosh^2 u - \sin^2 v}$<br>$\frac{a^2 (\cosh^2 u - \cos^2 v)}{(\cosh^2 u - \sin^2 v)^2}$   |

Table 2.02. Continuation

| No.                       | Transformation  | Fig. No. | Inver-sion of | $x$  | $y$  | $g_{11}$   |
|---------------------------|---|----------|---------------|--|--|--|
| <b>Elliptic Functions</b> |   |          |               |  |  |  |
| J 1                       | $z = a \operatorname{sn} w$   | 2.46     | J 3           | $\frac{a}{A} \operatorname{sn} u \operatorname{dn} v$<br>where<br>$A = 1 - \operatorname{dn}^2 u \operatorname{sn}^2 v$  | $\frac{a}{A} \operatorname{cn} u \operatorname{dn} u \operatorname{sn} v \operatorname{cn} v$  | $\frac{a^2 \Omega^2}{A^2}$<br>where<br>$\Omega^2 = (1 - \operatorname{sn}^2 u \operatorname{dn}^2 v) \times (\operatorname{dn}^2 v - k^2 \operatorname{sn}^2 u)$                                   |
| J 2                       | $\bar{z} = a \operatorname{cn} w$   | 2.47     | J 2           | $\frac{a}{A} \operatorname{cn} u \operatorname{cn} v$  | $\frac{a}{A} \operatorname{sn} u \operatorname{dn} u \operatorname{sn} v \operatorname{dn} v$  | $\frac{a^2 T^2}{A^2}$<br>where<br>$T^2 = (\operatorname{sn}^2 v + \operatorname{sn}^2 u \operatorname{cn}^2 v) \times (\operatorname{dn}^2 v - k^2 \operatorname{sn}^2 u)$                         |
| J 3                       | $\bar{z} = \frac{k^{\frac{1}{2}}}{i 2a} \left( \frac{1 + i k^{\frac{1}{2}} \operatorname{sn} w}{1 - i k^{\frac{1}{2}} \operatorname{sn} w} \right)$ | 2.48     | J 1           | $\frac{A}{a T} \operatorname{sn} u \operatorname{dn} v$<br>where<br>$T = \operatorname{sn}^2 u \operatorname{dn}^2 v + [(A/k)]$<br>$+ [\operatorname{cn} u \operatorname{dn} u \operatorname{sn} v \operatorname{cn} v]^2$   | $\frac{k^{\frac{1}{2}} H}{2a T}$<br>where<br>$H = (A^2/k)$<br>$- (\operatorname{sn}^2 u \operatorname{dn}^2 u + \operatorname{cn}^2 u \times \operatorname{dn}^2 u \operatorname{sn}^2 v \operatorname{cn}^2 v)$   | $\left(\frac{A \Omega}{a T}\right)^2$<br>where<br>$\Sigma = \operatorname{sn}^2 u \operatorname{dn}^2 v + \operatorname{cn}^2 u \operatorname{dn}^2 u \operatorname{sn}^2 v \operatorname{cn}^2 v$ |
| J 4                       | $\bar{z} = \frac{a}{\pi} \ln \left( \frac{1}{k \operatorname{sn}^2 w} \right)$  | 2.49     | —             | $\frac{a}{\pi} \ln \left( \frac{A^2}{k \Sigma} \right)$<br>$\times \tan^{-1} \left( \frac{\operatorname{cn} u \operatorname{dn} u \operatorname{sn} v \operatorname{cn} v}{\operatorname{sn} u \operatorname{dn} v} \right)$ | $\frac{2a}{\pi}$<br>$\Theta = \operatorname{sn}^2 u \operatorname{cn}^2 u \operatorname{dn}^2 u \operatorname{sn}^2 v \operatorname{cn}^2 v$<br>$- \operatorname{sn}^2 v \operatorname{dn}^2 v)^2 + \operatorname{sn}^2 v \operatorname{cn}^2 v$<br>$\times \operatorname{dn}^2 v (\operatorname{cn}^2 u + \operatorname{sn}^2 u \operatorname{dn}^2 u)^2$ | $\left(\frac{2a \Theta}{\pi \Sigma}\right)^2$  |

Table 2.02. Transformations

|     |  |      |     |   |  |
|-----|--|------|-----|---|--|
| J 5 | $\bar{z} = \frac{2a}{\pi} \ln \operatorname{cn} w$ | 2.20 | $-$ | $\frac{2a}{\pi} \times \tan^{-1} \left( \frac{\operatorname{sn} u \operatorname{dn} u \operatorname{sn} v \operatorname{dn} v}{\operatorname{cn} u \operatorname{cn} v} \right)$ $+ \frac{2a}{\pi} \times \tan^{-1} \left( \frac{\operatorname{sn}^2 u \operatorname{cn}^2 u \operatorname{dn}^2 u \operatorname{sn}^2 v}{\operatorname{cn}^2 v \operatorname{dn}^2 u} \right)$ $+ \frac{2a}{\pi} \times \tan^{-1} \left( \frac{\operatorname{sn}^2 u \operatorname{cn}^2 v \operatorname{dn}^2 u}{\operatorname{cn}^2 v \operatorname{dn}^2 v} \right)$ $+ \frac{2a}{\pi} \times \tan^{-1} \left( \frac{\operatorname{sn}^2 u \operatorname{cn}^2 v \operatorname{dn}^2 v}{\operatorname{cn}^2 v \operatorname{dn}^2 u} \right)$ | $\begin{aligned} & \left( \frac{2Ka}{\pi} \right)^2 \left\{ \frac{4k^4}{A'^4} \operatorname{sn}^2 u \operatorname{cn}^2 u \right. \\ & \quad \times \operatorname{dn}^2 u \operatorname{sn}^2 v' \operatorname{cn}^2 v' \operatorname{dn}^2 v' \\ & \quad + \left[ \operatorname{dn}^2 u - \frac{E}{K} + \frac{k^2 \operatorname{sn}^2 v'}{A'^2} \right] (\operatorname{dn}^2 u \\ & \quad \times \operatorname{cn}^2 v' (\operatorname{cn}^2 u - \operatorname{sn}^2 u \operatorname{dn}^2 u) \\ & \quad - k^2 \operatorname{sn}^2 u \cdot (1 - k^2 \operatorname{sn}^4 u)) \Big ^2 \} \end{aligned}$ <p style="text-align: center;">where</p> $A' = 1 - \operatorname{dn}^2 u \operatorname{sn}^2 v'$ $v' = v + K'$ $K = \int_0^{\pi/2} \frac{d\varphi}{(1 - k^2 \sin^2 \varphi)^{1/2}}$ $K' = \int_0^{\pi/2} \frac{d\varphi}{(1 - k'^2 \sin^2 \varphi)^{1/2}}$ $E = \int_0^{\pi/2} (1 - k^2 \sin^2 \varphi)^{1/2} d\varphi$ |
| J 6 | $\bar{z} = \frac{2Ka}{\pi} Z(w + iK') + ia$        | 2.21 | $-$ | $\begin{aligned} & \frac{2Ka}{\pi} \left\{ Z(u) + \frac{k^2}{A'} \right. \\ & \quad \times \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u \operatorname{sn} v' \Big\} \\ & \quad + \frac{\pi v}{2K'K} - \frac{1}{A'} \\ & \quad \times \operatorname{dn}^2 u \operatorname{sn} v' \operatorname{cn} v' \operatorname{dn} v' \} \end{aligned}$  |  |

With all elliptic functions, the modulus  $k$  is associated with the variable  $u$  and  $k'$  is associated with  $v$ . For example,  $\operatorname{sn} v \equiv \operatorname{sn}(v, k')$ ,  $\operatorname{dn} u \equiv \operatorname{dn}(u, k)$ .





Fig. 2.01. Tangent circles obtained by the transformation  $\bar{z} = 1/w$  (Transformation P4)

Fig. 2.02. Parabolas,  $z = \frac{1}{2}w^2$  (P 2)





Fig. 2.03. Cardioids,  $\bar{z} = \frac{1}{2}w^{-2}$  (P 3)

Fig. 2.04. Hyperbolas,  $z = \sqrt{2}w^{1/2}$  (P 4)





Fig. 2.05. 4-leaf roses,  $\bar{z} = \sqrt{2}w^{-\frac{1}{2}}$  (P 5)

Fig. 2.06. Circles,  $z = e^{iu}$  (E 1)




Fig. 2.07. Cassinian ovals,  $z = a(e^{iu} + 1)^{1/2}$  (E2)




Fig. 2.08. Inverse Cassinian ovals,  $\bar{z} = a(e^w + 1)^{-1/2}$  (E 3)




Fig. 2.99. Bipolar circles,  $\bar{z} = \frac{a(e^w + 1)}{e^w - 1}$  (E 4)




Fig. 2.10. Maxwell curves,  $z = \frac{a}{\pi} (w + 1 + e^w)$  (Eq. 5)




Fig. 2.11. Logarithmic curves,  $z = \frac{2a}{\pi} \ln w$  ( $L_1$ )




Fig. 2.12. In tan curves,  $z = \frac{2a}{\pi} \ln \tan w - ia$  (L.2)





Fig. 2.13. In  $\cosh$  curves,  $z = \frac{2a}{\pi} \ln \cosh w$  (L.3)

Fig. 2.44. Ellipses,  $z = a \cosh w$  (H 1)





Fig. 2.15. Inverse ellipses,  $\tilde{z} = u \operatorname{sech} w$  (H 2)

Fig. 2.16. sm curves,  $z = a \sin w$  (J1)




Fig. 2.17. cn curves,  $\tilde{z} = \alpha \operatorname{cn} w$  ( $j 2$ )

Table 2.02. Transformations




Fig. 2.18. Inverse sn,  $\bar{z} = \frac{k_3^1}{i2a} \left( \frac{1 + ik_3^1 \operatorname{sn} w}{1 - ik_3^1 \operatorname{sn} w} \right)$  (1.3)




Fig. 2.19. In sn curves,  $\bar{z} = \frac{a}{\pi} \ln \left( \frac{1}{k \sin^2 w} \right)$  (J 4)




Fig. 2.20. In cn curves,  $\bar{z} = \frac{2a}{\pi} \ln \operatorname{cn} w$  (J 5)




Fig. 2.21.  $\zeta$ -function curves,  $\bar{z} = \frac{2Ka}{\pi} Z(w + iK) + ia$  (J 6)

### Section III

## CYLINDRICAL SYSTEMS

The cylindrical coordinate systems are obtained by translating each of the maps, Figs. 2.01 to 2.21, in a direction perpendicular to the graphs, thus forming two orthogonal families of cylinders. The third family of coordinate surfaces consists of parallel planes,  $z = \text{const}$ . The coordinate axis that is parallel to the generators of the cylinders is called the  $z$ -axis in all cases.

The following coordinate systems are considered in this section:

| No.   | Name                                  | Fig. No.   |
|-------|---------------------------------------|------------|
| P 1 C | Tangent-cylinder coordinates          | 2.01       |
| P 2 C | Parabolic-cylinder coordinates        | 2.02, 1.04 |
| P 3 C | Cardioid-cylinder coordinates         | 2.03       |
| P 4 C | Hyperbolic-cylinder coordinates       | 2.04       |
| P 5 C | Rose coordinates                      | 2.05       |
| E 1 C | Circular-cylinder coordinates         | 2.06, 1.02 |
| E 2 C | Cassinian-oval coordinates            | 2.07       |
| E 3 C | Inverse Cassinian-oval coordinates    | 2.08       |
| E 4 C | Bi-cylindrical coordinates            | 2.09       |
| E 5 C | Maxwell-cylinder coordinates          | 2.10       |
| L 1 C | Logarithmic-cylinder coordinates      | 2.11       |
| L 2 C | $\ln \tan$ cylinder coordinates       | 2.12       |
| L 3 C | $\ln \cosh$ cylinder coordinates      | 2.13       |
| H 1 C | Elliptic-cylinder coordinates         | 2.14, 1.03 |
| H 2 C | Inverse elliptic-cylinder coordinates | 2.15       |
| J 1 C | sn-cylinder coordinates               | 2.16       |
| J 2 C | cn-cylinder coordinates               | 2.17       |
| J 3 C | Inverse sn-cylinder coordinates       | 2.18       |
| J 4 C | $\ln \text{sn}$ cylinder coordinates  | 2.19       |
| J 5 C | $\ln \text{cn}$ cylinder coordinates  | 2.20       |
| J 6 C | Zeta coordinates                      | 2.21       |

This list can be extended indefinitely. We have included here merely a few coordinate systems that seemed promising with regard to engineering applications. Table 3.01 lists the cylindrical coordinate systems obtained from the transformations of Section II. For each system, the metric coefficients are given and the separability data are evaluated. Table 3.02 presents the gradient, divergence, curl, and Laplacian for each of these coordinate systems. The appearance of the coordinate surfaces can be visualized by reference to Figs. 2.01 to 2.21.

### 3.01 CHARACTERISTICS

The equations of Table 3.02 are obtained from the general expressions of Section I, which are considerably simplified in this special case. For cylindrical systems with  $g_{11} = g_{22}$ ,

$$(ds)^2 = g_{11} [(du^1)^2 + (du^2)^2] + (dz)^2, \quad (3.01)$$

$$\text{grad } \varphi = \frac{1}{(g_{11})^{\frac{1}{2}}} \left[ \mathbf{a}_1 \frac{\partial \varphi}{\partial u^1} + \mathbf{a}_2 \frac{\partial \varphi}{\partial u^2} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}, \quad (3.02)$$

$$\text{div } \mathbf{E} = \frac{1}{g_{11}} \left\{ \frac{\partial}{\partial u^1} [(g_{11})^{\frac{1}{2}} E_1] + \frac{\partial}{\partial u^2} [(g_{11})^{\frac{1}{2}} E_2] \right\} + \frac{\partial E_z}{\partial z}, \quad (3.03)$$

$$\begin{aligned} \text{curl } \mathbf{E} = & \mathbf{a}_1 \left[ \frac{1}{(g_{11})^{\frac{1}{2}}} \frac{\partial E_z}{\partial u^2} - \frac{\partial E_2}{\partial z} \right] + \mathbf{a}_2 \left[ \frac{\partial E_1}{\partial z} - \frac{1}{(g_{11})^{\frac{1}{2}}} \frac{\partial E_z}{\partial u^1} \right] \\ & + \mathbf{a}_z \left[ \frac{\partial}{\partial u^1} [(g_{11})^{\frac{1}{2}} E_2] - \frac{\partial}{\partial u^2} [(g_{11})^{\frac{1}{2}} E_1] \right], \end{aligned} \quad (3.04)$$

$$\nabla^2 \varphi = \frac{1}{g_{11}} \left[ \frac{\partial^2 \varphi}{(\partial u^1)^2} + \frac{\partial^2 \varphi}{(\partial u^2)^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}. \quad (3.05)$$

No cylindrical coordinate system allows  $R$ -separation [5]. Also, except for the four cylindrical systems of Section I, no cylindrical system is known that allows separation of the Helmholtz equation [9] or that allows separation of the Laplace equation where  $\varphi = \varphi(u^1, u^2, u^3)$ . Therefore, *the only separation that can be considered in Table 3.02 is simple separation of the Laplace equation with  $\varphi = \varphi(u^1, u^2)$ .*

But if  $\varphi$  is independent of  $z$ , Eq. (3.05) gives for Laplace's equation,

$$\frac{\partial^2 \varphi}{(\partial u^1)^2} + \frac{\partial^2 \varphi}{(\partial u^2)^2} = 0, \quad (3.06)$$

which is the same as the equation obtained in rectangular coordinates. Particular solutions are

$$\varphi = e^{\pm p u^1} \begin{pmatrix} \sin \\ \cos \end{pmatrix} p u^2 \quad (3.07)$$

or

$$\varphi = e^{\pm p u^2} \begin{pmatrix} \sin \\ \cos \end{pmatrix} p u^1.$$

*These solutions apply to all the coordinate systems [18] of Table 3.02.*

A one-dimensional solution is also possible for Laplace's equation (but not for the Helmholtz equation) in cylindrical coordinates. If  $\varphi$  is independent of  $z$  and  $u^2$ , Eq. (3.05) gives for Laplace's equation,

$$\frac{d^2 \varphi}{(du^1)^2} = 0, \quad (3.08)$$

whose general solution is

$$\varphi = A + B u^1. \quad (3.09)$$

Thus there is no need of listing solutions under each coordinate system, as was done in Section I: Eqs. (3.07) and (3.09) apply to all the coordinates of Section III.

The one-dimensional solution, Eq. (3.09), is equivalent to the conventional method of field mapping by conformal transformation [17]. Here boundary conditions fix the potentials of a set of surfaces such as  $u^1 = \text{const}$ . But this approach is inadequate for the general case, where arbitrary potential distributions exist on the boundaries. In this case, the usual conformal-mapping technique fails; but its extension, by the development of new cylindrical coordinate systems, allows the treatment of two-dimensional field problems with arbitrary boundary conditions.

Table 3.01. Some cylindrical coordinate systems

TABLE 3.01. SOME CYLINDRICAL COORDINATE SYSTEMS

| No.   | Transformation   | Fig.<br>No. | Equations  | Metric Coefficients   | Simple Separability |                       | One-dimensional<br>Solution |
|-------|--|-------------|--|---|---------------------|-----------------------|-----------------------------|
|       |  |             |  |   | HELMHOLTZ           | LAPLACE               |                             |
|       |  |             |  |   | $\phi(n_1, n_2)$    | $\phi(n_1, n_2, n_3)$ |                             |
| P 1 C | $\bar{z} = 1/w$<br>Tangent cylinder                                  | 2.01        | $x = \frac{\mu}{\mu^2 + v^2}$<br>$y = \frac{v}{\mu^2 + v^2}$<br>$z = z$  | $g_{11} = g_{22} = (\mu^2 + v^2)^{-2}$<br>$g_{33} = 1$                      | ×                   | ×                     | ●<br>×                      |
| P 2 C | $z = \frac{1}{2}w^2$<br>Parabolic cylinder                           | 2.02        | $x = \frac{1}{2}(\mu^2 - v^2)$<br>$y = \mu v$<br>$z = z$   | $g_{11} = g_{22} = \mu^2 + v^2$<br>$g_{33} = 1$                             | ●<br>●              | ●<br>×                | ●<br>×                      |
| P 3 C | $\bar{z} = \frac{1}{2}w^{-2}$<br>Cardioid cylinder                   | 2.03        | $x = \frac{1}{2} \frac{\mu^2 - v^2}{(\mu^2 + v^2)^2}$<br>$y = \frac{\mu v}{(\mu^2 + v^2)^2}$<br>$z = z$                                    | $g_{11} = g_{22} = (\mu^2 + v^2)^{-3}$<br>$g_{33} = 1$                      | ×                   | ×                     | ●<br>×                      |
| P 4 C | $z = \sqrt[3]{2} w^{\frac{1}{3}}$<br>Hyperbolic-cylinder coordinates | 2.04        | $x = (\varrho + \mu)^{\frac{1}{3}}$<br>$y = (\varrho - \mu)^{\frac{1}{3}}$<br>$z = z$<br>where<br>$\varrho = +(\mu^2 + v^2)^{\frac{1}{3}}$ | $g_{11} = g_{22} = \frac{1}{2}(\mu^2 + v^2)^{-\frac{1}{3}}$<br>$g_{33} = 1$ | ×                   | ×                     | ●<br>×                      |

## Section III. Cylindrical systems

Table 3.01. Continuation

| No.   | Transformation<br>Fig.<br>No.                                   | Equations   | Metric Coefficients  | Simple Separability |           |                | One-dimensional<br>solution |
|-------|---|---|--|---------------------|-----------|----------------|-----------------------------|
|       |   |   |  | HELMHOLTZ           | LAPLACE   | HELM.<br>HOLTZ |                             |
| P 5 C | $\bar{z} = \sqrt{2} w^{-\frac{1}{2}}$<br>Rose<br>coordinates    | $x = \frac{1}{\varrho} (\varrho + \mu)^{\frac{1}{2}}$<br>$y = \frac{1}{\varrho} (\varrho - \mu)^{\frac{1}{2}}$<br>$z = z$<br>where<br>$\varrho = + (\mu^2 + v^2)^{\frac{1}{2}}$   | $g_{11} = g_{22} = \frac{1}{2(\mu^2 + v^2)^{\frac{3}{2}}}$<br>$g_{33} = 1$ | $\times$            | $\times$  | $\bullet$      | $\bullet$                   |
| E 1 C | $z = e^w$<br>Circular-cylinder<br>coordinates                   | $x = e^\eta \cos \psi$<br>$y = e^\eta \sin \psi$<br>$z = z$   | $g_{11} = g_{22} = e^{2\eta}$<br>$g_{33} = 1$                              | $\bullet$           | $\bullet$ | $\bullet$      | $\bullet$                   |
| E 2 C | $z = a(e^w + 1)^{\frac{1}{2}}$<br>Cassinian-oval<br>coordinates | $x = \frac{a}{\sqrt{2}} [\varrho_1 + (e^\eta \cos \psi + 1)]^{\frac{1}{2}}$<br>$y = \frac{a}{\sqrt{2}} [\varrho_1 - (e^\eta \cos \psi + 1)]^{\frac{1}{2}}$<br>$z = z$<br>where<br>$\varrho_1 = + (e^{2\eta} + 2e^\eta \cos \psi + 1)^{\frac{1}{2}}$ | $g_{11} = g_{22} = \frac{a^2 e^{2\eta}}{4 \varrho_1}$<br>$g_{33} = 1$      | $\times$            | $\times$  | $\bullet$      | $\bullet$                   |

Table 3.01. Some cylindrical coordinate systems

|       |   |   |  |  |
|-------|---|---|--|--|
| E 3 C | $\bar{z} = \frac{a}{(e^w + 1)^{\frac{1}{2}}}$ | $x = \frac{a}{\sqrt{2}\varrho} [\varrho + (e^\eta \cos \psi + 1)]^{\frac{1}{2}}$<br>$y = \frac{a}{\sqrt{2}\varrho} [\varrho - (e^\eta \cos \psi + 1)]^{\frac{1}{2}}$<br>$z = z$<br>where<br>$\varrho = + (e^{2\eta} + 2e^\eta \cos \psi + 1)^{\frac{1}{2}}$ | $g_{11} = g_{22} = \frac{a^2 e^{2\eta}}{4\varrho^3}$<br>$g_{33} = 1$   | $\times$<br>$\times$<br>$\times$<br>$\times$ |
| E 4 C | $\bar{z} = \frac{a(e^w + 1)}{e^w - 1}$        | $x = \frac{a \sinh \eta}{\cosh \eta - \cos \psi}$<br>$y = \frac{a \sin \psi}{\cosh \eta - \cos \psi}$<br>$z = z$  | $g_{11} = g_{22} = \frac{a^2}{(\cosh \eta - \cos \psi)^2}$<br>$g_{33} = 1$                                       | $\times$<br>$\times$<br>$\times$             |
| E 5 C | Maxwell-cylinder coordinates                  | $x = \frac{a}{\pi} (\eta + 1 + e^\eta \cos \psi)$<br>$y = \frac{a}{\pi} (\psi + e^\eta \sin \psi)$<br>$z = z$   | $g_{11} = g_{22} = \left(\frac{a}{\pi}\right)^2$<br>$\times (1 + 2e^\eta \cos \psi + e^{2\eta})$<br>$g_{33} = 1$ | $\times$<br>$\times$<br>$\times$             |
| L 1 C | $z = \frac{2a}{\pi} \ln w$                    | $x = \frac{a}{\pi} \ln (\mu^2 + \nu^2)$<br>$y = \frac{2a}{\pi} \tan^{-1}(v/\mu)$<br>$z = z$   | $g_{11} = g_{22} = \left(\frac{2a}{\pi}\right)^2 \frac{1}{\mu^2 + \nu^2}$<br>$g_{33} = 1$                        | $\times$<br>$\times$<br>$\times$             |

Table 3.01. Continuation

| No.   | Transformation                      | Fig.<br>No. | Equations   | Metric Coefficients  | Simple Separability |           |           | One-dimensional<br>Solution |
|-------|-------------------------------------|-------------|---|--|---------------------|-----------|-----------|-----------------------------|
|       |                                     |             |   |  | HELMHOLTZ           | LAPLACE   | LAPLACE   |                             |
| L 2 C | $z = \frac{2a}{\pi} \ln \tan w$     | 2.12        | $x = \frac{a}{\pi} \ln \left[ \frac{\sinh^2 \eta + \sin^2 \psi}{\sinh^2 \eta + \cos^2 \psi} \right]$<br>$y = \frac{2a}{\pi} \tan^{-1} \left( \frac{\sinh 2\eta}{\sin 2\psi} \right)$<br>$z = z$ | $g_{11} = g_{22} = \left( \frac{4a}{\pi} \right)^2$<br>$\times \frac{1}{\sinh^2 2\eta + \sin^2 2\psi}$<br>$g_{33} = 1$   | $\times$            | $\times$  | $\bullet$ | $\bullet$                   |
| L 3 C | $z = \frac{2a}{\pi} \ln \cosh w$    | 2.13        | $x = \frac{a}{\pi} \ln (\cosh^2 \eta - \sin^2 \psi)$<br>$y = \frac{2a}{\pi} \tan^{-1} (\tanh \eta \tan \psi)$<br>$z = z$  | $g_{11} = g_{22}$<br>$= \left( \frac{2a}{\pi} \right)^2 \left\{ \frac{\cosh^2 \eta \sinh^2 \eta}{(\cosh^2 \eta + [\sinh \eta \cosh \eta + \sin \psi \cos \psi]^2)} \right.$<br>$\left. - \frac{\sin^2 \psi}{\sin^2 \eta} \right\}$<br>$g_{33} = 1$ | $\times$            | $\times$  | $\bullet$ | $\bullet$                   |
| H 1 C | Elliptic-cylinder<br>coordinates    | 2.14        | $x = a \cosh \eta \cos \psi$<br>$y = a \sinh \eta \sin \psi$<br>$z = z$   | $g_{11} = g_{22}$<br>$= a^2 (\cosh^2 \eta - \cos^2 \psi)$<br>$g_{33} = 1$  | $\bullet$           | $\bullet$ | $\bullet$ | $\times$                    |
| H 2 C | $\bar{z} = a \operatorname{sech} w$ | 2.15        | $x = \frac{a \cosh \eta \cos \psi}{\cosh^2 \eta - \sin^2 \psi}$<br>$y = \frac{a \sinh \eta \sin \psi}{\cosh^2 \eta - \sin^2 \psi}$<br>$z = z$   | $g_{11} = g_{22} = \frac{a^2 (\cosh^2 \eta - \cos^2 \psi)}{(\cosh^2 \eta - \sin^2 \psi)^2}$<br>$g_{33} = 1$  | $\times$            | $\times$  | $\bullet$ | $\bullet$                   |

Table 3.01. Some cylindrical coordinate systems

|       |                                   |      |   |  |   |   |   |   |
|-------|-----------------------------------|------|---|--|---|---|---|---|
| J 1 C | $z = a \operatorname{sn} w$       | 2.16 | $x = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} v$<br>$y = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} v \operatorname{cn} v$<br>$z = z$<br>where<br>$A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 v$ | $g_{11} = g_{22} = \frac{a^2 \Omega^2}{A^2}$<br>$g_{33} = 1$<br>$\Omega^2 = (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 v)$<br>$\times (\operatorname{dn}^2 v - k^2 \operatorname{sn}^2 \mu)$   | ● | ● | × | × |
| J 2 C | $\bar{z} = a \operatorname{cn} w$ | 2.17 | $x = \frac{a}{A} \operatorname{cn} \mu \operatorname{cn} v$<br>$y = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \mu \operatorname{sn} v \operatorname{dn} v$<br>$z = z$   | $g_{11} = g_{22} = \frac{a^2 T^2}{A^2}$<br>$g_{33} = 1$<br>$T^2 = (\operatorname{sn}^2 v + \operatorname{sn}^2 \mu \operatorname{cn}^2 v)$<br>$\times (\operatorname{dn}^2 v - k^2 \operatorname{sn}^2 \mu)$   | ● | ● | × | × |
| J 3 C |                                   | 2.18 | $x = \frac{A}{a T} \operatorname{sn} \mu \operatorname{dn} v$<br>$y = \frac{k^{\frac{1}{2}} \Pi}{2 a T}$<br>$z = z$   | $g_{11} = g_{22} = \frac{A^2 \Omega^2}{a^2 T^2}$<br>$g_{33} = 1$   | ● | ● | × | × |
|       |                                   |      | $\bar{z} = \frac{k^{\frac{1}{2}}}{i 2 a}$<br>$\times \left( \frac{1 + i k^{\frac{1}{2}} \operatorname{sn} w}{1 - i k^{\frac{1}{2}} \operatorname{sn} w} \right)$  | $\mathcal{T} = \operatorname{sn}^2 \mu \operatorname{dn}^2 v + [(\Lambda / \bar{k})$<br>$+ \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} v \operatorname{cn} v]^2$<br>$\Pi = (\Lambda^2 / \bar{k}) - (\operatorname{sn}^2 \mu \operatorname{dn}^2 v)$<br>$+ \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 v \operatorname{cn}^2 v$ |   |   |   |   |

## Section III. Cylindrical systems

Table 3.01. Continuation

| No.   | Transformation<br>Fig.<br>No.                                     | Equations  | Metric Coefficients   | Simple Separability         |                    | One-dimensional<br>solution |
|-------|---|--|---|-----------------------------|--------------------|-----------------------------|
|       |   |  |   | HELMHOLTZ<br>HELM.<br>HOTLZ | LAPLACE<br>LAPLACE |                             |
| J 4 C | $\bar{z} = \frac{a}{\pi} \ln \left( \frac{1}{k \sin^2 w} \right)$ | $x = \frac{a}{\pi} \ln \left( \frac{A^2}{k \Sigma} \right)$<br>$y = \frac{2a}{\pi} \times \tan^{-1} \left( \frac{\operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu}{\operatorname{sn} \mu \operatorname{dn} \nu} \right)$<br>$z = z$ | $g_{11} = g_{22} = \left( \frac{2a \Theta}{\pi \Sigma} \right)^2$<br>$g_{33} = 1$<br>where<br>$\Sigma = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu$<br>$\Theta^2 = \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \times (\operatorname{cn}^2 \nu - \operatorname{sn}^2 \nu \operatorname{dn}^2 \nu)^2 + \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu \operatorname{dn}^2 \nu + (\operatorname{cm}^2 \mu + \operatorname{sn}^2 \mu \operatorname{dn}^2 \mu)^2$                           | ●                           | ×                  |                             |
| J 5 C | $\bar{z} = \frac{2a}{\pi} \ln \operatorname{cn} w$                | $x = \frac{a}{\pi} \ln (\mathcal{E}/A^2)$<br>$y = \frac{2a}{\pi} \times \tan^{-1} \left( \frac{\operatorname{sn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{dn} \nu}{\operatorname{cn} \mu \operatorname{cn} \nu} \right)$<br>$z = z$                   | $g_{11} = g_{22} = \left( \frac{2a \Theta'}{\pi \mathcal{E}} \right)^2$<br>$g_{33} = 1$<br>where<br>$\mathcal{E} = \operatorname{cn}^2 \mu \operatorname{cn}^2 \nu + \operatorname{sn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{dn}^2 \nu$<br>$\Theta'^2 = \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \times (\operatorname{dn}^2 \nu - k'^2 \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu)^2 + \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu \operatorname{dn}^2 \nu \times (\operatorname{dn}^2 \mu - k^2 \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu)^2$ | ●                           | ×                  |                             |

|       |  |   |           |          |          |   |          |
|-------|--|---|-----------|----------|----------|---|----------|
|       | $x = \frac{2Ka}{\pi} \left\{ Z(\mu) + \frac{k^2}{A'} \right.$<br>$\times \operatorname{sn}\mu \operatorname{cn}\mu \operatorname{dn}\mu \operatorname{sn}^2\nu' \left. \right\}$                                     |   |           |          |          |   |          |
|       | $y = \frac{2Ka}{\pi} \left\{ Z(\nu') \right.$<br>$+ \frac{\pi\nu}{2KK'} - \frac{1}{A'} \right.$<br>$\times \operatorname{dn}^2\mu \operatorname{sn}\nu' \operatorname{cn}\nu' \operatorname{dn}\nu' \left. \right\}$ | $g_{11} = g_{22} = \left( \frac{2Ka}{\pi} \right)^2 \left\{ \frac{4k^4}{A'^4} \right.$<br>$\times \operatorname{sn}^2\mu \operatorname{cn}^2\mu \operatorname{dn}^2\mu \operatorname{sn}^2\nu'$<br>$\times \operatorname{cn}^2\nu' \operatorname{dn}^2\nu' + \left[ \operatorname{dn}^2\mu - \frac{E}{K} \right]$<br>$+ \frac{k^2 \operatorname{sn}^2\nu'}{A'^2} (\operatorname{dn}^2\mu \operatorname{cn}^2\nu'$<br>$\times (\operatorname{cn}^2\mu - \operatorname{sn}^2\mu \operatorname{dn}^2\mu)$<br>$- k^2 \operatorname{sn}^2\mu (1 - k^2 \operatorname{sn}^4\mu)) \left. \right ^2 \left\} \right.$ | $\bullet$ | $\times$ | $\times$ | $\times$  | $\times$ |
| J 6 C | $\bar{z} = \frac{2Ka}{\pi}$<br>$\times Z(w + iK')$<br>$+ ia.$  | $z = z$<br>2.21<br>where<br>$A' = 1 - \operatorname{dn}^2\mu \operatorname{sn}^2\nu'$<br>$g_{33} = 1$<br>$\nu' = \nu + K'$  | $2.21$    | $\times$ | $\times$ | $K, K',$<br>and $E$<br>are complete elliptic<br>integrals |          |

Crosses indicate non-separability, dots indicate separability. The modulus  $k$  is associated with the variable  $\mu$ , and  $k'$  is associated with  $\nu$ : e.g.,  $\operatorname{sn}\mu \equiv \operatorname{sn}(\mu, k)$ ,  $\operatorname{sn}\nu \equiv \operatorname{sn}(\nu, k')$ .

**TABLE 3.02. IMPORTANT EQUATIONS FOR CYLINDRICAL SYSTEMS****P 1C. TANGENT-CYLINDER COORDINATES ( $\mu, \nu, z$ ), Fig. 2.01.**

$$\begin{cases} x = \frac{\mu}{\mu^2 + \nu^2}, \\ y = \frac{\nu}{\mu^2 + \nu^2}, \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = (\mu^2 + \nu^2)^{-2}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{1}{(\mu^2 + \nu^2)^2} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = (\mu^2 + \nu^2) \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = (\mu^2 + \nu^2)^2 \left[ \frac{\partial}{\partial \mu} \left( \frac{E_\mu}{\mu^2 + \nu^2} \right) + \frac{\partial}{\partial \nu} \left( \frac{E_\nu}{\mu^2 + \nu^2} \right) \right] + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\mu \left[ (\mu^2 + \nu^2) \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - (\mu^2 + \nu^2) \frac{\partial E_z}{\partial \mu} \right] \\ & + \mathbf{a}_z (\mu^2 + \nu^2)^2 \left[ \frac{\partial}{\partial \mu} \left( \frac{E_\nu}{\mu^2 + \nu^2} \right) - \frac{\partial}{\partial \nu} \left( \frac{E_\mu}{\mu^2 + \nu^2} \right) \right]. \end{aligned}$$

$$\nabla^2 \varphi = (\mu^2 + \nu^2)^2 \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**P 2C. PARABOLIC-CYLINDER COORDINATES (see Section I).****P 3C. CARDIOID-CYLINDER-COORDINATES ( $\mu, \nu, z$ ), Fig. 2.03.**

$$\begin{cases} x = \frac{1}{2} \frac{\mu^2 - \nu^2}{(\mu^2 + \nu^2)^2}, \\ y = \frac{\mu \nu}{(\mu^2 + \nu^2)^2}, \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = (\mu^2 + \nu^2)^{-3}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{1}{(\mu^2 + \nu^2)^3} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = (\mu^2 + \nu^2)^{\frac{1}{2}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = (\mu^2 + \nu^2)^3 \left\{ \frac{\partial}{\partial \mu} [(\mu^2 + \nu^2)^{-\frac{1}{2}} E_\mu] + \frac{\partial}{\partial \nu} [(\mu^2 + \nu^2)^{-\frac{1}{2}} E_\nu] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\mu \left[ (\mu^2 + \nu^2)^{\frac{1}{2}} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - (\mu^2 + \nu^2)^{\frac{1}{2}} \frac{\partial E_z}{\partial \mu} \right] \\ & + \mathbf{a}_z (\mu^2 + \nu^2)^3 \left[ \frac{\partial}{\partial \mu} ((\mu^2 + \nu^2)^{-\frac{1}{2}} E_\nu) - \frac{\partial}{\partial \nu} ((\mu^2 + \nu^2)^{-\frac{1}{2}} E_\mu) \right]. \end{aligned}$$

$$\nabla^2 \varphi = (\mu^2 + \nu^2)^3 \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**P 4C. HYPERBOLIC-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.04.**

$$\begin{cases} x = (\varrho + \mu)^{\frac{1}{2}}, \\ y = (\varrho - \mu)^{\frac{1}{2}}, \\ z = z, \quad \text{where } \varrho = +(\mu^2 + \nu^2)^{\frac{1}{2}}. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{1}{2}(\mu^2 + \nu^2)^{-\frac{1}{2}}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{1}{2(\mu^2 + \nu^2)^{\frac{1}{2}}} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{2}} \left\{ \frac{\partial}{\partial \mu} [(\mu^2 + \nu^2)^{-\frac{1}{4}} E_\mu] + \frac{\partial}{\partial \nu} [(\mu^2 + \nu^2)^{-\frac{1}{4}} E_\nu] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\mu \left[ \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_v}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \frac{\partial E_z}{\partial \mu} \right] \\ & + \mathbf{a}_z \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \left[ \frac{\partial}{\partial \mu} ((\mu^2 + \nu^2)^{-\frac{1}{4}} E_\nu) - \frac{\partial}{\partial \nu} ((\mu^2 + \nu^2)^{-\frac{1}{4}} E_\mu) \right]. \end{aligned}$$

$$\nabla^2 \varphi = 2(\mu^2 + \nu^2)^{\frac{1}{2}} \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**P 5C. ROSE-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.05.**

$$\begin{cases} x = \frac{1}{\varrho} (\varrho + \mu)^{\frac{1}{2}}, \\ y = \frac{1}{\varrho} (\varrho - \mu)^{\frac{1}{2}}, \\ z = z, \quad \text{where } \varrho = +(\mu^2 + \nu^2)^{\frac{1}{2}}. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{1}{2(\mu^2 + \nu^2)^{\frac{3}{2}}}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{1}{2(\mu^2 + \nu^2)^{\frac{3}{2}}} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \sqrt{2}(\mu^2 + \nu^2)^{\frac{3}{2}} \left\{ \frac{\partial}{\partial \mu} [(\mu^2 + \nu^2)^{-\frac{1}{4}} E_\mu] + \frac{\partial}{\partial \nu} [(\mu^2 + \nu^2)^{-\frac{1}{4}} E_\nu] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\mu \left[ \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_v}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \frac{\partial E_z}{\partial \mu} \right] \\ & + \mathbf{a}_z \sqrt{2}(\mu^2 + \nu^2)^{\frac{1}{4}} \left[ \frac{\partial}{\partial \mu} ((\mu^2 + \nu^2)^{-\frac{1}{4}} E_\nu) - \frac{\partial}{\partial \nu} ((\mu^2 + \nu^2)^{-\frac{1}{4}} E_\mu) \right]. \end{aligned}$$

$$\nabla^2 \varphi = 2(\mu^2 + \nu^2)^{\frac{3}{2}} \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**E 1C. CIRCULAR-CYLINDER COORDINATES** (see Section I).

**E 2C. CASSINIAN-OVAL COORDINATES** ( $\eta, \psi, z$ ), Fig. 2.07.

$$\begin{cases} x = \frac{a}{\sqrt{2}} [\varrho + (e^\eta \cos \psi + 1)]^{\frac{1}{2}}, \\ y = \frac{a}{\sqrt{2}} [\varrho - (e^\eta \cos \psi + 1)]^{\frac{1}{2}}, \\ z = z, \quad \text{where } \varrho = + (e^{2\eta} + 2e^\eta \cos \psi + 1)^{\frac{1}{2}}. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{a^2 e^{2\eta}}{4\varrho}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{a^2 e^{2\eta}}{4\varrho} [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{2\varrho^{\frac{1}{2}}}{a e^\eta} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{2\varrho}{a e^{2\eta}} \left\{ \frac{\partial}{\partial \eta} \left[ \frac{e^\eta}{\sqrt{\varrho}} E_\eta \right] + \frac{\partial}{\partial \psi} \left[ \frac{e^\eta}{\sqrt{\varrho}} E_\psi \right] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\eta \left[ \frac{2\varrho^{\frac{1}{2}}}{a e^\eta} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right] + \mathbf{a}_\psi \left[ \frac{\partial E_\eta}{\partial z} - \frac{2\varrho^{\frac{1}{2}}}{a e^\eta} \frac{\partial E_z}{\partial \eta} \right] \\ & + \mathbf{a}_z \frac{2\varrho}{a e^{2\eta}} \left[ \frac{\partial}{\partial \eta} \left( \frac{e^\eta}{\sqrt{\varrho}} E_\psi \right) - \frac{\partial}{\partial \psi} \left( \frac{e^\eta}{\sqrt{\varrho}} E_\eta \right) \right]. \end{aligned}$$

$$\nabla^2 \varphi = \frac{4\varrho}{a^2 e^{2\eta}} \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**E 3C. INVERSE CASSINIAN-OVAL COORDINATES** ( $\eta, \psi, z$ ), Fig. 2.08.

$$\begin{cases} x = \frac{a}{\sqrt{2}\varrho} [\varrho + (e^\eta \cos \psi + 1)]^{\frac{1}{2}}, \\ y = \frac{a}{\sqrt{2}\varrho} [\varrho - (e^\eta \cos \psi + 1)]^{\frac{1}{2}}, \\ z = z, \quad \text{where } \varrho = + (e^{2\eta} + 2e^\eta \cos \psi + 1)^{\frac{1}{2}}. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{a^2 e^{2\eta}}{4\varrho^3}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{a^2 e^{2\eta}}{4\varrho^3} [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{2\varrho^{\frac{3}{2}}}{a e^\eta} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{2\varrho^3}{a e^{2\eta}} \left\{ \frac{\partial}{\partial \eta} \left[ \frac{e^\eta}{\varrho^{\frac{3}{2}}} E_\eta \right] + \frac{\partial}{\partial \psi} \left[ \frac{e^\eta}{\varrho^{\frac{3}{2}}} E_\psi \right] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\eta \left[ \frac{2\varrho^{\frac{3}{2}}}{a e^\eta} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right] + \mathbf{a}_\psi \left[ \frac{\partial E_\eta}{\partial z} - \frac{2\varrho^{\frac{3}{2}}}{a e^\eta} \frac{\partial E_z}{\partial \eta} \right] \\ & + \mathbf{a}_z \frac{2\varrho^3}{a e^{2\eta}} \left[ \frac{\partial}{\partial \eta} \left( \frac{e^\eta}{\varrho^{\frac{3}{2}}} E_\psi \right) - \frac{\partial}{\partial \psi} \left( \frac{e^\eta}{\varrho^{\frac{3}{2}}} E_\eta \right) \right]. \end{aligned}$$

$$\nabla^2 \varphi = \frac{4\varrho^3}{a^2 e^{2\eta}} \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**E 4C. BI-CYLINDER COORDINATES  $(\eta, \psi, z)$ , Fig. 2.09.**

$$\begin{cases} x = \frac{a \sinh \eta}{\cosh \eta - \cos \psi}, \\ y = \frac{a \sin \psi}{\cosh \eta - \cos \psi}, \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^1 = \frac{a^2}{(\cosh \eta - \cos \psi)^2}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{a^2}{(\cosh \eta - \cos \psi)^2} [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\text{grad } \varphi = \frac{1}{a} (\cosh \eta - \cos \psi) \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{1}{a} (\cosh \eta - \cos \psi)^2 \\ &\times \left\{ \frac{\partial}{\partial \eta} [(\cosh \eta - \cos \psi)^{-1} E_\eta] + \frac{\partial}{\partial \psi} [(\cosh \eta - \cos \psi)^{-1} E_\psi] \right\} + \frac{\partial E_z}{\partial z}. \end{aligned}$$

$$\begin{aligned} \text{curl } \mathbf{E} &= \mathbf{a}_\eta \left[ \frac{1}{a} (\cosh \eta - \cos \psi) \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right] + \mathbf{a}_\psi \left[ \frac{\partial E_\eta}{\partial z} - \frac{1}{a} (\cosh \eta - \cos \psi) \frac{\partial E_z}{\partial \eta} \right] \\ &+ \mathbf{a}_z (\cosh \eta - \cos \psi)^2 \left[ \frac{\partial}{\partial \eta} \left( \frac{E_\psi}{\cosh \eta - \cos \psi} \right) - \frac{\partial}{\partial \psi} \left( \frac{E_\eta}{\cosh \eta - \cos \psi} \right) \right]. \end{aligned}$$

$$\nabla^2 \varphi = \frac{1}{a^2} (\cosh \eta - \cos \psi)^2 \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**E 5C. MAXWELL-CYLINDER COORDINATES  $(\eta, \psi, z)$ , Fig. 2.10.**

$$\begin{cases} x = \frac{a}{\pi} (\eta + 1 + e^\eta \cos \psi), \\ y = \frac{a}{\pi} (\psi + e^\eta \sin \psi), \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^1 = \left( \frac{a}{\pi} \right)^2 (1 + 2e^\eta \cos \psi + e^{2\eta}), \quad g_{33} = 1.$$

$$(ds)^2 = \left( \frac{a}{\pi} \right)^2 (1 + 2e^\eta \cos \psi + e^{2\eta}) [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\text{grad } \varphi = \frac{\pi}{a} (1 + 2e^\eta \cos \psi + e^{2\eta})^{-\frac{1}{2}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{\pi}{a} (1 + 2e^\eta \cos \psi + e^{2\eta})^{-1} \\ &\times \left\{ \frac{\partial}{\partial \eta} [(1 + 2e^\eta \cos \psi + e^{2\eta})^{\frac{1}{2}} E_\eta] + \frac{\partial}{\partial \psi} [(1 + 2e^\eta \cos \psi + e^{2\eta})^{\frac{1}{2}} E_\psi] \right\} + \frac{\partial E_z}{\partial z}. \end{aligned}$$

$$\begin{aligned} \text{curl } \mathbf{E} &= \mathbf{a}_\eta \left[ \frac{\pi}{a} (1 + 2e^\eta \cos \psi + e^{2\eta})^{-\frac{1}{2}} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right] \\ &+ \mathbf{a}_\psi \left[ \frac{\partial E_\eta}{\partial z} - \frac{\pi}{a} (1 + 2e^\eta \cos \psi + e^{2\eta})^{-\frac{1}{2}} \frac{\partial E_z}{\partial \eta} \right] + \mathbf{a}_z \frac{\pi}{a} (1 + 2e^\eta \cos \psi + e^{2\eta})^{-1} \\ &\times \left[ \frac{\partial}{\partial \eta} ((1 + 2e^\eta \cos \psi + e^{2\eta})^{\frac{1}{2}} E_\psi) - \frac{\partial}{\partial \psi} ((1 + 2e^\eta \cos \psi + e^{2\eta})^{\frac{1}{2}} E_\eta) \right]. \end{aligned}$$

$$\nabla^2 \varphi = \left( \frac{\pi}{a} \right)^2 (1 + 2e^\eta \cos \psi + e^{2\eta})^{-1} \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**L 1C. LOGARITHMIC-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.11.**

$$\begin{cases} x = \frac{a}{\pi} \ln(\mu^2 + \nu^2), \\ y = \frac{2a}{\pi} \tan^{-1}(\nu/\mu), \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \left( \frac{2a}{\pi} \right)^2 \frac{1}{\mu^2 + \nu^2}, \quad g_{33} = 1.$$

$$(ds)^2 = \left( \frac{2a}{\pi} \right)^2 \frac{1}{\mu^2 + \nu^2} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{\pi}{2a} (\mu^2 + \nu^2)^{\frac{1}{2}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{\pi}{2a} (\mu^2 + \nu^2) \left\{ \frac{\partial}{\partial \mu} [(\mu^2 + \nu^2)^{-\frac{1}{2}} E_\mu] + \frac{\partial}{\partial \nu} [(\mu^2 + \nu^2)^{-\frac{1}{2}} E_\nu] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} &= \mathbf{a}_\mu \left[ \frac{\pi}{2a} (\mu^2 + \nu^2)^{\frac{1}{2}} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \frac{\pi}{2a} (\mu^2 + \nu^2)^{\frac{1}{2}} \frac{\partial E_z}{\partial \mu} \right] \\ &\quad + \mathbf{a}_z \frac{\pi}{2a} (\mu^2 + \nu^2) \left[ \frac{\partial}{\partial \mu} \left( \frac{E_\nu}{(\mu^2 + \nu^2)^{\frac{1}{2}}} \right) - \frac{\partial}{\partial \nu} \left( \frac{E_\mu}{(\mu^2 + \nu^2)^{\frac{1}{2}}} \right) \right]. \end{aligned}$$

$$\nabla^2 \varphi = \left( \frac{\pi}{2a} \right)^2 (\mu^2 + \nu^2) \left[ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**L 2C. LN TAN-CYLINDER COORDINATES  $(\eta, \psi, z)$ , Fig. 2.12.**

$$\begin{cases} x = \frac{a}{\pi} \ln \left[ \frac{\sinh^2 \eta + \sin^2 \psi}{\sinh^2 \eta + \cos^2 \psi} \right], \\ y = \frac{2a}{\pi} \tan^{-1} \left( \frac{\sinh 2\eta}{\sin 2\psi} \right), \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \left( \frac{4a}{\pi} \right)^2 \frac{1}{\sinh^2 2\eta + \sin^2 2\psi}, \quad g_{33} = 1.$$

$$(ds)^2 = \left( \frac{4a}{\pi} \right)^2 \frac{1}{\sinh^2 2\eta + \sin^2 2\psi} [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{\pi}{4a} [\sinh^2 2\eta + \sin^2 2\psi]^{\frac{1}{2}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\begin{aligned} \operatorname{div} \mathbf{E} &= \frac{\pi}{4a} [\sinh^2 2\eta + \sin^2 2\psi] \left\{ \frac{\partial}{\partial \eta} [(\sinh^2 2\eta + \sin^2 2\psi)^{-\frac{1}{2}} E_\eta] \right. \\ &\quad \left. + \frac{\partial}{\partial \psi} [(\sinh^2 2\eta + \sin^2 2\psi)^{-\frac{1}{2}} E_\psi] \right\} + \frac{\partial E_z}{\partial z}. \end{aligned}$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} &= \mathbf{a}_\eta \left[ \frac{\pi}{4a} (\sinh^2 2\eta + \sin^2 2\psi)^{\frac{1}{2}} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right] \\ &\quad + \mathbf{a}_\psi \left[ \frac{\partial E_\eta}{\partial z} - \frac{\pi}{4a} (\sinh^2 2\eta + \sin^2 2\psi)^{\frac{1}{2}} \frac{\partial E_z}{\partial \eta} \right] + \mathbf{a}_z \frac{\pi}{4a} (\sinh^2 2\eta + \sin^2 2\psi) \\ &\quad \times \left[ \frac{\partial}{\partial \eta} ((\sinh^2 2\eta + \sin^2 2\psi)^{-\frac{1}{2}} E_\psi) - \frac{\partial}{\partial \psi} ((\sinh^2 2\eta + \sin^2 2\psi)^{-\frac{1}{2}} E_\eta) \right]. \end{aligned}$$

$$\nabla^2 \varphi = \left( \frac{\pi}{4a} \right)^2 (\sinh^2 2\eta + \sin^2 2\psi) \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**L 3C. LN COSH-CYLINDER COORDINATES  $(\eta, \psi, z)$ , Fig. 2.13.**

$$\begin{cases} x = \frac{a}{\pi} \ln(\cosh^2 \eta - \sin^2 \psi), \\ y = \frac{2a}{\pi} \tan^{-1}(\tanh \eta \tan \psi), \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \left( \frac{2a \Pi}{\pi} \right)^2, \quad g_{33} = 1.$$

where

$$\Pi^2 = \frac{\cosh^2 \eta \sinh^2 \eta + (\sinh \eta \cosh \eta + \sin \psi \cos \psi)^2}{(\cosh^2 \eta - \sin^2 \psi)^2}.$$

$$(ds)^2 = \left( \frac{2a \Pi}{\pi} \right)^2 [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\text{grad } \varphi = \frac{\pi}{2a \Pi} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\text{div } \mathbf{E} = \frac{\pi}{2a \Pi^2} \left\{ \frac{\partial}{\partial \eta} [\Pi E_\eta] + \frac{\partial}{\partial \psi} [\Pi E_\psi] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \text{curl } \mathbf{E} = & \mathbf{a}_\eta \left[ \frac{\pi}{2a \Pi} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right] + \mathbf{a}_\psi \left[ \frac{\partial E_\eta}{\partial z} - \frac{\pi}{2a \Pi} \frac{\partial E_z}{\partial \eta} \right] \\ & + \mathbf{a}_z \frac{\pi}{2a \Pi^2} \left[ \frac{\partial}{\partial \eta} [\Pi E_\psi] - \frac{\partial}{\partial \psi} [\Pi E_\eta] \right]. \end{aligned}$$

$$\nabla^2 \varphi = \left( \frac{\pi}{2a \Pi} \right)^2 \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.$$

**E 1C. ELLIPTIC-CYLINDER COORDINATES (see Section I).****H 2C. INVERSE ELLIPTIC-CYLINDER COORDINATES  $(\eta, \psi, z)$ , Fig. 2.15.**

$$\begin{cases} x = \frac{a \cosh \eta \cos \psi}{\cosh^2 \eta - \sin^2 \psi}, \\ y = \frac{a \sinh \eta \sin \psi}{\cosh^2 \eta - \sin^2 \psi}, \\ z = z. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{a^2 (\cosh^2 \eta - \cos^2 \psi)}{(\cosh^2 \eta - \sin^2 \psi)^2}, \quad g_{33} = 1.$$

$$(ds)^2 = \frac{a^2 (\cosh^2 \eta - \cos^2 \psi)}{(\cosh^2 \eta - \sin^2 \psi)^2} [(d\eta)^2 + (d\psi)^2] + (dz)^2.$$

$$\text{grad } \varphi = \frac{\cosh^2 \eta - \sin^2 \psi}{a (\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\psi \frac{\partial \varphi}{\partial \psi} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\begin{aligned} \text{div } \mathbf{E} = & \frac{(\cosh^2 \eta - \sin^2 \psi)^2}{a (\cosh^2 \eta - \cos^2 \psi)} \\ & \times \left\{ \frac{\partial}{\partial \eta} \left[ \frac{(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}}{\cosh^2 \eta - \sin^2 \psi} E_\eta \right] + \frac{\partial}{\partial \psi} \left[ \frac{(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}}{\cosh^2 \eta - \sin^2 \psi} E_\psi \right] \right\} + \frac{\partial E_z}{\partial z}. \end{aligned}$$

$$\begin{aligned}\operatorname{curl} \mathbf{E} &= \mathbf{a}_\eta \left\{ \frac{\cosh^2 \eta - \sin^2 \psi}{a(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z} \right\} \\ &\quad + \mathbf{a}_\psi \left\{ \frac{\partial E_\eta}{\partial z} - \frac{\cosh^2 \eta - \sin^2 \psi}{a(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}} \frac{\partial E_z}{\partial \eta} \right\} + \mathbf{a}_z \frac{(\cosh^2 \eta - \sin^2 \psi)^2}{a(\cosh^2 \eta - \cos^2 \psi)} \\ &\quad \times \left\{ \frac{\partial}{\partial \eta} \left[ \frac{(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}}{\cosh^2 \eta - \sin^2 \psi} E_\psi \right] - \frac{\partial}{\partial \psi} \left[ \frac{(\cosh^2 \eta - \cos^2 \psi)^{\frac{1}{2}}}{\cosh^2 \eta - \sin^2 \psi} E_\eta \right] \right\}. \\ \nabla^2 \varphi &= \frac{(\cosh^2 \eta - \sin^2 \psi)^2}{a^2(\cosh^2 \eta - \cos^2 \psi)} \left[ \frac{\partial^2 \varphi}{\partial \eta^2} + \frac{\partial^2 \varphi}{\partial \psi^2} \right] + \frac{\partial^2 \varphi}{\partial z^2}.\end{aligned}$$

J 1C. SN-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.16.

$$0 \leqq \mu \leqq K, \quad 0 \leqq \nu \leqq K', \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \nu, \\ y = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu, \\ z = z, \quad \text{where } A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{a^2 \Omega^2}{A^2}, \quad g_{33} = 1,$$

where

$$\Omega^2 = (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu) (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu).$$

$$(ds)^2 = \frac{a^2 \Omega^2}{A^2} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{A}{a \Omega} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{A^2}{a \Omega^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Omega}{A} E_\mu \right] + \frac{\partial}{\partial \nu} \left[ \frac{\Omega}{A} E_\nu \right] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned}\operatorname{curl} \mathbf{E} &= \mathbf{a}_\mu \left[ \frac{A}{a \Omega} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \frac{A}{a \Omega} \frac{\partial E_z}{\partial \mu} \right] \\ &\quad + \mathbf{a}_z \frac{A^2}{a \Omega^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Omega}{A} E_\nu \right] - \frac{\partial}{\partial \nu} \left[ \frac{\Omega}{A} E_\mu \right] \right\}.\end{aligned}$$

$$\nabla^2 \varphi = \frac{A^2}{a^2 \Omega^2} \left\{ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right\} + \frac{\partial^2 \varphi}{\partial z^2}.$$

J 2C. CN-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.17.

$$0 \leqq \mu \leqq K, \quad 0 \leqq \nu \leqq K', \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{a}{A} \operatorname{cn} \mu \operatorname{cn} \nu, \\ y = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{dn} \nu, \\ z = z, \quad \text{where } A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu. \end{cases}$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \frac{a^2 \Gamma^2}{A^2}, \quad g_{33} = 1,$$

where

$$\Gamma^2 = (\operatorname{sn}^2 \nu + \operatorname{sn}^2 \mu \operatorname{cn}^2 \nu) (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu).$$

$$(ds)^2 = \frac{a^2 \Gamma^2}{A^2} [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{A}{a \Gamma} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{A^2}{a \Gamma^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Gamma}{A} E_\mu \right] + \frac{\partial}{\partial \nu} \left[ \frac{\Gamma}{A} E_\nu \right] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} &= \mathbf{a}_\mu \left[ \frac{A}{a \Gamma} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \frac{A}{a \Gamma} \frac{\partial E_z}{\partial \mu} \right] \\ &\quad + \mathbf{a}_z \frac{A^2}{a \Gamma^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Gamma}{A} E_\nu \right] - \frac{\partial}{\partial \nu} \left[ \frac{\Gamma}{A} E_\mu \right] \right\}. \end{aligned}$$

$$\nabla^2 \varphi = \frac{A^2}{a^2 \Gamma^2} \left\{ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right\} + \frac{\partial^2 \varphi}{\partial z^2}.$$

J 3C. INVERSE SN-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.18.

$$0 \leqq \mu \leqq K, \quad 0 \leqq \nu \leqq K', \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{A}{a \mathcal{T}} \operatorname{sn} \mu \operatorname{dn} \nu, \\ y = \frac{k^{\frac{1}{2}} \Pi}{2a \mathcal{T}}, \\ z = z, \quad \text{where } A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu, \end{cases}$$

$$\Omega = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu,$$

$$\mathcal{T} = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + [(A/\sqrt{k}) + \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu],$$

$$\Pi = (A^2/k) - (\operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu).$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \left( \frac{A \Omega}{a \mathcal{T}} \right)^2, \quad g_{33} = 1.$$

$$(ds)^2 = \left( \frac{A \Omega}{a \mathcal{T}} \right)^2 [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{a \mathcal{T}}{A \Omega} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{a \mathcal{T}^2}{A^2 \Omega^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{A \Omega}{\mathcal{T}} E_\mu \right] + \frac{\partial}{\partial \nu} \left[ \frac{A \Omega}{\mathcal{T}} E_\nu \right] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} &= \mathbf{a}_\mu \left[ \frac{a \mathcal{T}}{A \Omega} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \frac{a \mathcal{T}}{A \Omega} \frac{\partial E_z}{\partial \mu} \right] \\ &\quad + \mathbf{a}_z \frac{a \mathcal{T}^2}{A^2 \Omega^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{A \Omega}{\mathcal{T}} E_\nu \right] - \frac{\partial}{\partial \nu} \left[ \frac{A \Omega}{\mathcal{T}} E_\mu \right] \right\}. \end{aligned}$$

$$\nabla^2 \varphi = \left( \frac{a \mathcal{T}}{A \Omega} \right)^2 \left\{ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right\} + \frac{\partial^2 \varphi}{\partial z^2}.$$

**J 4C. LN SN-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.19.**

$$0 \leq \mu \leq K, \quad 0 \leq \nu \leq K', \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{a}{\pi} \ln \left( \frac{\Lambda^2}{k \Sigma} \right), \\ y = \frac{2a}{\pi} \tan^{-1} \left( \frac{\operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu}{\operatorname{sn} \mu \operatorname{dn} \nu} \right), \\ z = z, \quad \text{where } \Lambda = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu, \end{cases}$$

$$\Sigma = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu.$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \left( \frac{2a\Theta}{\pi\Sigma} \right)^2, \quad g_{33} = 1,$$

where

$$\Theta^2 = \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu (\operatorname{cn}^2 \nu - \operatorname{sn}^2 \nu \operatorname{dn}^2 \nu)^2$$

$$+ \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu \operatorname{dn}^2 \nu (\operatorname{cn}^2 \mu + \operatorname{sn}^2 \mu \operatorname{dn}^2 \mu)^2.$$

$$(ds)^2 = \left( \frac{2a\Theta}{\pi\Sigma} \right)^2 [(d\mu)^2 + (d\nu)^2] + (dz)^2.$$

$$\operatorname{grad} \varphi = \frac{\pi\Sigma}{2a\Theta} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}.$$

$$\operatorname{div} \mathbf{E} = \frac{\pi\Sigma^2}{2a\Theta^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Theta}{\Sigma} E_\mu \right] + \frac{\partial}{\partial \nu} \left[ \frac{\Theta}{\Sigma} E_\nu \right] \right\} + \frac{\partial E_z}{\partial z}.$$

$$\begin{aligned} \operatorname{curl} \mathbf{E} = & \mathbf{a}_\mu \left[ \frac{\pi\Sigma}{2a\Theta} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \frac{\pi\Sigma}{2a\Theta} \frac{\partial E_z}{\partial \mu} \right] \\ & + \mathbf{a}_z \frac{\pi\Sigma^2}{2a\Theta^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Theta}{\Sigma} E_\nu \right] - \frac{\partial}{\partial \nu} \left[ \frac{\Theta}{\Sigma} E_\mu \right] \right\}. \end{aligned}$$

$$V^2 \varphi = \left( \frac{\pi\Sigma}{2a\Theta} \right)^2 \left\{ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right\} + \frac{\partial^2 \varphi}{\partial z^2}.$$

**J 5C. LN CN-CYLINDER COORDINATES  $(\mu, \nu, z)$ , Fig. 2.20.**

$$0 \leq \mu \leq K, \quad 0 \leq \nu \leq K', \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{a}{\pi} \ln (\Xi/\Lambda^2), \\ y = \frac{2a}{\pi} \tan^{-1} \left( \frac{\operatorname{sn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{dn} \nu}{\operatorname{cn} \mu \operatorname{cn} \nu} \right), \\ z = z, \quad \text{where } \Lambda = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu, \end{cases}$$

$$\Xi = \operatorname{cn}^2 \mu \operatorname{cn}^2 \nu + \operatorname{sn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{dn}^2 \nu.$$

$$g_{11} = g_{22} = g^{\frac{1}{2}} = \left( \frac{2a\Theta'}{\pi\Xi} \right)^2, \quad g_{33} = 1,$$

where

$$\Theta'^2 = \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu (\operatorname{dn}^2 \nu - k'^2 \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu)^2$$

$$+ \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu \operatorname{dn}^2 \nu (\operatorname{dn}^2 \mu - k^2 \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu)^2.$$

$$\begin{aligned}
 (ds)^2 &= \left( \frac{2a\Theta'}{\pi\Xi} \right)^2 [(d\mu)^2 + (d\nu)^2] + (dz)^2. \\
 \text{grad } \varphi &= \frac{\pi\Xi}{2a\Theta'} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_z \frac{\partial \varphi}{\partial z}. \\
 \text{div } \mathbf{E} &= \frac{\pi\Xi^2}{2a\Theta'^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Theta'}{\Xi} E_\mu \right] + \frac{\partial}{\partial \nu} \left[ \frac{\Theta'}{\Xi} E_\nu \right] \right\} + \frac{\partial E_z}{\partial z}. \\
 \text{curl } \mathbf{E} &= \mathbf{a}_\mu \left[ \frac{\pi\Xi}{2a\Theta'} \frac{\partial E_z}{\partial \nu} - \frac{\partial E_\nu}{\partial z} \right] + \mathbf{a}_\nu \left[ \frac{\partial E_\mu}{\partial z} - \frac{\pi\Xi}{2a\Theta'} \frac{\partial E_z}{\partial \mu} \right] \\
 &\quad + \mathbf{a}_z \frac{\pi\Xi^2}{2a\Theta'^2} \left\{ \frac{\partial}{\partial \mu} \left[ \frac{\Theta'}{\Xi} E_\nu \right] - \frac{\partial}{\partial \nu} \left[ \frac{\Theta'}{\Xi} E_\mu \right] \right\}. \\
 \nabla^2 \varphi &= \left( \frac{\pi\Xi}{2a\Theta'} \right)^2 \left\{ \frac{\partial^2 \varphi}{\partial \mu^2} + \frac{\partial^2 \varphi}{\partial \nu^2} \right\} + \frac{\partial^2 \varphi}{\partial z^2}.
 \end{aligned}$$

J 6C. ZETA-CYLINDER COORDINATES ( $\mu, \nu, z$ ), Fig. 2.21.

$$0 \leq \mu \leq K, \quad 0 \leq \nu \leq K', \quad -\infty < z < +\infty.$$

$$\begin{cases} x = \frac{2Ka}{\pi} \left\{ Z(\mu) + \frac{k^2}{A'} \operatorname{sn} \mu \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn}^2 \nu' \right\}, \\ y = \frac{2Ka}{\pi} \left\{ Z(\nu') + \frac{\pi\nu}{2KK'} - \frac{1}{A'} \operatorname{dn}^2 \mu \operatorname{sn} \nu' \operatorname{cn} \nu' \operatorname{dn} \nu' \right\}, \\ z = z, \quad \text{where } A' = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu', \end{cases}$$

$$\nu' = \nu + K',$$

$$K = \int_0^{\pi/2} \frac{d\varphi}{(1 - k^2 \sin^2 \varphi)^{\frac{1}{2}}},$$

$$K' = \int_0^{\pi/2} \frac{d\varphi}{(1 - k'^2 \sin^2 \varphi)^{\frac{1}{2}}},$$

$$E = \int_0^{\pi/2} (1 - k^2 \sin^2 \varphi)^{\frac{1}{2}} d\varphi.$$

$$\begin{aligned}
 g_{11} = g_{22} = g^{\frac{1}{2}} &= \left( \frac{2Ka}{\pi} \right)^2 \left\{ \frac{4k^4}{A'^4} \operatorname{sn}^2 \mu \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu' \operatorname{cn}^2 \nu' \operatorname{dn}^2 \nu' \right. \\
 &\quad + \left[ \operatorname{dn}^2 \mu - \frac{E}{K} + \frac{k^2 \operatorname{sn}^2 \nu'}{A'^2} (\operatorname{dn}^2 \mu \operatorname{cn}^2 \nu' [\operatorname{cn}^2 \mu - \operatorname{sn}^2 \mu \operatorname{dn}^2 \mu] \right. \\
 &\quad \left. \left. - k^2 \operatorname{sn}^2 \mu [1 - k^2 \operatorname{sn}^4 \mu]) \right]^2 \right\}.
 \end{aligned}$$

Other equations can be written with the help of the general relations of § 3.01.

## Section IV

### ROTATIONAL SYSTEMS

Each of the 21 transformations of Section II yields one or two rotational coordinate systems, obtained by twirling the plane map about an axis. A tabulation of the most interesting of these systems is given in this section.

#### 4.01 R-SEPARATION

In most cases, the rotational coordinate systems do not allow simple separation of the Laplace and Helmholtz equations, though they may allow *R*-separation of the Laplace equation. This subject is treated in previous publications [7], [9].

*Definition I.* If the assumption

$$\varphi = U^1(u^1) \cdot U^2(u^2) \cdot U^3(u^3)$$

permits the separation of the partial differential equation into three ordinary differential equations, the equation is said to be *simply separable*.

*Definition II.* If the assumption

$$\varphi = \frac{U^1(u^1) \cdot U^2(u^2) \cdot U^3(u^3)}{R(u^1, u^2, u^3)}$$

permits the separation of the partial differential equation into three ordinary differential equations, and if  $R \neq \text{const}$ , the equation is said to be *R-separable*.

*R*-separation introduces two new quantities,  $R(u^1, u^2, u^3)$  and  $Q(u^1, u^2, u^3)$ , which greatly improve the possibility of effecting separation. These quantities are defined by the relations,

$$\begin{cases} g_{ii} = \frac{S}{M_{i1}} Q, \\ \frac{g^{\frac{1}{2}}}{S} = f_1 f_2 f_3 (R)^2 Q. \end{cases} \quad (4.01)$$

Also,  $\alpha_1$  is not zero as it was in simple separation of the Laplace equation, but

$$\alpha_1 = -\frac{Q}{R} \sum_{i=1}^3 \frac{1}{f_i g_{ii}} \frac{\partial}{\partial u^i} \left( f_i \frac{\partial R}{\partial u^i} \right) = \text{const}. \quad (4.02)$$

Equation (4.03) constitutes the necessary and sufficient condition for separability of the Laplace equation in euclidean 3-space [7]. A comparison of the equations for simple separation and *R*-separation is given in Table 4.01.

No case is known in which the Helmholtz equation is *R*-separable, so the question that arises in this section is merely whether the Laplace equation is

$R$ -separable in a given rotational coordinate system. For example, in toroidal coordinates,

$$g_{11} = g_{22} = \frac{a^2}{(\cosh \eta - \cos \theta)^2},$$

$$g_{33} = \frac{a^2 \sinh^2 \eta}{(\cosh \eta - \cos \theta)^2}.$$

The complicated form of the denominator precludes the possibility of obtaining a Stäckel matrix for simple separation. But for  $R$ -separation, we let

$$Q = \frac{a^2}{(\cosh \eta - \cos \theta)^2}.$$

Then

$$\frac{M_{11}}{S} = \frac{M_{21}}{S} = 1, \quad \frac{M_{31}}{S} = \frac{1}{\sinh^2 \eta},$$

and the Stäckel determinant is very simple:

$$S = \begin{vmatrix} 1 & -1 & -\frac{1}{\sinh^2 \eta} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1.$$

Also,  $\alpha_1 = +\frac{1}{4}$  and  $R^2 = (\cosh \eta - \cos \theta)^{-1}$ . Requirements for  $R$ -separation of the Laplace equation are satisfied.

## 4.02 TABLES

Metric coefficients and separability data are listed in Table 4.02. Some of the rotational systems obtained from Figs. 2.01 to 2.21 do not allow separation of any kind. After discarding the non-separable cases, we find 4 systems in which Laplace's equation is simply separable and 10 that are  $R$ -separable. The simply separable coordinates have already been listed in Section I, and the  $R$ -separable cases are tabulated in this section. One additional system is introduced—the inversion of rectangular coordinates—making a total of 11 coordinate systems treated in Section 4:

| No.    | Name                                   | Fig. No. |
|--------|--|----------|
| P 1 R  | Tangent-sphere coordinates             | 4.01     |
| P 3 R  | Cardioid coordinates                   | 4.02     |
| E 4 Rx | Bispherical coordinates                | 4.03     |
| E 4 Ry | Toroidal coordinates                   | 4.04     |
| H 2 Rx | Inverse prolate spheroidal coordinates | 4.05     |
| H 2 Ry | Inverse oblate spheroidal coordinates  | 4.06     |
| — —    | 6-sphere coordinates                   | 4.07     |
| J 1 Rx | Bi-cyclide coordinates                 | 4.08     |
| J 1 Ry | Flat-ring cyclide coordinates          | 4.09     |
| J 2 R  | Disk-cyclide coordinates               | 4.10     |
| J 3 R  | Cap-cyclide coordinates                | 4.11     |

Details of these eleven coordinate systems are given in Table 4.03 and Figs. 4.01 to 4.11. Table 4.04 summarizes four rotational systems obtained from elliptic functions. It should be emphasized that these coordinates by no means constitute the totality of all  $R$ -separable systems.

TABLE 4.01. SEPARABILITY

for  $\varphi = \varphi(u^1, u^2, u^3)$ .

| Simple separability<br>of Helmholtz equation | $R$ -separability<br>of Laplace equation   |
|--|--|
| $\nabla^2 \varphi + \kappa^2 \varphi = 0$    | $\nabla^2 \varphi = 0$   |
| $\varphi = U^1 \cdot U^2 \cdot U^3$          | $\varphi = \frac{U^1 \cdot U^2 \cdot U^3}{R}$  |
| $g_{ii} = \frac{S}{M_{i1}}$                  | $g_{ii} = \frac{S}{M_{i1}} Q$  |
| $\frac{g^{\frac{1}{2}}}{S} = f_1 f_2 f_3$    | $\frac{g^{\frac{1}{2}}}{S} = f_1 f_2 f_3 (R)^2 Q$  |
| $\alpha_1 = \kappa^2$                        | $\alpha_1 = -\frac{Q}{R} \sum_{i=1}^3 \frac{1}{f_i g_{ii}} \frac{\partial}{\partial u^i} \left( f_i \frac{\partial R}{\partial u^i} \right)$ |

### Separated equations

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0.$$

### Stäckel matrix

$$[S] = \begin{bmatrix} \Phi_{11}(u^1) & \Phi_{12}(u^1) & \Phi_{13}(u^1) \\ \Phi_{21}(u^2) & \Phi_{22}(u^2) & \Phi_{23}(u^2) \\ \Phi_{31}(u^3) & \Phi_{32}(u^3) & \Phi_{33}(u^3) \end{bmatrix}.$$

Table 4.02. Some rotational coordinate systems

TABLE 4.02. SOME ROTATIONAL COORDINATE SYSTEMS

| No.                | Name<br>of Coordinate<br>System | Fig.<br>No. | Equations  | Metric Coefficients  | Separability |         | One-dimensional<br>Solution |          |
|--------------------|---------------------------------|-------------|--|--|--------------|---------|-----------------------------|----------|
|                    |                                 |             |  |  | HELMHOLTZ    | LAPLACE |                             |          |
| P <sub>1</sub> R   | Tangent-sphere                  | 4.01        | $x = \frac{\mu}{\mu^2 + \nu^2} \cos \psi$<br>Rot. about $y$ axis<br>$y = \frac{\mu}{\mu^2 + \nu^2} \sin \psi$<br>$z = \frac{\nu}{\mu^2 + \nu^2}$   | $g_{11} = g_{22} = (\mu^2 + \nu^2)^{-2}$<br>$g_{33} = \frac{\mu^2}{(\mu^2 + \nu^2)^2}$       | $\times$     | R       | $\times$                    | $\times$ |
| P <sub>2</sub> R   | Parabolic                       | 4.08        | $x = \mu \nu \cos \psi$<br>Rot. about $x$ axis<br>$y = \mu \nu \sin \psi$<br>$z = \frac{1}{2} (\mu^2 - \nu^2)$   | $g_{11} = g_{33} = \mu^2 + \nu^2$<br>$g_{33} = \mu^2 \nu^2$                                  | S            | S       | $\times$                    | ●<br>0   |
| * P <sub>3</sub> R | Cardioid                        | 4.02        | $x = \frac{\mu \nu}{(\mu^2 + \nu^2)^2} \cos \psi$<br>Rot. about $x$ axis<br>$y = \frac{\mu \nu}{(\mu^2 + \nu^2)^2} \sin \psi$<br>$z = \frac{1}{2} \frac{\mu^2 - \nu^2}{(\mu^2 + \nu^2)^2}$ | $g_{11} = g_{22} = (\mu^2 + \nu^2)^{-3}$<br>$g_{33} = \frac{\mu^2 \nu^2}{(\mu^2 + \nu^2)^4}$ | $\times$     | R       | $\times$                    | 0        |

Table 4.02. Continuation

| No.    | Name<br>of Coordinate<br>System | Fig.<br>No. | Equations  | Metric Coefficients   | Separability   |          | One-dimensional<br>Solution | $\alpha_1$ |
|--------|---------------------------------|-------------|--|---|----------------|----------|-----------------------------|------------|
|        |                                 |             |  |   | HELMHOLTZ      | LAPLACE  |                             |            |
|        |                                 |             |  |   | HELM.<br>HOLTZ | LAPLACE  |                             |            |
| P 4 R  | Hyperbolic                      | 2.04        | $x = (\varrho + \mu)^{\frac{1}{2}} \cos \psi$<br>$y = (\varrho + \mu)^{\frac{1}{2}} \sin \psi$<br>$z = (\varrho - \mu)^{\frac{1}{2}}$<br>where<br>$\varrho = +(\mu^2 + \nu^2)^{\frac{1}{2}}$ | $g_{11} = g_{22} = \frac{1}{2}(\mu^2 + \nu^2)^{-\frac{1}{2}}$<br>$g_{33} = \varrho + \mu$   | $\times$       | $\times$ | $\times$                    | $\times$   |
| E 4 R  | Spherical                       | 1.05        | $x = e^\eta \sin \theta \cos \psi$<br>$y = e^\eta \sin \theta \sin \psi$<br>$z = e^\eta \cos \theta$   | $g_{11} = g_{22} = e^{2\eta}$<br>$g_{33} = e^{2\eta} \sin^2 \theta$   | $S$            | $S$      | $\bullet$                   | 0          |
| E 4 Rx | Bispherical                     | 4.03        | $x = \frac{a \sin \theta \cos \psi}{\cosh \eta - \cos \theta}$<br>$y = \frac{a \sin \theta \sin \psi}{\cosh \eta - \cos \theta}$<br>$z = \frac{a \sinh \eta}{\cosh \eta - \cos \theta}$      | $g_{11} = g_{22} = \frac{a^2}{(\cosh \eta - \cos \theta)^2}$<br>$g_{33} = \frac{a^2 \sin^2 \theta}{(\cosh \eta - \cos \theta)^2}$ | $\times$       | $\times$ | R                           | $\times$   |

Table 4.02. Some rotational coordinate systems

|        |                                       |                                       |   |   |  |             |                                  |                                  |
|--------|---------------------------------------|---------------------------------------|---|---|--|-------------|----------------------------------|----------------------------------|
| E 4 Ry | Toroidal                              | 4.04<br>Rot.<br>about<br>$y$<br>axis  | $x = \frac{a \sinh \eta \cos \psi}{\cosh \eta - \cos \theta}$<br>$y = \frac{a \sinh \eta \sin \psi}{\cosh \eta - \cos \theta}$<br>$z = \frac{a \sin \theta}{\cosh \eta - \cos \theta}$  | $g_{11} = g_{22} = \frac{a^2}{(\cosh \eta - \cos \theta)^2}$<br>$g_{33} = \frac{a^2 \sinh^2 \eta}{(\cosh \eta - \cos \theta)^2}$  | $\times$<br>$\times$<br>$\times$   | R<br>R<br>R | $\times$<br>$\times$<br>$\times$ | $\times$<br>$\times$<br>$\times$ |
|        | H 1 Rx                                | Prolate<br>spheroidal                 | 1.06<br>Rot.<br>about<br>$x$<br>axis  | $x = a \sinh \eta \sin \theta \cos \psi$<br>$y = a \sinh \eta \sin \theta \sin \psi$<br>$z = a \cosh \eta \cos \theta$  | $g_{11} = g_{22} = \frac{a^2 (\cosh^2 \eta - \cos^2 \theta)}{a^2 (\sinh^2 \eta + \sin^2 \theta)}$<br>$g_{33} = a^2 \sinh^2 \eta \sin^2 \theta$ | S<br>S<br>S | S<br>S<br>S                      | $\bullet$<br>$\bullet$<br>0      |
|        |                                       |                                       |   |   | S<br>S<br>S  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |
|        |                                       |                                       | 1.07<br>Rot.<br>about<br>$y$<br>axis*   | $x = a \cosh \eta \sin \theta \cos \psi$<br>$y = a \cosh \eta \sin \theta \sin \psi$<br>$z = a \sinh \eta \cos \theta$  | $g_{11} = g_{22} = \frac{a^2 (\cosh^2 \eta - \sin^2 \theta)}{a^2 \cosh^2 \eta \sin^2 \theta}$<br>$g_{33} = a^2 \sinh^2 \eta \sin^2 \theta$     | S<br>S<br>S | S<br>S<br>S                      | $\bullet$<br>$\bullet$<br>0      |
|        |                                       |                                       |   |   |  | S<br>S<br>S | S<br>S<br>S                      | $\bullet$<br>$\bullet$<br>0      |
|        |                                       |                                       |   |   |  | S<br>S<br>S | S<br>S<br>S                      | $\bullet$<br>$\bullet$<br>0      |
| H 1 Ry | Oblate<br>spheroidal                  | 4.05<br>Rot.<br>about<br>$y$<br>axis* | $x = \frac{a \sinh \eta \sin \theta \cos \psi}{\cosh^2 \eta - \sin^2 \theta}$<br>$y = \frac{a \sinh \eta \sin \theta \sin \psi}{\cosh^2 \eta - \sin^2 \theta}$<br>$z = \frac{a \cosh \eta \cos \theta}{\cosh^2 \eta - \sin^2 \theta}$ | $g_{11} = g_{22} = \frac{a^2 (\sinh^2 \eta + \sin^2 \theta)}{(\cosh^2 \eta - \sin^2 \theta)^2}$<br>$g_{33} = \frac{a^2 \sinh^2 \eta \sin^2 \theta}{(\cosh^2 \eta - \sin^2 \theta)^2}$ | S<br>S<br>S  | R<br>R<br>R | $\times$<br>$\times$<br>$\times$ | $\times$<br>$\times$<br>$\times$ |
|        | S<br>S<br>S                           |                                       |   |   |  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |
|        | S<br>S<br>S                           |                                       |   |   |  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |
| H 2 Rx | Inversion<br>of prolate<br>spheroidal | 4.06<br>Rot.<br>about<br>$y$<br>axis* | $x = \frac{a \cosh \eta \sin \theta \cos \psi}{\cosh^2 \eta - \cos^2 \theta}$<br>$y = \frac{a \cosh \eta \sin \theta \sin \psi}{\cosh^2 \eta - \cos^2 \theta}$<br>$z = \frac{a \sinh \eta \cos \theta}{\cosh^2 \eta - \cos^2 \theta}$ | $g_{11} = g_{22} = \frac{a^2 (\cosh^2 \eta - \sin^2 \theta)}{(\cosh^2 \eta - \cos^2 \theta)^2}$<br>$g_{33} = \frac{a^2 \cosh^2 \eta \sin^2 \theta}{(\cosh^2 \eta - \cos^2 \theta)^2}$ | S<br>S<br>S  | R<br>R<br>R | $\times$<br>$\times$<br>$\times$ | $\times$<br>$\times$<br>$\times$ |
|        | S<br>S<br>S                           |                                       |   |   |  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |
|        | S<br>S<br>S                           |                                       |   |   |  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |
| H 2 Ry | Inversion<br>of oblate<br>spheroidal  | 4.06<br>Rot.<br>about<br>$y$<br>axis* | $x = \frac{a \cosh \eta \sin \theta \cos \psi}{\cosh^2 \eta - \cos^2 \theta}$<br>$y = \frac{a \cosh \eta \sin \theta \sin \psi}{\cosh^2 \eta - \cos^2 \theta}$<br>$z = \frac{a \sinh \eta \cos \theta}{\cosh^2 \eta - \cos^2 \theta}$ | $g_{11} = g_{22} = \frac{a^2 (\cosh^2 \eta - \cos^2 \theta)}{(\cosh^2 \eta - \cos^2 \theta)^2}$<br>$g_{33} = \frac{a^2 \cosh^2 \eta \sin^2 \theta}{(\cosh^2 \eta - \cos^2 \theta)^2}$ | S<br>S<br>S  | R<br>R<br>R | $\times$<br>$\times$<br>$\times$ | $\times$<br>$\times$<br>$\times$ |
|        | S<br>S<br>S                           |                                       |   |   |  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |
|        | S<br>S<br>S                           |                                       |   |   |  | S<br>S<br>S | $\bullet$<br>$\bullet$<br>0      |                                  |

\*  $\theta = v - \pi/2$  in this case, to make the equations agree with usual oblate nomenclature.

Table 4.02. Continuation

| No.  | Name<br>of Coordinate<br>System | Fig.<br>No.    | Equations   | Metric Coefficients  | Separability |         |         | One-dimensional<br>Solution | $\alpha_1$ |
|------|---------------------------------|----------------|---|--|--------------|---------|---------|-----------------------------|------------|
|      |                                 |                |   |  | HELMHOLTZ    | LAPLACE | LAPLACE |                             |            |
| 4.08 | J 1 Rx                          | $z = a \sin w$ | $x = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu$<br>$\times \operatorname{sn} \nu \operatorname{cn} \nu \cos \psi$<br>$y = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu$<br>$\times \operatorname{sn} \nu \operatorname{cn} \nu \sin \psi$<br>$z = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \nu$<br>where<br>$A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu$ | $g_{11} = g_{22} = \frac{a^2 \Omega^2}{A^2}$<br>$g_{33} = \frac{a^2}{A^2} \operatorname{cn}^2 \mu \operatorname{dn}^2 \nu$<br>$\times \operatorname{sn}^2 \nu \operatorname{cn}^2 \psi$<br>where<br>$\Omega^2 = (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu)$<br>$\times (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu)$ | R            | R       | R       | X                           | -2         |
| 4.09 | J 1 Ry                          | $z = a \sin w$ | $x = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \nu \cos \psi$<br>$y = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \nu \sin \psi$<br>$z = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu$<br>$\times \operatorname{sn} \nu \operatorname{cn} \nu$   | $g_{11} = g_{22} = \frac{a^2 \Omega^2}{A^2}$<br>$g_{33} = \frac{a^2}{A^2} \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu$   | X            | X       | R       | R                           | -1         |

Table 4.02. Some rotational coordinate systems

|  |  |                               |   |  |   |   |   |   |     |
|--|--|-------------------------------|---|--|---|---|---|---|-----|
|  | $\bar{z} = a \operatorname{cn} w$  | 4.10<br>Rot.<br>about<br>axis | $x = \frac{a}{A} \operatorname{cn} \mu \operatorname{cn} \nu \cos \psi$   | $g_{11} = g_{22} = \frac{a^2 I^2}{A^2}$  | R | R | X | X | — 1 |
|  |  |                               | $y = \frac{a}{A} \operatorname{cn} \mu \operatorname{cn} \nu \sin \psi$   | $g_{33} = \frac{a^2}{A^2} \operatorname{cn}^2 \mu \operatorname{cn}^2 \nu$<br>where  |   |   |   |   |     |
|  | $\bar{z} = \frac{k^{\frac{3}{2}}}{i 2a} \sqrt{\left( \frac{1 + ik^{\frac{3}{2}} \operatorname{sn} w}{1 - ik^{\frac{3}{2}} \operatorname{sn} w} \right)}$ | 4.11<br>Rot.<br>about<br>axis | $z = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \mu$<br>$\times \operatorname{sn} \nu \operatorname{dn} \nu$ | $I^2 = (\operatorname{sn}^2 \nu + \operatorname{sn}^2 \mu \operatorname{cn}^2 \nu)$<br>$\times (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu)$  | R | R | X | X | — 1 |
|  |  |                               |   |  |   |   |   |   |     |
|  | $\bar{z} = \frac{k^{\frac{3}{2}}}{i 2a} \sqrt{\left( \frac{1 + ik^{\frac{3}{2}} \operatorname{sn} w}{1 - ik^{\frac{3}{2}} \operatorname{sn} w} \right)}$ | 4.11<br>Rot.<br>about<br>axis | $x = \frac{A}{a \mathcal{T}} \operatorname{sn} \mu \operatorname{dn} \nu \cos \psi$                                   | $g_{11} = g_{22} = \frac{A^2 \Omega^2}{a^2 \mathcal{T}^2}$   | R | R | X | X | — 1 |
|  |  |                               | $y = \frac{A}{a \mathcal{T}} \operatorname{sn} \mu \operatorname{dn} \nu \sin \psi$                                   | $g_{33} = \frac{A^2}{a^2 \mathcal{T}^2} \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu$<br>where<br>$\mathcal{T} = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu$<br>$z = \frac{k^{\frac{3}{2}} \Pi}{2a \mathcal{T}}$ |   |   |   |   |     |

Crosses indicate non-separability,  $S$  indicates simple separability,  $R$  indicates  $R$ -separability. The modulus  $k$  is associated with  $\mu, k'$  is associated with  $\nu$ :  $\operatorname{sn} \mu \equiv \operatorname{sn}(\mu, k)$ ,  $\operatorname{sn} \nu \equiv \operatorname{sn}(\nu, k')$ .

TABLE 4.03

P 1R. TANGENT-SPHERE COORDINATES  $(\mu, \nu, \psi)$ , Fig. 4.01.




Fig. 4.01. Tangent-sphere coordinates  $(\mu, \nu, \psi)$ . This and the following figures are drawn as if the coordinate surfaces were filled with jelly and then one quarter were cut out (from  $\psi = 0$  to  $\psi = \pi/2$ ). The coordinate surfaces are tangent spheres ( $\nu = \pm \text{const}$ ), toroids without center opening ( $\mu = \text{const}$ ), and half-planes ( $\psi = \text{const}$ ). The solid arrow heads indicate the direction associated with an increase in the parameter

$$\begin{aligned} 0 &< \mu < +\infty, \\ -\infty &< \nu < +\infty, \\ 0 &\leq \psi < 2\pi. \end{aligned}$$

$$\left\{ \begin{array}{l} x = \frac{\mu \cos \psi}{\mu^2 + \nu^2}, \\ y = \frac{\mu \sin \psi}{\mu^2 + \nu^2}, \\ z = \frac{\nu}{\mu^2 + \nu^2}. \end{array} \right.$$

Coordinate surfaces

$$\left\{ \begin{array}{l} x^2 + y^2 + z^2 = \frac{1}{\mu} (x^2 + y^2)^{\frac{1}{2}} \\ \quad (\text{toroids without center opening, } \mu = \text{const}), \\ x^2 + y^2 + \left(z - \frac{1}{2\nu}\right)^2 = \frac{1}{4\nu^2} \\ \quad (\text{spheres tangent to } xy\text{-plane at origin, } \nu = \text{const}), \\ \tan \psi = y/x \\ \quad (\text{half planes, } \psi = \text{const}). \end{array} \right.$$

Stäckel matrix

$$[S] = \begin{bmatrix} 1 & -1 & -1/\mu^2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = 1,$$

$$M_{11} = 1, \quad M_{21} = 1, \quad M_{31} = 1/\mu^2.$$

### Metric coefficients

$$g_{11} = g_{22} = 1/(\mu^2 + \nu^2)^2, \quad g_{33} = \mu^2/(\mu^2 + \nu^2)^2, \quad g^{\frac{1}{2}} = \mu/(\mu^2 + \nu^2)^{\frac{3}{2}}.$$

$$R = (\mu^2 + \nu^2)^{-\frac{1}{2}}, \quad Q = (\mu^2 + \nu^2)^{-2},$$

$$f_1 = \mu, \quad f_2 = 1, \quad f_3 = 1, \quad \alpha_1 = 0.$$

### Important equations

$$(ds)^2 = \frac{(d\mu)^2 + (d\nu)^2 + \mu^2(d\psi)^2}{(\mu^2 + \nu^2)^2}.$$

$$\text{grad } \varphi = (\mu^2 + \nu^2) \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} + \frac{\mathbf{a}_\psi}{\mu} \frac{\partial \varphi}{\partial \psi} \right].$$

$$\text{div } \mathbf{E} = \frac{(\mu^2 + \nu^2)^3}{\mu} \left[ \frac{\partial}{\partial \mu} \left( \frac{\mu}{(\mu^2 + \nu^2)^2} E_\mu \right) + \mu \frac{\partial}{\partial \nu} \left( \frac{1}{(\mu^2 + \nu^2)^2} E_\nu \right) \right] + \frac{1}{\mu} (\mu^2 + \nu^2) \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{(\mu^2 + \nu^2)^2}{\mu} \begin{vmatrix} \mathbf{a}_\mu & \mathbf{a}_\nu & \mathbf{a}_\psi \mu \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ \frac{E_\mu}{\mu^2 + \nu^2} & \frac{E_\nu}{\mu^2 + \nu^2} & \frac{E_\psi \mu}{\mu^2 + \nu^2} \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{(\mu^2 + \nu^2)^3}{\mu} \left\{ \frac{\partial}{\partial \mu} \left( \frac{\mu}{\mu^2 + \nu^2} \frac{\partial \varphi}{\partial \mu} \right) + \mu \frac{\partial}{\partial \nu} \left( \frac{1}{\mu^2 + \nu^2} \frac{\partial \varphi}{\partial \nu} \right) \right\} + \frac{(\mu^2 + \nu^2)^2}{\mu^2} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{d u^i} \left( f_i \frac{d U^i}{d u^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = (\mu^2 + \nu^2)^{\frac{1}{2}} M \cdot N \cdot \Psi.$$

General case

$$\begin{cases} \frac{d^2 M}{d \mu^2} + \frac{1}{\mu} \frac{d M}{d \mu} - \left( \alpha_2 + \frac{\alpha_3}{\mu^2} \right) M = 0, \\ \frac{d^2 N}{d \nu^2} + \alpha_2 N = 0, \\ \frac{d^2 \Psi}{d \psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

$$If \alpha_2 = -q^2 \text{ and } \alpha_3 = p^2,$$

$$\frac{d^2 M}{d \mu^2} + \frac{1}{\mu} \frac{d M}{d \mu} + (q^2 - p^2/\mu^2) M = 0, \quad \{24\}^* \quad M = A \mathcal{J}_p(q\mu) + B \mathcal{J}_{-p}(q\mu)$$

$$\text{or } ** \quad M = A \mathcal{I}_n(q\mu) + B \mathcal{Y}_n(q\mu).$$

$$\frac{d^2 N}{d \nu^2} - q^2 N = 0, \quad \{04\} \quad N = A e^{q\nu} + B e^{-q\nu}.$$

$$\frac{d^2 \Psi}{d \psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

$$If \alpha_2 = +q^2 \text{ and } \alpha_3 = p^2,$$

$$\frac{d^2 M}{d \mu^2} + \frac{1}{\mu} \frac{d M}{d \mu} - (q^2 + p^2/\mu^2) M = 0, \quad \{24\} \quad M = A \mathcal{J}_p(i q \mu) + B \mathcal{J}_{-p}(i q \mu)$$

$$\text{or } ** \quad M = A \mathcal{I}_n(i q \mu) + B \mathcal{Y}_n(i q \mu).$$

$$\frac{d^2 N}{d \nu^2} + q^2 N = 0, \quad \{04\} \quad N = A \sin q\nu + B \cos q\nu.$$

$$\frac{d^2 \Psi}{d \psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

\* Designation in terms of singularities. See Section VI.

\*\* If  $p = n$ , an integer, use the second solution.

$$\text{If } \alpha_2 = 0 \text{ and } \alpha_3 = p^2,$$

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \frac{p^2}{\mu^2} M = 0, \quad \{04\} \quad M = A \mu^p + B \mu^{-p}.$$

$$\frac{d^2N}{d\nu^2} = 0, \quad \{04\} \quad N = A + B \nu.$$

$$\frac{d^2\Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p \psi + B \cos p \psi.$$

$$\text{If } \alpha_2 = \alpha_3 = 0,$$

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} = 0, \quad \{04\} \quad M = A + B \ln \mu.$$

$$\frac{d^2N}{d\nu^2} = 0, \quad \{04\} \quad N = A + B \nu.$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B \psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \alpha_2 M = 0, \\ \frac{d^2N}{d\nu^2} + \alpha_2 N = 0. \end{cases}$$

$$\text{If } \alpha_2 = -q^2,$$

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + q^2 M = 0, \quad \{14\} \quad M = A \mathcal{I}_0(q \mu) + B \mathcal{Y}_0(q \mu).$$

$$\frac{d^2N}{d\nu^2} - q^2 N = 0, \quad \{04\} \quad N = A e^{q\nu} + B e^{-q\nu}.$$

$$\text{If } \alpha_2 = +q^2,$$

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - q^2 M = 0, \quad \{14\} \quad M = A \mathcal{I}_0(i q \mu) + B \mathcal{Y}_0(i q \mu).$$

$$\frac{d^2N}{d\nu^2} + q^2 N = 0, \quad \{04\} \quad N = A \sin q \nu + B \cos q \nu.$$

$$\text{If } \alpha_2 = 0,$$

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} = 0, \quad \{04\} \quad M = A + B \ln \mu.$$

$$\frac{d^2N}{d\nu^2} = 0, \quad \{04\} \quad N = A + B \nu.$$

**P 3R. CARDIOID COORDINATES  $(\mu, \nu, \psi)$ , Fig. 4.02.**

$$0 \leq \mu < \infty,$$

$$0 \leq \nu < \infty,$$

$$0 \leq \psi < 2\pi.$$

$$\begin{cases} x = \frac{\mu\nu}{(\mu^2 + \nu^2)^2} \cos \psi, \\ y = \frac{\mu\nu}{(\mu^2 + \nu^2)^2} \sin \psi, \\ z = \frac{1}{2} \frac{(\mu^2 - \nu^2)}{(\mu^2 + \nu^2)^2}. \end{cases}$$

**Coordinate surfaces**

$$\begin{cases} x^2 + y^2 + z^2 \\ = \frac{1}{4\mu^2} [(x^2 + y^2 + z^2)^{\frac{1}{2}} + z] \\ (\text{cardioids of revolution intersecting positive half of } z\text{-axis, } \mu = \text{const}), \\ x^2 + y^2 + z^2 \\ = \frac{1}{4\nu^2} [(x^2 + y^2 + z^2)^{\frac{1}{2}} - z] \\ (\text{cardioids of revolution intersecting negative half of } z\text{-axis, } \nu = \text{const}), \\ \tan \psi = y/x \\ (\text{half planes, } \psi = \text{const}). \end{cases}$$




Fig. 4.02. Cardioid coordinates  $(\mu, \nu, \psi)$ . The coordinate surfaces are cardioids of revolution ( $\mu = \text{const}$  and  $\nu = \text{const}$ ), and half-planes ( $\psi = \text{const}$ )

**Stäckel matrix**

$$[S] = \begin{bmatrix} \mu^2 & -1 & -1/\mu^2 \\ \nu^2 & 1 & -1/\nu^2 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = \mu^2 + \nu^2, \quad M_{11} = 1, \quad M_{21} = 1, \quad M_{31} = \frac{\mu^2 + \nu^2}{\mu^2 \nu^2}.$$

**Metric coefficients**

$$g_{11} = g_{22} = 1/(\mu^2 + \nu^2)^3, \quad g_{33} = \mu^2 \nu^2 / (\mu^2 + \nu^2)^4, \quad g^{\frac{1}{2}} = \mu \nu / (\mu^2 + \nu^2)^{\frac{5}{2}}.$$

$$R = 1/(\mu^2 + \nu^2), \quad Q = 1/(\mu^2 + \nu^2)^4,$$

$$f_1 = \mu, \quad f_2 = \nu, \quad f_3 = 1, \quad \alpha_1 = 0.$$

**Important equations**

$$(ds)^2 = \frac{1}{(\mu^2 + \nu^2)^3} [(d\mu)^2 + (d\nu)^2] + \frac{\mu^2 \nu^2}{(\mu^2 + \nu^2)^4} (d\psi)^2.$$

$$\text{grad } \varphi = (\mu^2 + \nu^2)^{\frac{3}{2}} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_\psi \frac{(\mu^2 + \nu^2)}{\mu \nu} \frac{\partial \varphi}{\partial \psi}.$$

$$\text{div } \mathbf{E} = \frac{(\mu^2 + \nu^2)^5}{\mu \nu} \left[ \nu \frac{\partial}{\partial \mu} \left( \frac{\mu}{(\mu^2 + \nu^2)^{\frac{5}{2}}} E_\mu \right) + \mu \frac{\partial}{\partial \nu} \left( \frac{\nu}{(\mu^2 + \nu^2)^{\frac{5}{2}}} E_\nu \right) \right] + \frac{(\mu^2 + \nu^2)^2}{\mu \nu} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{(\mu^2 + \nu^2)^{\frac{7}{2}}}{\mu \nu} \begin{vmatrix} \mathbf{a}_\mu & \mathbf{a}_\nu & \frac{\mathbf{a}_\psi \mu \nu}{(\mu^2 + \nu^2)^{\frac{1}{2}}} \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ \frac{E_\mu}{(\mu^2 + \nu^2)^{\frac{3}{2}}} & \frac{E_\nu}{(\mu^2 + \nu^2)^{\frac{3}{2}}} & \frac{E_\psi \mu \nu}{(\mu^2 + \nu^2)^2} \end{vmatrix},$$

$$\nabla^2 \varphi = \frac{(\mu^2 + \nu^2)^5}{\mu \nu} \times \left\{ \nu \frac{\partial}{\partial \mu} \left( \frac{\mu}{(\mu^2 + \nu^2)^2} \frac{\partial \varphi}{\partial \mu} \right) + \mu \frac{\partial}{\partial \nu} \left( \frac{\nu}{(\mu^2 + \nu^2)^2} \frac{\partial \varphi}{\partial \nu} \right) \right\} + \frac{(\mu^2 + \nu^2)^4}{\mu^2 \nu^2} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = (\mu^2 + \nu^2) M \cdot N \cdot \Psi.$$

General case

$$\begin{cases} \frac{d^2 M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \left( \alpha_2 + \frac{\alpha_3}{\mu^2} \right) M = 0, \\ \frac{d^2 N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + \left( \alpha_2 - \frac{\alpha_3}{\nu^2} \right) N = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

$$If \alpha_2 = q^2 \text{ and } \alpha_3 = p^2,$$

$$\frac{d^2 M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - (q^2 + p^2/\mu^2) M = 0, \quad \{24\} \quad M = A \mathcal{J}_p(i q \mu) + B \mathcal{J}_{-p}(i q \mu) \\ \text{or*} \quad M = A \mathcal{J}_n(i q \mu) + B \mathcal{Y}_n(i q \mu).$$

$$\frac{d^2 N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + (q^2 - p^2/\nu^2) N = 0, \quad \{24\} \quad N = A \mathcal{J}_p(q \nu) + B \mathcal{J}_{-p}(q \nu) \\ \text{or*} \quad N = A \mathcal{J}_n(q \nu) + B \mathcal{Y}_n(q \nu).$$

$$\frac{d^2 \Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p \psi + B \cos p \psi.$$

$$If \alpha_2 = -q^2 \text{ and } \alpha_3 = p^2,$$

$$\frac{d^2 M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + (q^2 - p^2/\mu^2) M = 0, \quad \{24\} \quad M = A \mathcal{J}_p(q \mu) + B \mathcal{J}_{-p}(q \mu) \\ \text{or*} \quad M = A \mathcal{J}_n(q \mu) + B \mathcal{Y}_n(q \mu).$$

$$\frac{d^2 N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} - (q^2 + p^2/\nu^2) N = 0, \quad \{24\} \quad N = A \mathcal{J}_p(i q \nu) + B \mathcal{J}_{-p}(i q \nu) \\ \text{or*} \quad N = A \mathcal{J}_n(i q \nu) + B \mathcal{Y}_n(i q \nu).$$

$$\frac{d^2 \Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p \psi + B \cos p \psi.$$

\* If  $p = n$ , an integer, use the second solution.

If  $\alpha_2 = 0$  and  $\alpha_3 = p^2$ ,

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \frac{p^2}{\mu^2} M = 0, \quad \{04\} \quad M = A\mu^p + B\mu^{-p}.$$

$$\frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} - \frac{p^2}{\nu^2} N = 0, \quad \{04\} \quad N = A\nu^p + B\nu^{-p}.$$

$$\frac{d^2\Psi}{d\psi^2} + p^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p\psi + B \cos p\psi.$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} = 0, \quad \{01\} \quad M = A + B \ln \mu.$$

$$\frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} = 0, \quad \{01\} \quad N = A + B \ln \nu.$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{01\} \quad \Psi = A + B\psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - \alpha_2 M = 0, \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + \alpha_2 N = 0. \end{cases}$$

If  $\alpha_2 = q^2$ ,

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} - q^2 M = 0, \quad \{14\} \quad M = A\mathcal{J}_0(iq\mu) + B\mathcal{Y}_0(iq\mu).$$

$$\frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + q^2 N = 0, \quad \{14\} \quad N = A\mathcal{J}_0(q\nu) + B\mathcal{Y}_0(q\nu).$$

If  $\alpha_2 = -q^2$ ,

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + q^2 M = 0, \quad \{14\} \quad M = A\mathcal{J}_0(q\mu) + B\mathcal{Y}_0(q\mu).$$

$$\frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} - q^2 N = 0, \quad \{14\} \quad N = A\mathcal{J}_0(iq\nu) + B\mathcal{Y}_0(iq\nu).$$

If  $\alpha_2 = 0$ ,

$$\frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} = 0, \quad \{01\} \quad M = A + B \ln \mu.$$

$$\frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} = 0, \quad \{01\} \quad N = A + B \ln \nu.$$

E 4 Rx. BISPERICAL COORDINATES ( $\eta, \theta, \psi$ ), Fig. 4.03.




Fig. 4.03. Bispherical coordinates  $(\eta, \theta, \psi)$ . The spheres are designated by  $\eta = \text{const}$ , half-planes by  $\psi = \text{const}$ . The surfaces  $\theta = \text{const}$  are spindles if  $\theta < \pi/2$ , a sphere at  $\theta = \pi/2$ , and apple-shaped surfaces if  $\theta > \pi/2$

$$-\infty < \eta < +\infty,$$

$$0 \leq \theta < \pi,$$

$$0 \leq \psi < 2\pi.$$

$$\left\{ \begin{array}{l} x = \frac{a \sin \theta \cos \psi}{\cosh \eta - \cos \theta}, \\ y = \frac{a \sin \theta \sin \psi}{\cosh \eta - \cos \theta}, \\ z = \frac{a \sinh \eta}{\cosh \eta - \cos \theta}. \end{array} \right.$$

Coordinate surfaces

$$\left\{ \begin{array}{l} x^2 + y^2 + (z - a \coth \eta)^2 = \frac{a^2}{\sinh^2 \eta} \\ \quad (\text{spheres, } \eta = \text{const}), \\ x^2 + y^2 + z^2 - 2a(x^2 + y^2)^{\frac{1}{2}} \cot \theta = a^2 \\ \quad (\text{apple-shaped surfaces for } \theta < \pi/2, \text{ spindle-shaped surfaces for } \theta > \pi/2. \theta = \text{const}), \\ \tan \psi = y/x \\ \quad (\text{half-planes, } \psi = \text{const}). \end{array} \right.$$

Stäckel matrix

$$[S] = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1/\sin^2 \theta \\ 0 & 0 & 1 \end{bmatrix},$$

$$S = 1, \quad M_{11} = 1,$$

$$M_{21} = 1, \quad M_{31} = 1/\sin^2 \theta.$$

Metric coefficients

$$g_{11} = g_{22} = \frac{a^2}{(\cosh \eta - \cos \theta)^2},$$

$$g_{33} = \frac{a^2 \sin^2 \theta}{(\cosh \eta - \cos \theta)^2},$$

$$g^{\frac{1}{2}} = \frac{a^3 \sin \theta}{(\cosh \eta - \cos \theta)^3}.$$

$$R = [\cosh \eta - \cos \theta]^{-\frac{1}{2}}, \quad Q = a^2/[\cosh \eta - \cos \theta]^2,$$

$$f_1 = 1, \quad f_2 = \sin \theta, \quad f_3 = a, \quad \alpha_1 = -\frac{1}{4}.$$

Important equations

$$(ds)^2 = \frac{a^2}{(\cosh \eta - \cos \theta)^2} [(d\eta)^2 + (d\theta)^2 + \sin^2 \theta (d\psi)^2].$$

$$\text{grad } \varphi = \frac{1}{a} (\cosh \eta - \cos \theta) \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} + \frac{\mathbf{a}_\psi}{\sin \theta} \frac{\partial \varphi}{\partial \psi} \right].$$

$$\operatorname{div} \mathbf{E} = \frac{(\cosh \eta - \cos \theta)^3}{a \sin \theta} \left[ \sin \theta \frac{\partial}{\partial \eta} \left( \frac{E_\eta}{(\cosh \eta - \cos \theta)^2} \right) + \frac{\partial}{\partial \theta} \left( \frac{\sin \theta E_\eta}{(\cosh \eta - \cos \theta)^2} \right) \right] + \frac{(\cosh \eta - \cos \theta)}{a \sin \theta} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{(\cosh \eta - \cos \theta)^2}{a \sin \theta} \begin{vmatrix} \mathbf{a}_\eta & \mathbf{a}_\theta & \mathbf{a}_\psi \sin \theta \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ \frac{E_\eta}{\cosh \eta - \cos \theta} & \frac{E_\theta}{\cosh \eta - \cos \theta} & \frac{E_\psi \sin \theta}{\cosh \eta - \cos \theta} \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{(\cosh \eta - \cos \theta)^3}{a^2 \sin \theta} \left\{ \sin \theta \frac{\partial}{\partial \eta} \left( \frac{1}{\cosh \eta - \cos \theta} \frac{\partial \varphi}{\partial \eta} \right) + \frac{\partial}{\partial \theta} \left( \frac{\sin \theta}{\cosh \eta - \cos \theta} \frac{\partial \varphi}{\partial \theta} \right) \right\} + \frac{(\cosh \eta - \cos \theta)^2}{a^2 \sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = -\frac{1}{4}$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = (\cosh \eta - \cos \theta)^{\frac{1}{2}} H \cdot \Theta \cdot \Psi.$$

General case

$$\begin{cases} \frac{d^2 H}{d\eta^2} - (\frac{1}{4} + \alpha_2) H = 0, \\ \frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2 H}{d\eta^2} - (p + \frac{1}{2})^2 H = 0, \quad \{04\} \quad H = A e^{(p+\frac{1}{2})\eta} + B e^{-(p+\frac{1}{2})\eta}.$$

$$(1 - \xi^2) \frac{d^2 \Theta}{d\xi^2} - 2\xi \frac{d\Theta}{d\xi} + \left[ p(p+1) - \frac{q^2}{1 - \xi^2} \right] \Theta = 0,$$

$\{222\}$  where  $\xi = \cos \theta$ ,

$$\Theta = A \mathcal{P}_p^q(\cos \theta) + B \mathcal{Q}_p^q(\cos \theta).$$

$$\frac{d^2 \Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = 0$ ,

$$\frac{d^2 H}{d\eta^2} - (p + \frac{1}{2})^2 H = 0, \quad \{04\} \quad H = A e^{(p+\frac{1}{2})\eta} + B e^{-(p+\frac{1}{2})\eta}.$$

$$(1 - \xi^2) \frac{d^2 \Theta}{d\xi^2} - 2\xi \frac{d\Theta}{d\xi} + p(p+1) \Theta = 0, \quad \{112\} \quad \Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta).$$

$$\frac{d^2 \Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B \psi.$$

$$If \alpha_2 = \alpha_3 = 0,$$

$$\begin{aligned} \frac{d^2H}{d\eta^2} - \frac{1}{4}H &= 0, & \{04\} \quad H &= A e^{\eta/2} + B e^{-\eta/2}. \\ \frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} &= 0, & \{01\} \quad \Theta &= A + B \ln \cot(\theta/2). \\ \frac{d^2\Psi}{d\psi^2} &= 0, & \{01\} \quad \Psi &= A + B \psi. \end{aligned}$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2H}{d\eta^2} - (\frac{1}{4} + \alpha_2)H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} + \alpha_2 \Theta = 0. \end{cases}$$

$$If \alpha_2 = p(p+1),$$

$$\begin{aligned} \frac{d^2H}{d\eta^2} - (p + \frac{1}{2})^2 H &= 0, & \{04\} \quad H &= A e^{(p+\frac{1}{2})\eta} + B e^{-(p+\frac{1}{2})\eta}. \\ (1 - \xi^2) \frac{d^2\Theta}{d\xi^2} - 2\xi \frac{d\Theta}{d\xi} + p(p+1)\Theta &= 0, & \{112\} \quad \Theta &= A \mathcal{P}_p(\cos\theta) + B \mathcal{Q}_p(\cos\theta). \end{aligned}$$

$$If \alpha_2 = 0,$$

$$\begin{aligned} \frac{d^2H}{d\eta^2} - \frac{1}{4}H &= 0, & \{04\} \quad H &= A e^{\eta/2} + B e^{-\eta/2}. \\ \frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} &= 0, & \{01\} \quad \Theta &= A + B \ln \cot(\theta/2). \end{aligned}$$

E 4 Ry. TOROIDAL COORDINATES  $(\eta, \theta, \psi)$ , Fig. 4.04.

$$0 \leq \eta < +\infty, \quad -\pi < \theta \leq +\pi, \quad 0 \leq \psi < 2\pi.$$

$$\begin{cases} x = \frac{a \sinh \eta \cos \psi}{\cosh \eta - \cos \theta}, \\ y = \frac{a \sinh \eta \sin \psi}{\cosh \eta - \cos \theta}, \\ z = \frac{a \sin \theta}{\cosh \eta - \cos \theta}. \end{cases}$$

Coordinate surfaces

$$\begin{cases} x^2 + y^2 + z^2 + a^2 = 2a(x^2 + y^2)^{\frac{1}{2}} \coth \eta & \text{(toroids, } \eta = \text{const}), \\ x^2 + y^2 + (z - a \cot \theta)^2 = \frac{a^2}{\sin^2 \theta} & \text{(spherical bowls, } \theta = \text{const}), \\ \tan \psi = y/x & \text{(half-planes, } \psi = \text{const}). \end{cases}$$

Stäckel matrix,

$$[S] = \begin{bmatrix} 1 & -1 & -1/\sinh^2 \eta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = 1, \quad M_{11} = 1, \quad M_{21} = 1, \quad M_{31} = 1/\sinh^2 \eta.$$

## Metric coefficients

$$g_{11} = g_{22} = \frac{a^2}{(\cosh \eta - \cos \theta)^2},$$

$$g_{33} = \frac{a^2 \sinh^2 \eta}{(\cosh \eta - \cos \theta)^2},$$

$$g_{\psi}^1 = \frac{a^3 \sinh \eta}{(\cosh \eta - \cos \theta)^3}.$$

$$R = [\cosh \eta - \cos \theta]^{-\frac{1}{2}}, \quad Q = a^2/(\cosh \eta - \cos \theta)^2,$$

$$f_1 = \sinh \eta, \quad f_2 = 1, \quad f_3 = a, \quad \alpha_1 = +\frac{1}{4}.$$




Fig. 4.04. Toroidal coordinates  $(\eta, \theta, \psi)$ . Coordinate surfaces are toroids ( $\eta = \text{const}$ ), spherical bowls ( $\theta = \text{const}$ ), and half-planes ( $\psi = \text{const}$ )

Important equations,

$$(ds)^2 = \frac{a^2}{(\cosh \eta - \cos \theta)^2} [(d\eta)^2 + (d\theta)^2 + \sinh^2 \eta (d\psi)^2].$$

$$\text{grad } \varphi = \frac{1}{a} (\cosh \eta - \cos \theta) \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} + \frac{\mathbf{a}_\psi}{\sinh \eta} \frac{\partial \varphi}{\partial \psi} \right].$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{(\cosh \eta - \cos \theta)^3}{a \sinh \eta} \left[ \frac{\partial}{\partial \eta} \left( \frac{\sinh \eta}{(\cosh \eta - \cos \theta)^2} E_\eta \right) \right. \\ &\quad \left. + \sinh \eta \frac{\partial}{\partial \theta} \left( \frac{E_\theta}{(\cosh \eta - \cos \theta)^2} \right) \right] + \frac{\cosh \eta - \cos \theta}{a \sinh \eta} \frac{\partial E_\psi}{\partial \psi}. \end{aligned}$$

$$\operatorname{curl} \mathbf{E} = \frac{(\cosh \eta - \cos \theta)^2}{a \sinh \eta} \begin{vmatrix} \mathbf{a}_\eta & \mathbf{a}_\theta & \mathbf{a}_\psi \sinh \eta \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_\eta & E_\theta & \frac{E_\psi \sinh \eta}{\cosh \eta - \cos \theta} \end{vmatrix}.$$

$$\begin{aligned} \nabla^2 \varphi &= \frac{(\cosh \eta - \cos \theta)^3}{a^2 \sinh \eta} \left\{ \frac{\partial}{\partial \eta} \left( \frac{\sinh \eta}{\cosh \eta - \cos \theta} \frac{\partial \varphi}{\partial \eta} \right) \right. \\ &\quad \left. + \sinh \eta \frac{\partial}{\partial \theta} \left( \frac{1}{\cosh \eta - \cos \theta} \frac{\partial \varphi}{\partial \theta} \right) \right\} + \frac{(\cosh \eta - \cos \theta)^2}{a^2 \sinh^2 \eta} \frac{\partial^2 \varphi}{\partial \psi^2}. \end{aligned}$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = \frac{1}{4}$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = (\cosh \eta - \cos \theta)^{\frac{1}{2}} H \cdot \Theta \cdot \Psi.$$

**General case**

$$\begin{cases} \frac{d^2 H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + \left( \frac{1}{4} - \alpha_2 - \frac{\alpha_3}{\sinh^2 \eta} \right) H = 0, \\ \frac{d^2 \Theta}{d\theta^2} + \alpha_2 \Theta = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$(\xi^2 - 1) \frac{d^2 H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - \left[ \frac{q^2}{\xi^2 - 1} + (p^2 - \frac{1}{4}) \right] H = 0,$$

{222} where  $\xi = \cosh \eta$ ,

$$H = A \mathcal{P}_{p-\frac{1}{2}}^q(\cosh \eta) + B \mathcal{Q}_{p-\frac{1}{2}}^q(\cosh \eta).$$

$$\frac{d^2 \Theta}{d\theta^2} + p^2 \Theta = 0, \quad \{04\} \quad \Theta = A \sin p \theta + B \cos p \theta.$$

$$\frac{d^2 \Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin p \psi + B \cos p \psi.$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = 0$ ,

$$(\xi^2 - 1) \frac{d^2 H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - (p^2 - \frac{1}{4}) H = 0,$$

{112}  $H = A \mathcal{P}_{p-\frac{1}{2}}(\cosh \eta) + B \mathcal{Q}_{p-\frac{1}{2}}(\cosh \eta)$ .

$$\frac{d^2 \Theta}{d\theta^2} + p^2 \Theta = 0, \quad \{04\} \quad \Theta = A \sin p \theta + B \cos p \theta.$$

$$\frac{d^2 \Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B \psi.$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$(\xi^2 - 1) \frac{d^2 H}{d\xi^2} + 2\xi \frac{dH}{d\xi} + \frac{1}{4} H = 0, \quad \{112\} \quad H = A \mathcal{P}_{-\frac{1}{2}}(\cosh \eta) + B \mathcal{Q}_{-\frac{1}{2}}(\cosh \eta).$$

$$\frac{d^2 \Theta}{d\theta^2} = 0, \quad \{01\} \quad \Theta = A + B \theta.$$

$$\frac{d^2 \Psi}{d\psi^2} = 0, \quad \{01\} \quad \Psi = A + B \psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2 H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + (\frac{1}{4} - \alpha_2) H = 0, \\ \frac{d^2 \Theta}{d\theta^2} + \alpha_2 \Theta = 0. \end{cases}$$

If  $\alpha_2 = p^2$ ,

$$(\xi^2 - 1) \frac{d^2 H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - (p^2 - \frac{1}{4}) H = 0, \quad \{112\} \quad H = A \mathcal{P}_{p-\frac{1}{2}}(\cosh \eta) + B \mathcal{Q}_{p-\frac{1}{2}}(\cosh \eta).$$

$$\frac{d^2 \Theta}{d\theta^2} + p^2 \Theta = 0, \quad \{04\} \quad \Theta = A \sin p \theta + B \cos p \theta.$$

If  $\alpha_2 = 0$ ,

$$\frac{d^2 H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + \frac{1}{4} H = 0, \quad \{112\} \quad H = A \mathcal{P}_{-\frac{1}{2}}(\cosh \eta) + B \mathcal{Q}_{-\frac{1}{2}}(\cosh \eta).$$

$$\frac{d^2 \Theta}{d\theta^2} = 0, \quad \{01\} \quad \Theta = A + B \theta.$$

## H 2 Rx. INVERSE PROLATE SPHEROIDAL COORDINATES ( $\eta, \theta, \psi$ ), Fig. 4.05.

$$0 \leq \eta < +\infty, \quad 0 \leq \theta \leq \pi, \quad 0 \leq \psi < 2\pi.$$

$$\begin{cases} x = \frac{a \sinh \eta \sin \theta \cos \psi}{\cosh^2 \eta - \sin^2 \theta}, \\ y = \frac{a \sinh \eta \sin \theta \sin \psi}{\cosh^2 \eta - \sin^2 \theta}, \\ z = \frac{a \cosh \eta \cos \theta}{\cosh^2 \eta - \sin^2 \theta}. \end{cases}$$

Coordinate surfaces

$$\begin{cases} x^2 + y^2 + z^2 = a \left[ \frac{x^2 + y^2}{\sinh^2 \eta} + \frac{z^2}{\cosh^2 \eta} \right]^{\frac{1}{2}} & \text{(rotation cyclides, } \eta = \text{const}), \\ x^2 + y^2 + z^2 = a \left[ -\frac{x^2 + y^2}{\sin^2 \theta} + \frac{z^2}{\cos^2 \theta} \right]^{\frac{1}{2}} & \text{(rotation cyclides, } \theta = \text{const}), \\ \tan \psi = y/x & \text{(half-planes, } \psi = \text{const}). \end{cases}$$

## Stäckel matrix

$$[S] = \begin{bmatrix} a^2 \sinh^2 \eta & -1 & -1/\sinh^2 \eta \\ a^2 \sin^2 \theta & 1 & -1/\sin^2 \theta \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = a^2(\sinh^2 \eta + \sin^2 \theta), \quad M_{11} = 1, \quad M_{21} = 1, \quad M_{31} = \frac{\sinh^2 \eta + \sin^2 \theta}{\sinh^2 \eta \sin^2 \theta}.$$




Fig. 4.05. Inverse prolate spheroidal coordinates  $(\eta, \theta, \psi)$ . The surfaces  $\eta = \text{const}$  are inversions of prolate spheroids; the surfaces  $\theta = \text{const}$  are inversions of hyperboloids of two sheets

## Metric coefficients

$$g_{11} = g_{22} = \frac{a^2(\sinh^2 \eta + \sin^2 \theta)}{(\cosh^2 \eta - \sin^2 \theta)^2},$$

$$g_{33} = \frac{a^2 \sinh^2 \eta \sin^2 \theta}{(\cosh^2 \eta - \sin^2 \theta)^2},$$

$$g^1 = \frac{a^3 \sinh \eta \sin \theta (\sinh^2 \eta + \sin^2 \theta)}{(\cosh^2 \eta - \sin^2 \theta)^3},$$

$$R = [\cosh^2 \eta - \sin^2 \theta]^{-\frac{1}{2}}, \quad Q = [\cosh^2 \eta - \sin^2 \theta]^{-2},$$

$$f_1 = \sinh \eta, \quad f_2 = \sin \theta, \quad f_3 = a, \quad \alpha_1 = 0.$$

### Important equations

$$(ds)^2 = \frac{a^2}{\Omega^2} [\Pi[(d\eta)^2 + (d\theta)^2] + \sinh^2 \eta \sin^2 \theta (d\psi)^2],$$

where

$$\Pi = \sinh^2 \eta + \sin^2 \theta,$$

$$\Omega = \cosh^2 \eta - \sin^2 \theta.$$

$$\text{grad } \varphi = \frac{\Omega}{a} \left[ \Pi^{-\frac{1}{2}} \left( \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} \right) + \frac{\mathbf{a}_\psi}{\sinh \eta \sin \theta} \frac{\partial \varphi}{\partial \psi} \right].$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{\Omega^3}{a \Pi \sinh \eta \sin \theta} \left[ \sin \theta \frac{\partial}{\partial \eta} \left( \frac{\Pi^{\frac{1}{2}} \sinh \eta}{\Omega^2} E_\eta \right) + \sinh \eta \frac{\partial}{\partial \theta} \left( \frac{\Pi^{\frac{1}{2}} \sin \theta}{\Omega^2} E_\theta \right) \right] \\ &\quad + \frac{\Omega}{a \sinh \eta \sin \theta} \frac{\partial E_\psi}{\partial \psi}. \end{aligned}$$

$$\text{curl } \mathbf{E} = \frac{\Omega^2}{a \Pi^{\frac{1}{2}} \sinh \eta \sin \theta} \begin{vmatrix} \mathbf{a}_\eta & \mathbf{a}_\theta & \mathbf{a}_\psi \Pi^{-\frac{1}{2}} \sinh \eta \sin \theta \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_\eta \frac{\Pi^{\frac{1}{2}}}{\Omega} & E_\theta \frac{\Pi^{\frac{1}{2}}}{\Omega} & E_\psi \frac{\sinh \eta \sin \theta}{\Omega} \end{vmatrix}.$$

$$\begin{aligned} \nabla^2 \varphi &= \frac{\Omega^3}{a^2 \Pi \sinh \eta \sin \theta} \left\{ \sin \theta \frac{\partial}{\partial \eta} \left( \frac{\sinh \eta}{\Omega} \frac{\partial \varphi}{\partial \eta} \right) + \sinh \eta \frac{\partial}{\partial \theta} \left( \frac{\sin \theta}{\Omega} \frac{\partial \varphi}{\partial \theta} \right) \right\} \\ &\quad + \frac{\Omega^2}{a^2 \sinh^2 \eta \sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2}. \end{aligned}$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = (\cosh^2 \eta - \sin^2 \theta)^{\frac{1}{2}} H \cdot \Theta \cdot \Psi.$$

**General case**

$$\begin{cases} \frac{d^2 H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} - \left( \alpha_2 + \frac{\alpha_3}{\sinh^2 \eta} \right) H = 0, \\ \frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$(\xi^2 - 1) \frac{d^2 H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - \left[ p(p+1) + \frac{q^2}{\xi^2 - 1} \right] H = 0,$$

{222} where  $\xi = \cosh \eta$ ,

$$H = A \mathcal{P}_p^q(\cosh \eta) + B \mathcal{Q}_p^q(\cosh \eta).$$

$$(1 - \zeta^2) \frac{d^2\Theta}{d\zeta^2} - 2\zeta \frac{d\Theta}{d\zeta} + \left[ p(p+1) - \frac{q^2}{\zeta^2 - 1} \right] \Theta = 0,$$

$\{222\}$  where  $\zeta = \cos \theta$ ,

$$\Theta = A \mathcal{P}_p^q(\cos \theta) + B \mathcal{Q}_p^q(\cos \theta).$$

$$\frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0, \quad \{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

*If*  $\alpha_2 = p(p+1)$  *and*  $\alpha_3 = 0$ ,

$$(\xi^2 - 1) \frac{d^2H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - p(p+1) H = 0,$$

$\{112\}$   $H = A \mathcal{P}_p(\cosh \eta) + B \mathcal{Q}_p(\cosh \eta).$

$$(1 - \zeta^2) \frac{d^2\Theta}{d\zeta^2} - 2\zeta \frac{d\Theta}{d\zeta} + p(p+1) \Theta = 0,$$

$\{112\}$   $\Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta).$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{01\} \quad \Psi = A + B\psi.$$

*If*  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} = 0, \quad \{01\} \quad H = A + B \ln \coth(\eta/2).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} = 0, \quad \{01\} \quad \Theta = A + B \ln \cot(\theta/2).$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{01\} \quad \Psi = A + B\psi.$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} \frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} - \alpha_2 H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \alpha_2 \Theta = 0. \end{cases}$$

*If*  $\alpha_2 = p(p+1)$ ,

$$(\xi^2 - 1) \frac{d^2H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - p(p+1) H = 0,$$

$\{112\}$  where  $\xi = \cosh \eta$ ,

$$H = A \mathcal{P}_p(\cosh \eta) + B \mathcal{Q}_p(\cosh \eta).$$

$$(1 - \zeta^2) \frac{d^2\Theta}{d\zeta^2} - 2\zeta \frac{d\Theta}{d\zeta} + p(p+1) \Theta = 0,$$

$\{112\}$  where  $\zeta = \cos \theta$ ,

$$\Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta).$$

*If*  $\alpha_2 = 0$ ,

$$\frac{d^2H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} = 0, \quad \{01\} \quad H = A + B \ln \coth(\eta/2).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} = 0, \quad \{01\} \quad \Theta = A + B \ln \cot(\theta/2).$$

H 2Ry. INVERSE OBLATE SPHEROIDAL COORDINATES  $(\eta, \theta, \psi)$ , Fig. 4.06.

$$0 \leq \eta < \infty,$$

$$0 \leq \theta \leq \pi,$$

$$0 \leq \psi < 2\pi.$$

$$\left\{ \begin{array}{l} x = \frac{a \cosh \eta \sin \theta \cos \psi}{\cosh^2 \eta - \cos^2 \theta}, \\ y = \frac{a \cosh \eta \sin \theta \sin \psi}{\cosh^2 \eta - \cos^2 \theta}, \\ z = \frac{a \sinh \eta \cos \theta}{\cosh^2 \eta - \cos^2 \theta}. \end{array} \right.$$

Coordinate surfaces

$$\left\{ \begin{array}{l} x^2 + y^2 + z^2 \\ = a \left[ \frac{x^2 + y^2}{\cosh^2 \eta} + \frac{z^2}{\sinh^2 \eta} \right]^{\frac{1}{2}} \\ (\text{rotation cyclides, } \eta = \text{const}), \end{array} \right.$$

$$\left\{ \begin{array}{l} x^2 + y^2 + z^2 \\ = a \left[ \frac{x^2 + y^2}{\sin^2 \theta} - \frac{z^2}{\cos^2 \theta} \right]^{\frac{1}{2}} \\ (\text{rotation cyclides, } \theta = \text{const}), \end{array} \right.$$

$$\tan \psi = y/x$$

(half-planes,  $\psi = \text{const}$ ).




Fig. 4.06. Inverse oblate spheroidal coordinates  $(\eta, \theta, \psi)$ . The surfaces  $\eta = \text{const}$  are inversions of oblate spheroids; the surfaces  $\theta = \text{const}$  are inversions of hyperboloids of one sheet

**Stäckel matrix**

$$[S] = \begin{bmatrix} a^2 \cosh^2 \eta & -1 & 1/\cosh^2 \eta \\ -a^2 \sin^2 \theta & 1 & -1/\sin^2 \theta \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = a^2(\cosh^2 \eta - \sin^2 \theta), \quad M_{11} = 1, \quad M_{21} = 1, \quad M_{31} = \frac{\cosh^2 \eta - \sin^2 \theta}{\cosh^2 \eta \sin^2 \theta}.$$

**Metric coefficients**

$$g_{11} = g_{22} = \frac{a^2(\cosh^2 \eta - \sin^2 \theta)}{(\cosh^2 \eta - \cos^2 \theta)^2},$$

$$g_{33} = \frac{a^2 \cosh^2 \eta \sin^2 \theta}{(\cosh^2 \eta - \cos^2 \theta)^2},$$

$$g^{\frac{1}{2}} = \frac{a^3 \cosh \eta \sin \theta (\cosh^2 \eta - \sin^2 \theta)}{(\cosh^2 \eta - \cos^2 \theta)^3}.$$

$$R = [\cosh^2 \eta - \cos^2 \theta]^{-\frac{1}{2}}, \quad Q = 1/(\cosh^2 \eta - \cos^2 \theta)^2,$$

$$f_1 = \cosh \eta, \quad f_2 = \sin \theta, \quad f_3 = a, \quad \alpha_1 = 0.$$

### Important equations

$$(ds)^2 = \frac{a^2}{\Sigma^2} \{ \Omega [(d\eta)^2 + (d\theta)^2] + \cosh^2 \eta \sin^2 \theta (d\psi)^2 \},$$

where

$$\Omega = \cosh^2 \eta - \sin^2 \theta,$$

$$\Sigma = \cosh^2 \eta - \cos^2 \theta.$$

$$\text{grad } \varphi = \frac{\Sigma}{a} \left\{ \Omega^{-\frac{1}{2}} \left[ \mathbf{a}_\eta \frac{\partial \varphi}{\partial \eta} + \mathbf{a}_\theta \frac{\partial \varphi}{\partial \theta} \right] + \frac{\mathbf{a}_\psi}{\cosh \eta \sin \theta} \frac{\partial \varphi}{\partial \psi} \right\}.$$

$$\begin{aligned} \text{div } \mathbf{E} &= \frac{\Sigma^3}{a \Omega \cosh \eta \sin \theta} \left[ \sin \theta \frac{\partial}{\partial \eta} \left( \frac{\Omega^{\frac{1}{2}} \cosh \eta}{\Sigma^2} E_\eta \right) + \cosh \eta \frac{\partial}{\partial \theta} \left( \frac{\Omega^{\frac{1}{2}} \sin \theta}{\Sigma^2} E_\theta \right) \right] \\ &\quad + \frac{\Sigma}{a \cosh \eta \sin \theta} E_\psi. \end{aligned}$$

$$\text{curl } \mathbf{E} = \frac{\Sigma^2}{a \Omega^{\frac{1}{2}} \cosh \eta \sin \theta} \begin{vmatrix} \mathbf{a}_\eta & \mathbf{a}_\theta & \mathbf{a}_\psi \Omega^{-\frac{1}{2}} \cosh \eta \sin \theta \\ \frac{\partial}{\partial \eta} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \psi} \\ E_\eta \frac{\Omega^{\frac{1}{2}}}{\Sigma} & E_\theta \frac{\Omega^{\frac{1}{2}}}{\Sigma} & E_\psi \frac{\cosh \eta \sin \theta}{\Sigma} \end{vmatrix}.$$

$$\begin{aligned} \nabla^2 \varphi &= \frac{\Sigma^3}{a^2 \Omega \cosh \eta \sin \theta} \left\{ \sin \theta \frac{\partial}{\partial \eta} \left( \frac{\cosh \eta}{\Sigma} \frac{\partial \varphi}{\partial \eta} \right) + \cosh \eta \frac{\partial}{\partial \theta} \left( \frac{\sin \theta}{\Sigma} \frac{\partial \varphi}{\partial \theta} \right) \right\} \\ &\quad + \frac{\Sigma^2}{a^2 \cosh^2 \eta \sin^2 \theta} \frac{\partial^2 \varphi}{\partial \psi^2}. \end{aligned}$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = H(\eta)$ ,  $U^2 = \Theta(\theta)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = (\cosh^2 \eta - \cos^2 \theta)^{\frac{1}{2}} H \cdot \Theta \cdot \Psi.$$

**General case**

$$\begin{cases} \frac{d^2 H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} + \left( -\alpha_2 + \frac{\alpha_3}{\cosh^2 \eta} \right) H = 0, \\ \frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left( \alpha_2 - \frac{\alpha_3}{\sin^2 \theta} \right) \Theta = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

If  $\alpha_2 = p(p+1)$  and  $\alpha_3 = q^2$ ,

$$(\xi^2 - 1) \frac{d^2 H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - \left[ p(p+1) - \frac{q^2}{\xi^2 - 1} \right] H = 0,$$

{222} where  $\xi = i \sinh \eta$ ,

$$H = A \mathcal{P}_p^q(i \sinh \eta) + B \mathcal{Q}_p^q(i \sinh \eta).$$

$$(1 - \zeta^2) \frac{d^2\Theta}{d\zeta^2} - 2\zeta \frac{d\Theta}{d\zeta} + \left[ p(p+1) - \frac{q^2}{\zeta^2 - 1} \right] \Theta = 0,$$

$\{222\}$  where  $\zeta = \cos \theta$ ,

$$\Theta = A \mathcal{P}_p^q(\cos \theta) + B \mathcal{Q}_p^q(\cos \theta).$$

$$\frac{d^2\Psi}{d\psi^2} + q^2 \Psi = 0,$$

$$\{04\} \quad \Psi = A \sin q\psi + B \cos q\psi.$$

*If*  $\alpha_2 = p(p+1)$  and  $\alpha_3 = 0$ ,

$$(\xi^2 - 1) \frac{d^2H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - p(p+1) H = 0,$$

$$\{112\} \quad H = A \mathcal{P}_p(i \sinh \eta) + B \mathcal{Q}_p(i \sinh \eta).$$

$$(1 - \zeta^2) \frac{d^2\Theta}{d\zeta^2} - 2\zeta \frac{d\Theta}{d\zeta} + p(p+1) \Theta = 0,$$

$$\{112\} \quad \Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta).$$

$$\frac{d^2\Psi}{d\psi^2} = 0,$$

$$\{04\} \quad \Psi = A + B\psi.$$

*If*  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} = 0, \quad \{04\} \quad H = A + B \cot^{-1}(\sinh \eta).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} = 0, \quad \{04\} \quad \Theta = A + B \ln \cot(\theta/2).$$

$$\frac{d^2\Psi}{d\psi^2} = 0, \quad \{04\} \quad \Psi = A + B\psi.$$

**For  $\varphi$  independent of  $\Psi$ ,**

$$\begin{cases} \frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} - \alpha_2 H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \alpha_2 \Theta = 0. \end{cases}$$

*If*  $\alpha_2 = p(p+1)$ ,

$$(\xi^2 - 1) \frac{d^2H}{d\xi^2} + 2\xi \frac{dH}{d\xi} - p(p+1) H = 0,$$

$$\{112\} \quad H = A \mathcal{P}_p(i \sinh \eta) + B \mathcal{Q}_p(i \sinh \eta).$$

$$(1 - \zeta^2) \frac{d^2\Theta}{d\zeta^2} - 2\zeta \frac{d\Theta}{d\zeta} + p(p+1) \Theta = 0,$$

$$\{112\} \quad \Theta = A \mathcal{P}_p(\cos \theta) + B \mathcal{Q}_p(\cos \theta).$$

*If*  $\alpha_2 = 0$ ,

$$\frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} = 0, \quad \{04\} \quad H = A + B \cot^{-1}(\sinh \eta).$$

$$\frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} = 0, \quad \{04\} \quad \Theta = A + B \ln \cot(\theta/2).$$

6-SPHERE COORDINATES  $(u, v, w)$ , Fig. 4.07.




Fig. 4.07. 6-sphere coordinates  $(u, v, w)$ . The inversion of rectangular coordinates. All coordinate surfaces are spheres

$$\begin{aligned} -\infty &< u < +\infty, \\ -\infty &< v < +\infty, \\ -\infty &< w < +\infty. \end{aligned}$$

$$\left\{ \begin{array}{l} x = \frac{u}{u^2 + v^2 + w^2}, \\ y = \frac{v}{u^2 + v^2 + w^2}, \\ z = \frac{w}{u^2 + v^2 + w^2}. \end{array} \right.$$

Coordinate surfaces

$$\begin{aligned} \left( x - \frac{1}{2u} \right)^2 + y^2 + z^2 &= \frac{1}{4u^2} \\ (\text{spheres tangent to } yz\text{-plane at origin, } u = \text{const}), \\ x^2 + \left( y - \frac{1}{2v} \right)^2 + z^2 &= \frac{1}{4v^2} \\ (\text{spheres tangent to } xz\text{-plane at origin, } v = \text{const}), \\ x^2 + y^2 + \left( z - \frac{1}{2w} \right)^2 &= \frac{1}{4w^2} \\ (\text{spheres tangent to } xy\text{-plane at origin, } w = \text{const}). \end{aligned}$$

**Stäckel matrix**

$$[S] = \begin{bmatrix} 0 & -1 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

$$S = 1, \quad M_{11} = M_{21} = M_{31} = 1.$$

**Metric coefficients,**

$$g_{11} = g_{22} = g_{33} = 1/[u^2 + v^2 + w^2]^2, \quad g^{\frac{1}{2}} = 1/[u^2 + v^2 + w^2]^3.$$

$$R = [u^2 + v^2 + w^2]^{-\frac{1}{2}}, \quad Q = 1/[u^2 + v^2 + w^2]^2,$$

$$f_1 = f_2 = f_3 = 1, \quad \alpha_1 = 0.$$

**Important equations,**

$$(ds)^2 = \frac{(du)^2 + (dv)^2 + (dw)^2}{[u^2 + v^2 + w^2]^2}.$$

$$\text{grad } \varphi = (u^2 + v^2 + w^2) \left[ \mathbf{a}_u \frac{\partial \varphi}{\partial u} + \mathbf{a}_v \frac{\partial \varphi}{\partial v} + \mathbf{a}_w \frac{\partial \varphi}{\partial w} \right].$$

$$\begin{aligned} \text{div } \mathbf{E} &= (u^2 + v^2 + w^2)^3 \left[ \frac{\partial}{\partial u} \left( \frac{E_u}{(u^2 + v^2 + w^2)^2} \right) \right. \\ &\quad \left. + \frac{\partial}{\partial v} \left( \frac{E_v}{(u^2 + v^2 + w^2)^2} \right) + \frac{\partial}{\partial w} \left( \frac{E_w}{(u^2 + v^2 + w^2)^2} \right) \right]. \end{aligned}$$

$$\text{curl } \mathbf{E} = (u^2 + v^2 + w^2)^2 \begin{vmatrix} \mathbf{a}_u & \mathbf{a}_v & \mathbf{a}_w \\ \frac{\partial}{\partial u} & \frac{\partial}{\partial v} & \frac{\partial}{\partial w} \\ \frac{E_u}{u^2 + v^2 + w^2} & \frac{E_v}{u^2 + v^2 + w^2} & \frac{E_w}{u^2 + v^2 + w^2} \end{vmatrix}.$$

$$\nabla^2 \varphi = (u^2 + v^2 + w^2)^3 \left[ \frac{\partial}{\partial u} \left( \frac{1}{u^2 + v^2 + w^2} \frac{\partial \varphi}{\partial u} \right) + \frac{\partial}{\partial v} \left( \frac{1}{u^2 + v^2 + w^2} \frac{\partial \varphi}{\partial v} \right) + \frac{\partial}{\partial w} \left( \frac{1}{u^2 + v^2 + w^2} \frac{\partial \varphi}{\partial w} \right) \right].$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = 0$  and  $U^1 = U(u)$ ,  $U^2 = V(v)$ ,  $U^3 = W(w)$ ,

$$\varphi = (u^2 + v^2 + w^2)^{\frac{1}{2}} U \cdot V \cdot W.$$

**General case**

$$\begin{cases} \frac{d^2 U}{d u^2} - (\alpha_2 + \alpha_3) U = 0, \\ \frac{d^2 V}{d v^2} + \alpha_2 V = 0, \\ \frac{d^2 W}{d w^2} + \alpha_3 W = 0. \end{cases}$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\frac{d^2 X}{d x^2} - (p^2 + q^2) X = 0, \quad \{04\} \quad X = A e^{(p^2 + q^2)^{\frac{1}{2}} x} + B e^{-(p^2 + q^2)^{\frac{1}{2}} x}$$

$$\frac{d^2 Y}{d y^2} + p^2 Y = 0, \quad \{04\} \quad Y = A \sin p y + B \cos p y.$$

$$\frac{d^2 Z}{d z^2} + q^2 Z = 0, \quad \{04\} \quad Z = A \sin q z + B \cos q z.$$

If  $\alpha_2 = -p^2$  and  $\alpha_3 = -q^2$ ,

$$\frac{d^2 X}{d x^2} + (p^2 + q^2) X = 0, \quad \{04\} \quad X = A \sin (p^2 + q^2)^{\frac{1}{2}} x + B \cos (p^2 + q^2)^{\frac{1}{2}} x.$$

$$\frac{d^2 Y}{d y^2} - p^2 Y = 0, \quad \{04\} \quad Y = A e^{p y} + B e^{-p y}.$$

$$\frac{d^2 Z}{d z^2} - q^2 Z = 0, \quad \{04\} \quad Z = A e^{q z} + B e^{-q z}.$$

If  $\alpha_2 = p^2$  and  $\alpha_3 = 0$ ,

$$\frac{d^2 X}{d x^2} - p^2 X = 0, \quad \{04\} \quad X = A e^{p x} + B e^{-p x}.$$

$$\frac{d^2 Y}{d y^2} + p^2 Y = 0, \quad \{04\} \quad Y = A \sin p y + B \cos p y.$$

$$\frac{d^2 Z}{d z^2} = 0, \quad \{01\} \quad Z = A + B z.$$

If  $\alpha_2 = \alpha_3 = 0$ ,

$$\frac{d^2 X}{d x^2} = \frac{d^2 Y}{d y^2} = \frac{d^2 Z}{d z^2} = 0, \quad X = A + B x, \quad Y = A + B y, \quad Z = A + B z.$$

J 1 Rx. BI-CYCLIDE COORDINATES  $(\mu, \nu, \psi)$ , Fig. 4.08.




Fig. 4.08. Bi-cyclide coordinates  $(\mu, \nu, \psi)$ . Similar to bispherical coordinates (Fig. 4.03), but with 4th-degree surfaces instead of 2nd-degree for  $\mu = \text{const}$ . Corresponding to the two spheres of bispherical coordinates are two cyclides  $\mu = \pm \text{const}$ . Corresponding to the spindles or apple-shaped surfaces are the cyclides  $\nu = \text{const}$

$$0 \leq \mu \leq K, \quad 0 \leq \nu \leq K', \quad 0 \leq \psi < 2\pi.$$

$$\begin{cases} x = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu \cos \psi, \\ y = \frac{a}{A} \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu \sin \psi, \\ z = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \nu, \end{cases}$$

$$A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu.$$

#### Coordinate surfaces

$$\left\{ \begin{array}{l} (x^2 + y^2 + z^2)^2 + \frac{a^2}{k^4} \frac{[(1 - k^2)^2 - 2(1 - k^2) \operatorname{dn}^2 \mu + (1 + k^2) \operatorname{dn}^4 \mu]}{\operatorname{dn}^2 \mu \operatorname{cn}^2 \mu} (x^2 + y^2) \\ \quad - a^2 \left( \operatorname{sn}^2 \mu + \frac{1}{k^2 \operatorname{sn}^2 \mu} \right) z^2 + \frac{a^4}{k^2} = 0 \quad (\text{bi-cyclides, } \mu = \text{const}), \\ \left[ \frac{\operatorname{cn}^2 \nu}{a^2 \operatorname{sn}^2 \nu} (x^2 + y^2) + \frac{\operatorname{dn}^2 \nu}{a^2} z^2 \right]^2 - \frac{2 \operatorname{cn}^2 \nu}{a^2 \operatorname{sn}^2 \nu} (x^2 + y^2) - \frac{2 \operatorname{dn}^2 \nu}{a^2} z^2 + 1 = 0 \\ \quad (\text{rotation-cyclides, } \nu = \text{const}), \\ \tan \psi = y/x \quad (\text{half planes, } \psi = \text{const}). \end{array} \right.$$

**Stäckel matrix**

$$[S] = \begin{bmatrix} -k^2 \operatorname{sn}^2 \mu & -1 & -\frac{k'^2 \operatorname{sn}^2 \mu}{\operatorname{cn}^2 \mu \operatorname{dn}^2 \mu} \\ \operatorname{dn}^2 \nu & 1 & -\frac{\operatorname{dn}^2 \nu}{\operatorname{sn}^2 \nu \operatorname{cn}^2 \nu} \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = \operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu = \operatorname{dn}^2 \mu - k'^2 \operatorname{sn}^2 \nu,$$

$$M_{11} = M_{21} = 1, \quad M_{31} = \frac{\Omega^2}{\operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu}.$$

**Metric coefficients**

$$g_{11} = g_{22} = \frac{a^2 \Omega^2}{A^2},$$

$$g_{33} = \frac{a^2}{A^2} \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu,$$

$$g^{\frac{1}{2}} = \frac{a^3 \Omega^2}{A^3} \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu,$$

where

$$\Omega^2 = (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu) (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu).$$

$$R = A^{-\frac{1}{2}}, \quad Q = \frac{a^2}{A^2} (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu),$$

$$f_1 = \operatorname{cn} \mu \operatorname{dn} \mu, \quad f_2 = \operatorname{sn} \nu \operatorname{cn} \nu, \quad f_3 = a, \quad \alpha_1 = -2.$$

**Important equations**

$$(ds)^2 = \frac{a^2 \Omega^2}{A^2} [(d\mu)^2 + (d\nu)^2] + \frac{a^2}{A^2} \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu (d\psi)^2.$$

$$\operatorname{grad} \varphi = \frac{A}{a \Omega} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_\psi \frac{A}{a \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu} \frac{\partial \varphi}{\partial \psi}.$$

$$\operatorname{div} \mathbf{E} = \frac{A^3}{a \Omega^2 \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu} \left\{ \operatorname{sn} \nu \operatorname{cn} \nu \frac{\partial}{\partial \mu} \left[ \frac{\Omega \operatorname{cn} \mu \operatorname{dn} \mu}{A^2} E_\mu \right] + \operatorname{cn} \mu \operatorname{dn} \mu \frac{\partial}{\partial \nu} \left[ \frac{\Omega \operatorname{sn} \nu \operatorname{cn} \nu}{A^2} E_\nu \right] \right\} + \frac{A}{a \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{A^2}{a \Omega^2 \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu} \begin{vmatrix} \mathbf{a}_\mu \Omega & \mathbf{a}_\nu \Omega & \mathbf{a}_\psi \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ E_\mu \Omega / A & E_\nu \Omega / A & E_\psi \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu / A \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{A^3}{a^2 \Omega^2 \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu} \left\{ \operatorname{sn} \nu \operatorname{cn} \nu \frac{\partial}{\partial \mu} \left[ \frac{\operatorname{cn} \mu \operatorname{dn} \mu}{A} \frac{\partial \varphi}{\partial \mu} \right] + \operatorname{cn} \mu \operatorname{dn} \mu \frac{\partial}{\partial \nu} \left[ \frac{\operatorname{sn} \nu \operatorname{cn} \nu}{A} \frac{\partial \varphi}{\partial \nu} \right] \right\} + \frac{A^2}{a^2 \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = -2$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = A^{\frac{1}{2}} M N \Psi.$$

**General case**

$$\begin{cases} \frac{d^2M}{d\mu^2} - \frac{\operatorname{sn} \mu (\operatorname{dn}^2 \mu + k^2 \operatorname{cn}^2 \mu)}{\operatorname{cn} \mu \operatorname{dn} \mu} \frac{dM}{d\mu} + \left[ 2k^2 \operatorname{sn}^2 \mu - \alpha_2 - \alpha_3 \frac{k'^4 \operatorname{sn}^2 \mu}{\operatorname{cn}^2 \mu \operatorname{dn}^2 \mu} \right] M = 0, \\ \frac{d^2N}{d\nu^2} + \frac{\operatorname{dn} \nu (\operatorname{cn}^2 \nu - \operatorname{sn}^2 \nu)}{\operatorname{sn} \nu \operatorname{cn} \nu} \frac{dN}{d\nu} + \left[ -2 \operatorname{dn}^2 \nu + \alpha_2 - \alpha_3 \frac{\operatorname{dn}^2 \nu}{\operatorname{sn}^2 \nu \operatorname{cn}^2 \nu} \right] N = 0, \\ \frac{d^2\Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

Substitution of  $\operatorname{sn}^2 \mu = z$  in the first equation and  $\operatorname{dn}^2 \nu = z$  in the second, reduces both to the canonical form

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z - a_1} + \frac{2}{z - a_2} + \frac{2}{z - a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2 + \bar{A}_3 z^3}{(z - a_1)(z - a_2)^2(z - a_3)^2} \right] Z = 0,$$

where, for  $M$  with  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} a_1 &= 0, & a_2 &= b = 1, & a_3 &= c = 1/k^2, \\ \bar{A}_0 &= -p^2 c, & \bar{A}_1 &= p^2 c(1+c) - q^2(1-c)^2 + 2c, \\ \bar{A}_2 &= -[p^2 c + 2(1+c)], & \bar{A}_3 &= 2. \end{aligned}$$

For  $N$  with  $\alpha_2 = p'^2$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} a_1 &= 0, & a_2 &= b = 1, & a_3 &= c = k^2, \\ \bar{A}_0 &= -p'^2 c, & \bar{A}_1 &= p'^2(1+c) - q^2(1-c)^2 + 2c, \\ \bar{A}_2 &= -[p'^2 + 2(1+c)], & \bar{A}_3 &= 2. \end{aligned}$$

Evidently  $p'^2 = p^2 c$ . This is a Bôcher equation with designation {1222}, and the general solution is

$$Z = A \mathcal{U}_p^q(k, z) + B \mathcal{V}_p^q(k, z).$$

**General case,  $\varphi = \varphi(u^1, u^2, u^3)$** 

$$\begin{cases} M = A \mathcal{U}_p^q(k, \operatorname{sn}^2 \mu) + B \mathcal{V}_p^q(k, \operatorname{sn}^2 \mu), \\ N = A \mathcal{U}_{p'}^q(1/k, \operatorname{dn}^2 \nu) + B \mathcal{V}_{p'}^q(1/k, \operatorname{dn}^2 \nu), \\ \Psi = A \sin q \psi + B \cos q \psi, \quad \text{where } p'^2 = p^2/k^2. \end{cases}$$

**For  $\varphi$  independent of  $\Psi$ ,**

$$\begin{cases} M = A \mathcal{U}_p(k, \operatorname{sn}^2 \mu) + B \mathcal{V}_p(k, \operatorname{sn}^2 \mu), \\ N = A \mathcal{U}_{p'}(1/k, \operatorname{dn}^2 \nu) + B \mathcal{V}_{p'}(1/k, \operatorname{dn}^2 \nu). \end{cases}$$

**J 1 Ry. FLAT-RING CYCLIDE COORDINATES  $(\mu, \nu, \psi)$ , Fig. 4.09.**

$$0 \leqq \mu \leqq K, \quad 0 \leqq \nu \leqq K', \quad 0 \leqq \psi < 2\pi.$$

$$\begin{cases} x = \frac{a}{\Lambda} \operatorname{sn} \mu \operatorname{dn} \nu \cos \psi, \\ y = \frac{a}{\Lambda} \operatorname{sn} \mu \operatorname{dn} \nu \sin \psi, \\ z = \frac{a}{\Lambda} \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu, \end{cases}$$

where

$$\Lambda = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu.$$

## Coordinate surfaces

$$\left\{ \begin{array}{l} (x^2 + y^2 + z^2)^2 + \frac{a^2}{k^4} \frac{[(1 - k^2)^2 - 2(1 - k^2) \operatorname{dn}^2 \mu + (1 + k^2) \operatorname{dn}^4 \mu]}{\operatorname{dn}^2 \mu \operatorname{cn}^2 \mu} z^2 \\ \quad - a^2 \left( \operatorname{sn}^2 \mu + \frac{1}{k^2 \operatorname{sn}^2 \mu} \right) (x^2 + y^2) + \frac{a^4}{k^2} = 0 \quad (\text{flat-ring cyclides, } \mu = \text{const}), \\ \left[ \frac{\operatorname{dn}^2 \nu}{a^2} (x^2 + y^2) + \frac{\operatorname{cn}^2 \nu}{a^2 \operatorname{sn}^2 \nu} z^2 \right]^2 - \frac{2 \operatorname{cn}^2 \nu}{a^2 \operatorname{sn}^2 \nu} z^2 - \frac{2 \operatorname{dn}^2 \nu}{a^2} (x^2 + y^2) + 1 = 0 \\ \quad (\text{rotation-cyclides, } \nu = \text{const}), \\ \tan \psi = y/x \quad (\text{half-planes, } \psi = \text{const}). \end{array} \right.$$




Fig. 4.09. Flat-ring cyclide coordinates  $(\mu, \nu, \psi)$ . Similar to toroidal coordinates (Fig. 4.04) but with 4th-degree surfaces instead of 2nd-degree for  $\nu = \text{const}$ . The toroids of circular cross-section are replaced by flattened rings ( $\mu = \text{const}$ ), and the spherical bowls of toroidal coordinates are replaced by rotation cyclides ( $\nu = \text{const}$ )

## Stäckel matrix

$$[S] = \begin{bmatrix} -k^2 \operatorname{sn}^2 \mu & -1 & -\left(k^2 \operatorname{sn}^2 \mu + \frac{1}{\operatorname{sn}^2 \mu}\right) \\ \operatorname{dn}^2 \nu & 1 & \left(\operatorname{dn}^2 \nu + \frac{k^2}{\operatorname{dn}^2 \nu}\right) \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = \operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu = \operatorname{dn}^2 \mu - k'^2 \operatorname{sn}^2 \nu,$$

$$M_{11} = M_{21} = 1, \quad M_{31} = \frac{\Omega^2}{\operatorname{sn}^2 \mu \operatorname{dn}^2 \nu}.$$

**Metric coefficients**

$$g_{11} = g_{22} = a^2 \Omega^2 / A^2,$$

$$g_{33} = \frac{a^2}{A^2} \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu,$$

$$g^1 = \frac{a^3 \Omega^2}{A^3} \operatorname{sn} \mu \operatorname{dn} \nu,$$

where

$$\Omega^2 = (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu) (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu).$$

$$R = A^{-\frac{1}{2}}, \quad Q = \frac{a^2}{A^2} (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu),$$

$$f_1 = \operatorname{sn} \mu, \quad f_2 = \operatorname{dn} \nu, \quad f_3 = a, \quad \alpha_1 = -1.$$

**Important equations**

$$(ds)^2 = \frac{a^2 \Omega^2}{A^2} [(d\mu)^2 + (d\nu)^2] + \frac{a^2}{A^2} \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu (d\psi)^2.$$

$$\operatorname{grad} \varphi = \frac{A}{a \Omega} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \frac{\mathbf{a}_\psi A}{a \operatorname{sn} \mu \operatorname{dn} \nu} \frac{\partial \varphi}{\partial \psi}.$$

$$\operatorname{div} \mathbf{E} = \frac{A^3}{a \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu} \times \left\{ \operatorname{dn} \nu \frac{\partial}{\partial \mu} \left[ \frac{\Omega \operatorname{sn} \mu}{A^2} E_\mu \right] + \operatorname{sn} \mu \frac{\partial}{\partial \nu} \left[ \frac{\Omega \operatorname{dn} \nu}{A^2} E_\nu \right] \right\} + \frac{A}{a \operatorname{sn} \mu \operatorname{dn} \nu} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{A^2}{a \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu} \begin{vmatrix} \mathbf{a}_\mu \Omega & \mathbf{a}_\nu \Omega & \mathbf{a}_\psi \operatorname{sn} \mu \operatorname{dn} \nu \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ E_\mu \Omega / A & E_\nu \Omega / A & E_\psi \operatorname{sn} \mu \operatorname{dn} \nu / A \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{A^3}{a^2 \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu} \times \left\{ \operatorname{dn} \nu \frac{\partial}{\partial \mu} \left[ \frac{\operatorname{sn} \mu}{A} \frac{\partial \varphi}{\partial \mu} \right] + \operatorname{sn} \mu \frac{\partial}{\partial \nu} \left[ \frac{\operatorname{dn} \nu}{A} \frac{\partial \varphi}{\partial \nu} \right] \right\} + \frac{A^2}{a^2 \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

**SEPARATION OF LAPLACE'S EQUATION,  $\nabla^2 \varphi = 0$ .**

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = -1$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = A^{\frac{1}{2}} M N \Psi.$$

**General case**

$$\begin{cases} \frac{d^2 M}{d\mu^2} + \frac{\operatorname{cn} \mu \operatorname{dn} \mu}{\operatorname{sn} \mu} \frac{dM}{d\mu} + \left[ k^2 \operatorname{sn}^2 \mu - \alpha_2 - \alpha_3 \left( k^2 \operatorname{sn}^2 \mu + \frac{1}{\operatorname{sn}^2 \mu} \right) \right] M = 0, \\ \frac{d^2 N}{d\nu^2} - \frac{k'^2 \operatorname{sn} \nu \operatorname{cn} \nu}{\operatorname{dn} \nu} \frac{dN}{d\nu} + \left[ -\operatorname{dn}^2 \nu + \alpha_2 + \alpha_3 \left( \operatorname{dn}^2 \nu + \frac{k^2}{\operatorname{dn}^2 \nu} \right) \right] N = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

Substitution of  $\operatorname{sn}^2 \mu = z$  in the first equation and  $\operatorname{dn}^2 \nu = z$  in the second, reduces both to the canonical form

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{2}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z-a_1)(z-a_2)(z-a_3)^2} \right] Z = 0,$$

where, for  $M$  with  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} a_1 &= b = 1, & a_2 &= c = 1/k^2, & a_3 &= 0, \\ \bar{A}_0 &= -q^2 c, & \bar{A}_1 &= -p^2 c, & \bar{A}_2 &= 1 - q^2. \end{aligned}$$

For  $N$ , with  $\alpha_2 = p'^2$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} a_1 &= b = 1, & a_2 &= c = k^2, & a_3 &= 0, \\ \bar{A}_0 &= -q^2 c, & \bar{A}_1 &= p'^2, & \bar{A}_2 &= 1 - q^2. \end{aligned}$$

These are Bôcher equations with designation {1122}, and their solutions are Wangerin functions:

$$Z = A \mathcal{S}_p^q(k, z) + B \mathcal{T}_p^q(k, z).$$

**General case,**  $\varphi = \varphi(u^1, u^2, u^3)$ ,

$$\begin{cases} M = A \mathcal{S}_p^q(k, \operatorname{sn}^2 \mu) + B \mathcal{T}_p^q(k, \operatorname{sn}^2 \mu), \\ N = A \mathcal{S}_{p'}^q(1/k, \operatorname{dn}^2 \nu) + B \mathcal{T}_{p'}^q(1/k, \operatorname{dn}^2 \nu), \\ \Psi = A \sin q \psi + B \cos q \psi, \quad \text{where } p'^2 = p^2/k^2. \end{cases}$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} M = A \mathcal{S}_p(k, \operatorname{sn}^2 \mu) + B \mathcal{T}_p(k, \operatorname{sn}^2 \mu), \\ N = A \mathcal{S}_{p'}(1/k, \operatorname{dn}^2 \nu) + B \mathcal{T}_{p'}(1/k, \operatorname{dn}^2 \nu). \end{cases}$$

## J 2R. DISK-CYCLIDE COORDINATES ( $\mu, \nu, \psi$ ), Fig. 4.10.

$$0 \leqq \mu \leqq K, \quad 0 \leqq \nu \leqq K', \quad 0 \leqq \psi < 2\pi.$$

$$\begin{cases} x = \frac{a}{A} \operatorname{cn} \mu \operatorname{cn} \nu \cos \psi, \\ y = \frac{a}{A} \operatorname{cn} \mu \operatorname{cn} \nu \sin \psi, \\ z = \frac{a}{A} \operatorname{sn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{dn} \nu, \\ A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu. \end{cases}$$

### Coordinate surfaces

$$\begin{cases} \left( \frac{x^2 + y^2}{a^2 \operatorname{cn}^2 \mu} + \frac{k^2 \operatorname{sn}^2 \mu}{a^2 \operatorname{dn}^2 \mu} z^2 \right)^2 - \frac{2}{a^2 \operatorname{cn}^2 \mu} (x^2 + y^2) - \frac{2k^2 \operatorname{sn}^2 \mu}{a^2 \operatorname{dn}^2 \mu} z^2 + 1 = 0 \\ \quad (\text{rotation-cyclides, } \mu = \text{const}), \\ \left( \frac{\operatorname{cn}^2 \nu}{a^2} (x^2 + y^2) + \frac{k'^2 \operatorname{sn}^2 \nu}{a^2 \operatorname{dn}^2 \nu} z^2 \right)^2 - \frac{2 \operatorname{cn}^2 \nu}{a^2} (x^2 + y^2) - \frac{2k'^2 \operatorname{sn}^2 \nu}{a^2 \operatorname{dn}^2 \nu} z^2 + 1 = 0 \\ \quad (\text{disk-cyclides, } \nu = \text{const}), \\ \tan \psi = y/x \\ \quad (\text{half-planes, } \psi = \text{const}). \end{cases}$$

**Stäckel matrix**

$$[S] = \begin{bmatrix} -k^2 \operatorname{sn}^2 \mu & -1 & \left( k^2 \operatorname{cn}^2 \mu - \frac{k'^2}{\operatorname{cn}^2 \mu} \right) \\ \operatorname{dn}^2 \nu & 1 & \left( k'^2 \operatorname{cn}^2 \nu - \frac{k^2}{\operatorname{cn}^2 \nu} \right) \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = \operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu = \operatorname{dn}^2 \mu - k'^2 \operatorname{sn}^2 \nu,$$

$$M_{11} = M_{21} = 1, \quad M_{31} = \frac{\Gamma^2}{\operatorname{cn}^2 \mu \operatorname{cn}^2 \nu}.$$




Fig. 4.10. Disk-cyclide coordinates  $(\mu, \nu, \psi)$ . Coordinate surfaces are cyclides  $(\mu = \text{const}$  and  $\nu = \text{const})$  and half-planes  $(\psi = \text{const})$

**Metric coefficients**

$$g_{11} = g_{22} = a^2 \Gamma^2 / A^2,$$

$$g_{33} = \frac{a^2}{A^2} \operatorname{cn}^2 \mu \operatorname{cn}^2 \nu,$$

$$g^{\frac{1}{2}} = \frac{a^3 \Gamma^2}{A^3} \operatorname{cn} \mu \operatorname{cn} \nu,$$

where

$$\Gamma^2 = (\operatorname{sn}^2 \nu + \operatorname{sn}^2 \mu \operatorname{cn}^2 \nu) (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu).$$

$$R = A^{-\frac{1}{2}}, \quad Q = \frac{a^2}{A^2} (\operatorname{sn}^2 \nu + \operatorname{sn}^2 \mu \operatorname{cn}^2 \nu),$$

$$f_1 = \operatorname{cn} \mu, \quad f_2 = \operatorname{cn} \nu, \quad f_3 = a, \quad \alpha_1 = -1.$$

**Important equations**

$$(ds)^2 = \frac{a^2 \Gamma^2}{A^2} [(d\mu)^2 + (d\nu)^2] + \frac{a^2}{A^2} \operatorname{cn}^2 \mu \operatorname{cn}^2 \nu (d\psi)^2.$$

$$\operatorname{grad} \varphi = \frac{A}{a \Gamma} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_\psi \frac{A}{a \operatorname{cn} \mu \operatorname{cn} \nu} \frac{\partial \varphi}{\partial \psi}.$$

$$\operatorname{div} \mathbf{E} = \frac{A^3}{a \Gamma^2 \operatorname{cn} \mu \operatorname{cn} \nu} \\ \times \left\{ \operatorname{cn} \nu \frac{\partial}{\partial \mu} \left[ \frac{\Gamma \operatorname{cn} \mu}{A^2} E_\mu \right] + \operatorname{cn} \mu \frac{\partial}{\partial \nu} \left[ \frac{\Gamma \operatorname{cn} \nu}{A^2} E_\nu \right] \right\} + \frac{A}{a \operatorname{cn} \mu \operatorname{cn} \nu} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{A^2}{a \Gamma^2 \operatorname{cn} \mu \operatorname{cn} \nu} \begin{vmatrix} \mathbf{a}_\mu \Gamma & \mathbf{a}_\nu \Gamma & \mathbf{a}_\psi \operatorname{cn} \mu \operatorname{cn} \nu \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ E_\mu \Gamma / A & E_\nu \Gamma / A & E_\psi \operatorname{cn} \mu \operatorname{cn} \nu / A \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{A^3}{a^2 \Gamma^2 \operatorname{cn} \mu \operatorname{cn} \nu} \\ \times \left\{ \operatorname{cn} \nu \frac{\partial}{\partial \mu} \left[ \frac{\operatorname{cn} \mu}{A} \frac{\partial \varphi}{\partial \mu} \right] + \operatorname{cn} \mu \frac{\partial}{\partial \nu} \left[ \frac{\operatorname{cn} \nu}{A} \frac{\partial \varphi}{\partial \nu} \right] \right\} + \frac{A^2}{a^2 \operatorname{cn}^2 \mu \operatorname{cn}^2 \nu} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

**SEPARATION OF LAPLACE'S EQUATION,**  $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{d u^i} \left( f_i \frac{d U^i}{d u^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = -1$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = A^{\frac{1}{2}} M N \Psi.$$

**General case**

$$\begin{cases} \frac{d^2 M}{d \mu^2} - \frac{\operatorname{sn} \mu \operatorname{dn} \mu}{\operatorname{cn} \mu} \frac{d M}{d \mu} + \left[ k^2 \operatorname{sn}^2 \mu - \alpha_2 + \alpha_3 \left( k^2 \operatorname{cn}^2 \mu - \frac{k'^2}{\operatorname{cn}^2 \mu} \right) \right] M = 0, \\ \frac{d^2 N}{d \nu^2} - \frac{\operatorname{sn} \nu \operatorname{dn} \nu}{\operatorname{cn} \nu} \frac{d N}{d \nu} + \left[ -\operatorname{dn}^2 \nu + \alpha_2 + \alpha_3 \left( k'^2 \operatorname{cn}^2 \nu - \frac{k^2}{\operatorname{cn}^2 \nu} \right) \right] N = 0, \\ \frac{d^2 \Psi}{d \psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

Substitution of  $\operatorname{cn}^2 \mu = z$  and  $\operatorname{cn}^2 \nu = z$  in the first two equations reduces them to the canonical form

$$\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z - a_1} + \frac{1}{z - a_2} + \frac{2}{z - a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z - a_1)(z - a_2)(z - a_3)} \right] Z = 0,$$

where, for  $M$  with  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$a_1 = b = 1, \quad a_2 = c = -(k'/k)^2, \quad a_3 = 0,$$

$$\bar{A}_0 = -q^2 c, \quad \bar{A}_1 = p^2(1 - c) - 1, \quad \bar{A}_2 = 1 - q^2.$$

For  $N$ , with  $\alpha_2 = p'^2$  and  $\alpha_3 = q^2$ ,

$$a_1 = b = 1, \quad a_2 = c = -(k/k')^2, \quad a_3 = 0,$$

$$\bar{A}_0 = -q^2 c, \quad \bar{A}_1 = -[p'^2(1 - c) + c], \quad \bar{A}_2 = 1 - q^2.$$

Evidently  $p'^2 = 1 - p^2$ . These are Bôcher equations with designation {1122}, and their solutions are the Wangerin functions,

$$Z = A \mathcal{S}_p^q(k, z) + B \mathcal{T}_p^q(k, z).$$

For  $\varphi = \varphi(u^1, u^2, u^3)$ ,

$$\begin{cases} M = A \mathcal{S}_p^q(i k/k', \operatorname{cn}^2 \mu) + B \mathcal{T}_p^q(i k/k', \operatorname{cn}^2 \mu), \\ N = A \mathcal{S}_{p'}^q(i k'/k, \operatorname{cn}^2 \nu) + B \mathcal{T}_{p'}^q(i k'/k, \operatorname{cn}^2 \nu), \\ \Psi = A \sin q \psi + B \cos q \psi, \quad \text{where } p'^2 = 1 - p^2. \end{cases}$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} M = A \mathcal{S}_p(i k/k', \operatorname{cn}^2 \mu) + B \mathcal{T}_p(i k/k', \operatorname{cn}^2 \mu), \\ N = A \mathcal{S}_{p'}(i k'/k, \operatorname{cn}^2 \nu) + B \mathcal{T}_{p'}(i k'/k, \operatorname{cn}^2 \nu). \end{cases}$$

J 3 R. CAP-CYCLIDE COORDINATES  $(\mu, \nu, \psi)$ , Fig. 4.11.




Fig. 4.11. Cap-cyclide coordinates  $(\mu, \nu, \psi)$ . An inversion of bi-cyclide coordinates (Fig. 4.08). Coordinate surfaces are cap cyclides  $(\nu = \text{const})$ , ring cyclides  $(\mu = \text{const})$ , and half-planes  $(\psi = \text{const})$

$$0 \leq \mu \leq K, \quad 0 \leq \nu \leq K', \quad 0 \leq \psi < 2\pi.$$

$$\begin{cases} x = \frac{A}{a \mathcal{T}} \operatorname{sn} \mu \operatorname{dn} \nu \cos \psi, \\ y = \frac{A}{a \mathcal{T}} \operatorname{sn} \mu \operatorname{dn} \nu \sin \psi, \\ z = \frac{k^{\frac{1}{2}} \Pi}{2a \mathcal{T}}, \end{cases}$$

where

$$A = 1 - \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu,$$

$$\mathcal{T} = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + [(A/\sqrt{k}) + \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu]^2,$$

$$\Pi = (A^2/k) - (\operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + \operatorname{cn}^2 \mu \operatorname{dn}^2 \mu \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu).$$

### Coordinate surfaces

$$(x^2 + y^2 + z^2)^2 + A(x^2 + y^2) + Bz^2 + C = 0,$$

where

$$A = -(R_1 + R_2),$$

$$B = \frac{4(R_1 + R_2) \operatorname{sn}^2 \mu - \left(4a^2 R_1 R_2 + \frac{k^2}{4a^2}\right)(1/k + \operatorname{sn}^2 \mu)^2}{k(1/k - \operatorname{sn}^2 \mu)^2}.$$

$$C = R_1 R_2,$$

$$R_1 = \frac{k(1 - \operatorname{dn}^2 \mu/(1+k))^2 \operatorname{sn}^2 \mu}{a^2 \left[ k \operatorname{sn}^2 \mu + \frac{1}{\sqrt{k}} \left( 1 - \frac{\operatorname{dn}^2 \mu}{1+k} \right) + (\operatorname{cn} \mu \operatorname{dn} \mu) \frac{\sqrt{k}}{1+k} \right]^2}.$$

$$R_2 = \frac{k \left( 1 - \frac{\operatorname{dn}^2 \mu}{1+k} \right)^2 \operatorname{sn}^2 \mu}{a^2 \left[ k \operatorname{sn}^2 \mu + \frac{1}{\sqrt{k}} \left( 1 - \frac{\operatorname{dn}^2 \mu}{1+k} \right) - \frac{\sqrt{k}}{1+k} \operatorname{cn} \mu \operatorname{dn} \mu \right]^2} \quad (\text{ring-cyclides, } \mu = \text{const}).$$

$$(x^2 + y^2 + z^2)^2 + A(x^2 + y^2) + Bz^2 + C = 0,$$

where

$$A = \frac{k^2}{8a^2} \left[ \frac{(1+k')(\operatorname{cn}^4 \nu + 6k \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu + k^2 \operatorname{sn}^4 \nu) \left( \frac{(1-k' \operatorname{sn}^2 \nu)}{k} - \frac{(1+k' \operatorname{sn}^2 \nu)}{1+k'} \right)^2}{\operatorname{dn}^2 \nu (\operatorname{cn}^2 \nu - k^2 \operatorname{sn}^2 \nu)^2} - 1 \right],$$

$$B = -\frac{k}{2a^2} \frac{(\operatorname{cn}^4 \nu + 6k \operatorname{sn}^2 \nu \operatorname{cn}^2 \nu + k^2 \operatorname{sn}^4 \nu)}{(\operatorname{cn}^2 \nu - k^2 \operatorname{sn}^2 \nu)^2},$$

$$C = \frac{k^2}{16a^4} \quad (\text{cap-cyclides, } \nu = \text{const}).$$

$$\tan \psi = y/x$$

(half-planes,  $\psi = \text{const}$ ).

### Stäckel matrix

$$[S] = \begin{bmatrix} -k^2 \operatorname{sn}^2 \mu & -1 & -\left(k^2 \operatorname{sn}^2 \mu + \frac{1}{\operatorname{sn}^2 \mu}\right) \\ \operatorname{dn}^2 \nu & 1 & \left(\operatorname{dn}^2 \nu + \frac{k^2}{\operatorname{dn}^2 \nu}\right) \\ 0 & 0 & 1 \end{bmatrix}.$$

$$S = \operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu = \operatorname{dn}^2 \mu - k'^2 \operatorname{sn}^2 \nu,$$

$$M_{11} = M_{21} = 1, \quad M_{31} = \frac{\Omega^2}{\operatorname{sn}^2 \mu \operatorname{dn}^2 \nu}.$$

### Metric coefficients

$$g_{11} = g_{22} = \left(\frac{A}{a \mathcal{T}}\right)^2 \Omega^2,$$

$$g_{33} = \left(\frac{A}{a \mathcal{T}}\right)^2 \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu,$$

$$g^{\frac{1}{2}} = \left(\frac{A}{a \mathcal{T}}\right)^{\frac{3}{2}} \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu,$$

where

$$\Omega^2 = (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu) (\operatorname{dn}^2 \nu - k^2 \operatorname{sn}^2 \mu),$$

$$\Upsilon = \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu + [A/\sqrt{k} + \operatorname{cn} \mu \operatorname{dn} \mu \operatorname{sn} \nu \operatorname{cn} \nu]^2.$$

$$R = A^{\frac{1}{2}} \Upsilon^{-\frac{1}{2}}, \quad Q = \left(\frac{A}{a \Upsilon}\right)^2 (1 - \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu),$$

$$f_1 = \operatorname{sn} \mu, \quad f_2 = \operatorname{dn} \nu, \quad f_3 = 1/a, \quad \alpha_1 = -1.$$

### Important equations

$$(ds)^2 = \left(\frac{A}{a \Upsilon}\right)^2 \Omega^2 [(d\mu)^2 + (d\nu)^2] + \left(\frac{A}{a \Upsilon}\right)^2 \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu (d\psi)^2.$$

$$\operatorname{grad} \varphi = \frac{a \Upsilon}{A \Omega} \left[ \mathbf{a}_\mu \frac{\partial \varphi}{\partial \mu} + \mathbf{a}_\nu \frac{\partial \varphi}{\partial \nu} \right] + \mathbf{a}_\psi \frac{a \Upsilon}{A \operatorname{sn} \mu \operatorname{dn} \nu} \frac{\partial \varphi}{\partial \psi}.$$

$$\operatorname{div} \mathbf{E} = \frac{a \Upsilon^3}{A^3 \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu} \times \left\{ \operatorname{dn} \nu \frac{\partial}{\partial \mu} \left[ \frac{A^2 \Omega \operatorname{sn} \mu}{\Upsilon^2} E_\mu \right] + \operatorname{sn} \mu \frac{\partial}{\partial \nu} \left[ \frac{A^2 \Omega \operatorname{dn} \nu}{\Upsilon^2} E_\nu \right] \right\} + \frac{a \Upsilon}{A \operatorname{sn} \mu \operatorname{dn} \nu} \frac{\partial E_\psi}{\partial \psi}.$$

$$\operatorname{curl} \mathbf{E} = \frac{a \Upsilon^2}{A^2 \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu} \begin{vmatrix} \mathbf{a}_\mu \Omega & \mathbf{a}_\nu \Omega & \mathbf{a}_\psi \operatorname{sn} \mu \operatorname{dn} \nu \\ \frac{\partial}{\partial \mu} & \frac{\partial}{\partial \nu} & \frac{\partial}{\partial \psi} \\ E_\mu A \Omega / \Upsilon & E_\nu A \Omega / \Upsilon & E_\psi A \operatorname{sn} \mu \operatorname{dn} \nu / \Upsilon \end{vmatrix}.$$

$$\nabla^2 \varphi = \frac{a^2 \Upsilon^3}{A^3 \Omega^2 \operatorname{sn} \mu \operatorname{dn} \nu} \times \left\{ \operatorname{dn} \nu \frac{\partial}{\partial \mu} \left[ \frac{A \operatorname{sn} \mu}{\Upsilon} \frac{\partial \varphi}{\partial \mu} \right] + \operatorname{sn} \mu \frac{\partial}{\partial \nu} \left[ \frac{A \operatorname{dn} \nu}{\Upsilon} \frac{\partial \varphi}{\partial \nu} \right] \right\} + \frac{a^2 \Upsilon^2}{A^2 \operatorname{sn}^2 \mu \operatorname{dn}^2 \nu} \frac{\partial^2 \varphi}{\partial \psi^2}.$$

### SEPARATION OF LAPLACE'S EQUATION, $\nabla^2 \varphi = 0$ .

$$\frac{1}{f_i} \frac{d}{du^i} \left( f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^3 \Phi_{ij} \alpha_j = 0,$$

where  $\alpha_1 = -1$  and  $U^1 = M(\mu)$ ,  $U^2 = N(\nu)$ ,  $U^3 = \Psi(\psi)$ ,

$$\varphi = A^{-\frac{1}{2}} \Upsilon^{\frac{1}{2}} M N \Psi.$$

### General case

$$\begin{cases} \frac{d^2 M}{d\mu^2} + \frac{\operatorname{cn} \mu \operatorname{dn} \mu}{\operatorname{sn} \mu} \frac{dM}{d\mu} + \left[ k^2 \operatorname{sn}^2 \mu - \alpha_2 - \alpha_3 \left( k^2 \operatorname{sn}^2 \mu + \frac{1}{\operatorname{sn}^2 \mu} \right) \right] M = 0, \\ \frac{d^2 N}{d\nu^2} - \frac{k'^2 \operatorname{sn} \nu \operatorname{cn} \nu}{\operatorname{dn} \nu} \frac{dN}{d\nu} + \left[ -\operatorname{dn}^2 \nu + \alpha_2 + \alpha_3 \left( \operatorname{dn}^2 \nu + \frac{k^2}{\operatorname{sn}^2 \nu} \right) \right] N = 0, \\ \frac{d^2 \Psi}{d\psi^2} + \alpha_3 \Psi = 0. \end{cases}$$

Substitution of  $\operatorname{sn}^2 \mu = z$  in the first equation, and  $\operatorname{dn}^2 \nu = z$  in the second equation, reduces both to the canonical form

$$\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z - a_1} + \frac{1}{z - a_2} + \frac{2}{z - a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z - a_1)(z - a_2)(z - a_3)^2} \right] Z = 0,$$

where, for  $M$  with  $\alpha_2 = p^2$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} a_1 &= b_1 = 1, \quad a_2 = c = 1/k^2, \quad a_3 = 0, \\ \bar{A}_0 &= -q^2 c, \quad \bar{A}_1 = -p^2 c, \quad \bar{A}_2 = 1 - q^2. \end{aligned}$$

For  $N$ , with  $\alpha_2 = p'^2$  and  $\alpha_3 = q^2$ ,

$$\begin{aligned} a_1 &= b = 1, \quad a_2 = c = k^2, \quad a_3 = 0, \\ \bar{A}_0 &= -q^2 c, \quad \bar{A}_1 = -p'^2, \quad \bar{A}_2 = 1 - q^2. \end{aligned}$$

Here  $p'^2 = p^2/k^2$ . These are Bôcher equations with designation {1122}, and their solutions are Wangerin functions,

$$Z = A \mathcal{S}_p^q(k, z) + B \mathcal{T}_p^q(k, z).$$

For  $\varphi = \varphi(u^1, u^2, u^3)$ ,

$$\begin{cases} M = A \mathcal{S}_p^q(k, \operatorname{sn}^2 \mu) + B \mathcal{T}_p^q(k, \operatorname{sn}^2 \mu), \\ N = A \mathcal{S}_p^q(i/k, \operatorname{dn}^2 \nu) + B \mathcal{T}_p^q(i/k, \operatorname{dn}^2 \nu), \\ \Psi = A \sin q \psi + B \cos q \psi, \quad \text{where } p'^2 = p^2/k^2. \end{cases}$$

For  $\varphi$  independent of  $\psi$ ,

$$\begin{cases} M = A \mathcal{S}_p(k, \operatorname{sn}^2 \mu) + B \mathcal{T}_p(k, \operatorname{sn}^2 \mu), \\ N = A \mathcal{S}_{p'}(i/k, \operatorname{dn}^2 \nu) + B \mathcal{T}_{p'}(i/k, \operatorname{dn}^2 \nu). \end{cases}$$

A summary of the coordinates obtained from elliptic functions is given in Table 4.04. Further information on the differential equations and their solutions may be found in Section VII.

TABLE 4.04.  
SEPARATION EQUATIONS OBTAINED IN FOUR COORDINATE SYSTEMS

|   |     | $a_1$ | $a_2$           | $a_3$       | $\bar{A}_0$ | $\bar{A}_1$                      | $\bar{A}_2$         | $\bar{A}_3$ |
|---|-----|-------|-----------------|-------------|-------------|----------------------------------|---------------------|-------------|
| J1 Rx<br>{1222}<br>$p'^2 = p^2/k^2$       | $M$ | 0     | 1               | $1/k^2 = c$ | $-p^2 c^2$  | $[p^2 c(1+c) - q^2(1-c)^2 + 2c]$ | $[-p^2 c + 2(1+c)]$ | 2           |
|   | $N$ | 0     | 1               | $k^2 = c$   | $-p'^2 c$   | $[p'^2(1+c) - q^2(1-c)^2 + 2c]$  | $[-p'^2 + 2(1+c)]$  | 2           |
| J1 Ry, J3 R<br>{1122}<br>$p'^2 = p^2/k^2$ | $M$ | 1     | $1/k^2 = c$     | 0           | $-q^2 c$    | $-p^2 c$                         | $1 - q^2$           | 0           |
|   | $N$ | 1     | $k^2 = c$       | 0           | $-q^2 c$    | $-p'^2$                          | $1 - q^2$           | 0           |
| J2 R<br>{1122}<br>$p'^2 = 1 - p^2$        | $M$ | 1     | $-(k'/k)^2 = c$ | 0           | $-q^2 c$    | $[p^2(1-c) - 1]$                 | $1 - q^2$           | 0           |
|   | $N$ | 1     | $-(k/k')^2 = c$ | 0           | $-q^2 c$    | $[-p'^2(1-c) + c]$               | $1 - q^2$           | 0           |

## Section V

### THE VECTOR HELMHOLTZ EQUATION

The vector Helmholtz equation, which occurs particularly in electromagnetic theory [19], is more complicated than the scalar Helmholtz equation and its separation presents new problems.

#### 5.01 CYLINDRICAL SYSTEMS

The vector Laplacian in general orthogonal coordinates is expressed by Eq. (1.11). For a *cylindrical* coordinate system, these results can be simplified because  $g_{33}$  is always unity [9]. If  $g_{11} = g_{22}$  also,

$$\begin{aligned}\mathcal{T} &= \frac{1}{g_{11}} \left\{ \frac{\partial}{\partial u^1} [(g_{11})^{\frac{1}{2}} E_1] + \frac{\partial}{\partial u^2} [(g_{11})^{\frac{1}{2}} E_2] \right\} + \frac{\partial E_z}{\partial z}, \\ \Gamma_1 &= \frac{\partial E_3}{\partial u^2} - (g_{11})^{\frac{1}{2}} \frac{\partial E_2}{\partial z}, \\ \Gamma_2 &= (g_{11})^{\frac{1}{2}} \frac{\partial E_1}{\partial z} - \frac{\partial E_z}{\partial u^1}, \\ \Gamma_3 &= \frac{1}{g_{11}} \left\{ \frac{\partial}{\partial u^1} [(g_{11})^{\frac{1}{2}} E_2] - \frac{\partial}{\partial u^2} [(g_{11})^{\frac{1}{2}} E_1] \right\}.\end{aligned}$$

Substitution into Eq. (1.11) gives the vector Laplacian in any *cylindrical* system with  $g_{11} = g_{22}$ :

$$\begin{aligned}(\hat{\star} \hat{\star} \mathbf{E})_1 &= \frac{1}{g_{11}} \left\{ \frac{\partial^2 E_1}{(\partial u^1)^2} + \frac{\partial^2 E_1}{(\partial u^2)^2} \right\} + \frac{\partial^2 E_1}{\partial z^2} + \frac{E_1}{2(g_{11})^2} \left[ \frac{\partial^2 g_{11}}{(\partial u^1)^2} + \frac{\partial^2 g_{11}}{(\partial u^2)^2} \right] \\ &\quad - \frac{3E_1}{4(g_{11})^3} \left[ \left( \frac{\partial g_{11}}{\partial u^1} \right)^2 + \left( \frac{\partial g_{11}}{\partial u^2} \right)^2 \right] + \frac{1}{(g_{11})^2} \left[ \frac{\partial g_{11}}{\partial u^2} \frac{\partial E_2}{\partial u^1} - \frac{\partial g_{11}}{\partial u^1} \frac{\partial E_2}{\partial u^2} \right],\end{aligned} \quad (5.01)$$

$$\begin{aligned}(\hat{\star} \hat{\star} \mathbf{E})_2 &= \frac{1}{g_{11}} \left\{ \frac{\partial^2 E_2}{(\partial u^1)^2} + \frac{\partial^2 E_2}{(\partial u^2)^2} \right\} + \frac{\partial^2 E_2}{\partial z^2} + \frac{E_2}{2(g_{11})^2} \left[ \frac{\partial^2 g_{11}}{(\partial u^1)^2} + \frac{\partial^2 g_{11}}{(\partial u^2)^2} \right] \\ &\quad - \frac{3E_2}{4(g_{11})^3} \left[ \left( \frac{\partial g_{11}}{\partial u^1} \right)^2 + \left( \frac{\partial g_{11}}{\partial u^2} \right)^2 \right] + \frac{1}{(g_{11})^2} \left[ \frac{\partial g_{11}}{\partial u^1} \frac{\partial E_1}{\partial u^2} - \frac{\partial g_{11}}{\partial u^2} \frac{\partial E_1}{\partial u^1} \right],\end{aligned} \quad (5.02)$$

$$(\hat{\star} \hat{\star} \mathbf{E})_z = \frac{1}{g_{11}} \left\{ \frac{\partial^2 E_z}{(\partial u^1)^2} + \frac{\partial^2 E_z}{(\partial u^2)^2} \right\} + \frac{\partial^2 E_z}{\partial z^2}. \quad (5.03)$$

Equation (5.03) shows that the  $z$ -component of the vector Laplacian has exactly the form of the scalar Laplacian:

$$(\hat{\star} \hat{\star} \mathbf{E})_z = \nabla^2 E_z. \quad (5.04)$$

But the other components of  $\hat{\star} \hat{\star} \mathbf{E}$  contain additional terms [20].

Details for the separation of the vector Helmholtz equation in cylindrical systems are given in Table 5.01. For rectangular coordinates, the three components of the vector Helmholtz equation are identical with the scalar Helmholtz equation. Thus separation presents no problem and the solutions are those given

in Section I. Circular-cylinder coordinates, though less satisfactory than rectangular coordinates, still allow solutions in a number of cases. For the other cylindrical systems, separation appears to be possible only for the  $z$ -component and to occur only in the four cylindrical systems in which the scalar wave equation is known to separate. Solutions for  $E_z$  are identical with those previously given for  $\varphi$ .

## 5.02 ROTATIONAL SYSTEMS

A simplification of Eq. (1.11) is also effected in coordinate systems having axial symmetry. Here the metric coefficients do not contain the third independent variable  $u^3 = \psi$ . If we also stipulate that  $g_{11} = g_{22}$ , then

$$\begin{aligned}\mathcal{T} &= \frac{1}{(g_{11})^{\frac{1}{2}}} \left[ \frac{\partial E_1}{\partial u^1} + \frac{\partial E_2}{\partial u^2} \right] + \frac{1}{(g_{33})^{\frac{1}{2}}} \frac{\partial E_\psi}{\partial \psi} \\ &\quad + \frac{E_1}{2(g_{11})^{\frac{1}{2}}} \left[ \frac{1}{g_{11}} \frac{\partial g_{11}}{\partial u^1} + \frac{1}{g_{33}} \frac{\partial g_{33}}{\partial u^1} \right] + \frac{E_2}{2(g_{11})^{\frac{1}{2}}} \left[ \frac{1}{g_{11}} \frac{\partial g_{11}}{\partial u^2} + \frac{1}{g_{33}} \frac{\partial g_{33}}{\partial u^2} \right], \\ I_1 &= \frac{\partial E_\psi}{\partial u^2} - \left( \frac{g_{11}}{g_{33}} \right)^{\frac{1}{2}} \frac{\partial E_2}{\partial \psi} + \frac{E_\psi}{2g_{33}} \frac{\partial g_{33}}{\partial u^2}, \\ I_2 &= \left( \frac{g_{11}}{g_{33}} \right)^{\frac{1}{2}} \frac{\partial E_1}{\partial \psi} - \frac{\partial E_\psi}{\partial u^1} - \frac{E_\psi}{2g_{33}} \frac{\partial g_{33}}{\partial u^1}, \\ I_3 &= \left( \frac{g_{33}}{g_{11}} \right)^{\frac{1}{2}} \left[ \frac{\partial E_2}{\partial u^1} - \frac{\partial E_1}{\partial u^2} + \frac{1}{2g_{11}} \left( \frac{\partial g_{11}}{\partial u^1} E_2 - \frac{\partial g_{11}}{\partial u^2} E_1 \right) \right],\end{aligned}$$

Therefore,

$$\begin{aligned}(\star\star\star \mathbf{E})_1 &= \frac{1}{g_{11}} \left[ \frac{\partial^2 E_1}{(\partial u^1)^2} + \frac{\partial^2 E_1}{(\partial u^2)^2} \right] + \frac{1}{g_{33}} \frac{\partial^2 E_1}{\partial \psi^2} + \frac{1}{2g_{11}g_{33}} \left[ \frac{\partial g_{33}}{\partial u^1} \frac{\partial E_1}{\partial u^1} + \frac{\partial g_{33}}{\partial u^2} \frac{\partial E_2}{\partial u^2} \right] \\ &\quad + \frac{E_1}{2g_{11}} \left\{ \frac{1}{g_{11}} \left[ \frac{\partial^2 g_{11}}{(\partial u^1)^2} + \frac{\partial^2 g_{11}}{(\partial u^2)^2} \right] - \frac{3}{2(g_{11})^2} \left[ \left( \frac{\partial g_{11}}{\partial u^1} \right)^2 + \left( \frac{\partial g_{11}}{\partial u^2} \right)^2 \right] \right\} \\ &\quad - \frac{1}{2g_{11}g_{33}} \left[ \frac{\partial g_{11}}{\partial u^1} \frac{\partial g_{33}}{\partial u^1} - \frac{\partial g_{11}}{\partial u^2} \frac{\partial g_{33}}{\partial u^2} \right] + \frac{1}{g_{33}} \left[ \frac{\partial^2 g_{33}}{(\partial u^1)^2} - \frac{1}{g_{33}} \left( \frac{\partial g_{33}}{\partial u^1} \right)^2 \right] \\ &\quad + \frac{1}{(g_{11})^2} \left[ \frac{\partial g_{11}}{\partial u^2} \frac{\partial E_2}{\partial u^1} - \frac{\partial g_{11}}{\partial u^1} \frac{\partial E_2}{\partial u^2} \right] \\ &\quad + \frac{E_2}{2g_{11}g_{22}} \left[ \frac{\partial^2 g_{33}}{\partial u^1 \partial u^2} - \frac{\partial g_{33}}{\partial u^2} \left( \frac{1}{g_{11}} \frac{\partial g_{11}}{\partial u^1} + \frac{1}{g_{33}} \frac{\partial g_{33}}{\partial u^1} \right) \right] \\ &\quad - \frac{1}{(g_{11})^{\frac{1}{2}}(g_{33})^{\frac{3}{2}}} \frac{\partial g_{33}}{\partial u^1} \frac{\partial E_\psi}{\partial \psi}. \\ (\star\star\star \mathbf{E})_2 &= \frac{1}{g_{11}} \left[ \frac{\partial^2 E_2}{(\partial u^1)^2} + \frac{\partial^2 E_2}{(\partial u^2)^2} \right] + \frac{1}{g_{33}} \frac{\partial^2 E_2}{\partial \psi^2} + \frac{1}{2g_{11}g_{33}} \left[ \frac{\partial g_{33}}{\partial u^1} \frac{\partial E_2}{\partial u^1} + \frac{\partial g_{33}}{\partial u^2} \frac{\partial E_2}{\partial u^2} \right] \\ &\quad + \frac{E_2}{2g_{11}} \left\{ \frac{1}{g_{11}} \left[ \frac{\partial^2 g_{11}}{(\partial u^1)^2} + \frac{\partial^2 g_{11}}{(\partial u^2)^2} \right] - \frac{3}{2(g_{11})^2} \left[ \left( \frac{\partial g_{11}}{\partial u^1} \right)^2 + \left( \frac{\partial g_{11}}{\partial u^2} \right)^2 \right] \right\} \\ &\quad - \frac{1}{2g_{11}g_{33}} \left[ \frac{\partial g_{11}}{\partial u^2} \frac{\partial g_{33}}{\partial u^2} - \frac{\partial g_{11}}{\partial u^1} \frac{\partial g_{33}}{\partial u^1} \right] + \frac{1}{g_{33}} \left[ \frac{\partial^2 g_{33}}{(\partial u^2)^2} - \frac{1}{g_{33}} \left( \frac{\partial g_{33}}{\partial u^2} \right)^2 \right] \\ &\quad + \frac{1}{(g_{11})^2} \left[ \frac{\partial g_{11}}{\partial u^1} \frac{\partial E_1}{\partial u^2} - \frac{\partial g_{11}}{\partial u^2} \frac{\partial E_1}{\partial u^1} \right] \\ &\quad + \frac{E_1}{2g_{11}g_{33}} \left[ \frac{\partial^2 g_{33}}{\partial u^1 \partial u^2} - \frac{\partial g_{33}}{\partial u^1} \left( \frac{1}{g_{11}} \frac{\partial g_{11}}{\partial u^2} + \frac{1}{g_{33}} \frac{\partial g_{33}}{\partial u^2} \right) \right] \\ &\quad - \frac{1}{(g_{11})^{\frac{1}{2}}(g_{33})^{\frac{3}{2}}} \frac{\partial g_{33}}{\partial u^2} \frac{\partial E_\psi}{\partial \psi},\end{aligned}$$

$$\begin{aligned} (\nabla^2 \mathbf{E})_\psi &= \frac{1}{g_{11}} \left[ \frac{\partial^2 E_\psi}{(\partial u^1)^2} + \frac{\partial^2 E_\psi}{(\partial u^2)^2} \right] + \frac{1}{g_{33}} \frac{\partial^2 E_\psi}{\partial \psi^2} + \frac{1}{2g_{11} g_{33}} \left[ \frac{\partial g_{33}}{\partial u^1} \frac{\partial E_\psi}{\partial u^1} + \frac{\partial g_{33}}{\partial u^2} \frac{\partial E_\psi}{\partial u^2} \right] \\ &\quad + \frac{E_\psi}{2g_{11} g_{33}} \left\{ \left[ \frac{\partial^2 g_{33}}{(\partial u^1)^2} + \frac{\partial^2 g_{33}}{(\partial u^2)^2} \right] - \frac{1}{g_{33}} \left[ \left( \frac{\partial g_{33}}{\partial u^1} \right)^2 + \left( \frac{\partial g_{33}}{\partial u^2} \right)^2 \right] \right\} \\ &\quad + \frac{1}{(g_{11})^{\frac{1}{2}} (g_{33})^{\frac{3}{2}}} \left[ \frac{\partial g_{33}}{\partial u^1} \frac{\partial E_1}{\partial \psi} + \frac{\partial g_{33}}{\partial u^2} \frac{\partial E_2}{\partial \psi} \right]. \end{aligned}$$

If the field is axially symmetric, the  $\psi$ -component of the vector Helmholtz equation reduces to

$$\begin{aligned} \frac{1}{g_{11}} \left[ \frac{\partial^2 E_\psi}{(\partial u^1)^2} + \frac{\partial^2 E_\psi}{(\partial u^2)^2} \right] + \frac{1}{2g_{11} g_{33}} \left[ \frac{\partial g_{33}}{\partial u^1} \frac{\partial E_\psi}{\partial u^1} + \frac{\partial g_{33}}{\partial u^2} \frac{\partial E_\psi}{\partial u^2} \right] \\ + \left[ \kappa^2 + \frac{1}{2g_{11} g_{33}} \left\{ \left[ \frac{\partial^2 g_{33}}{(\partial u^1)^2} + \frac{\partial^2 g_{33}}{(\partial u^2)^2} \right] - \frac{1}{g_{33}} \left[ \left( \frac{\partial g_{33}}{\partial u^1} \right)^2 + \left( \frac{\partial g_{33}}{\partial u^2} \right)^2 \right] \right\} \right] E_\psi = 0. \end{aligned}$$

This equation is simply separable in the rotational coordinate systems in which the scalar Helmholtz equation separates. Details are given in Table 5.02.

TABLE 5.01.  
THE VECTOR HELMHOLTZ EQUATION IN CYLINDRICAL SYSTEMS

FIG. 1.01. RECTANGULAR COORDINATES ( $x, y, z$ )

$$\begin{aligned} g_{11} &= g_{22} = g_{33} = g^{\frac{1}{2}} = 1, \\ T &= \frac{\partial E_1}{\partial u^1} + \frac{\partial E_2}{\partial u^2} + \frac{\partial E_3}{\partial u^3}, \\ I_1 &= \frac{\partial E_3}{\partial u^2} - \frac{\partial E_2}{\partial u^3}, \\ I_2 &= \frac{\partial E_1}{\partial u^3} - \frac{\partial E_3}{\partial u^1}, \\ I_3 &= \frac{\partial E_2}{\partial u^1} - \frac{\partial E_1}{\partial u^2}. \end{aligned}$$

$$\nabla^2 \mathbf{E} = \mathbf{a}_x V^2 E_x + \mathbf{a}_y V^2 E_y + \mathbf{a}_z V^2 E_z.$$

Thus the vector Helmholtz equation in rectangular coordinates reduces to three ordinary scalar Helmholtz equations:

$$\begin{cases} V^2 E_x + \kappa^2 E_x = 0, \\ V^2 E_y + \kappa^2 E_y = 0, \\ V^2 E_z + \kappa^2 E_z = 0. \end{cases}$$

Solutions are given in Section I.

FIG. 1.02. CIRCULAR-CYLINDER COORDINATES ( $r, \psi, z$ )

$$\begin{aligned} g_{11} &= 1, \quad g_{22} = r^2, \quad g_{33} = 1, \quad g^{\frac{1}{2}} = r. \\ T &= \frac{\partial E_r}{\partial r} + \frac{E_r}{r} + \frac{1}{r} \frac{\partial E_\psi}{\partial \psi} + \frac{\partial E_z}{\partial z}, \\ I_1 &= \frac{1}{r} \frac{\partial E_z}{\partial \psi} - \frac{\partial E_\psi}{\partial z}, \\ I_2 &= r \left[ \frac{\partial E_r}{\partial z} - \frac{\partial E_z}{\partial r} \right], \\ I_3 &= \frac{\partial E_\psi}{\partial r} + \frac{E_\psi}{r} - \frac{1}{r} \frac{\partial E_r}{\partial \psi}. \end{aligned}$$

From Eq. (1.11), the vector Helmholtz equation is

$$\begin{aligned}\frac{\partial^2 E_r}{\partial r^2} + \frac{1}{r} \frac{\partial E_r}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_r}{\partial \psi^2} + \frac{\partial^2 E_r}{\partial z^2} - \frac{2}{r^2} \frac{\partial E_\varphi}{\partial \psi} + (\kappa^2 - 1/r^2) E_r &= 0, \\ \frac{\partial^2 E_\varphi}{\partial r^2} + \frac{1}{r} \frac{\partial E_\varphi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_\varphi}{\partial \psi^2} + \frac{\partial^2 E_\varphi}{\partial z^2} + \frac{2}{r^2} \frac{\partial E_r}{\partial \psi} + (\kappa^2 - 1/r^2) E_\varphi &= 0, \\ \frac{\partial^2 E_z}{\partial r^2} + \frac{1}{r} \frac{\partial E_z}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_z}{\partial \psi^2} + \frac{\partial^2 E_z}{\partial z^2} + \kappa^2 E_z &= 0.\end{aligned}$$

These equations can be written

$$\left\{ \begin{array}{l} \nabla^2 E_r - \frac{2}{r^2} \frac{\partial E_\varphi}{\partial \psi} + (\kappa^2 - 1/r^2) E_r = 0, \\ \nabla^2 E_\varphi + \frac{2}{r^2} \frac{\partial E_r}{\partial \psi} + (\kappa^2 - 1/r^2) E_\varphi = 0, \\ \nabla^2 E_z + \kappa^2 E_z = 0. \end{array} \right.$$

### Solutions

For  $\mathbf{E} = \mathbf{a}_r E_r$ ,

$$\left\{ \begin{array}{l} \frac{\partial^2 E_r}{\partial r^2} + \frac{1}{r} \frac{\partial E_r}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_r}{\partial \psi^2} + \frac{\partial^2 E_r}{\partial z^2} + (\kappa^2 - 1/r^2) E_r = 0, \\ \frac{\partial E_r}{\partial \psi} = 0. \end{array} \right.$$

Because of the second equation,  $E_r$  must be independent of  $\psi$  and the vector Helmholtz equation is

$$\frac{\partial^2 E_r}{\partial r^2} + \frac{1}{r} \frac{\partial E_r}{\partial r} + \frac{\partial^2 E_r}{\partial z^2} + (\kappa^2 - 1/r^2) E_r = 0.$$

Thus, for  $\mathbf{E} = \mathbf{a}_r E_r(r, z)$ , separation equations and solutions are

$$\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + (q^2 - 1/r^2) R = 0, \quad \{24\} \quad R = A J_1(q r) + B Y_1(q r).$$

$$\frac{d^2 Z}{dz^2} + (\kappa^2 - q^2) Z = 0, \quad \{04\} \quad Z = A \sin(\kappa^2 - q^2)^{1/2} z + B \cos(\kappa^2 - q^2)^{1/2} z.$$

For  $\mathbf{E} = \mathbf{a}_r E_r(r)$ ,

$$\frac{d^2 E_r}{dr^2} + \frac{1}{r} \frac{dE_r}{dr} + (\kappa^2 - 1/r^2) E_r = 0, \quad \{24\} \quad E_r = A J_1(\kappa r) + B Y_1(\kappa r).$$

For  $\mathbf{E} = \mathbf{a}_\varphi E_\varphi$ ,

$$\left\{ \begin{array}{l} \frac{\partial^2 E_\varphi}{\partial r^2} + \frac{1}{r} \frac{\partial E_\varphi}{\partial r} + \frac{\partial^2 E_\varphi}{\partial z^2} + \frac{1}{r^2} \frac{\partial^2 E_\varphi}{\partial \psi^2} + (\kappa^2 - 1/r^2) E_\varphi = 0, \\ \frac{\partial E_\varphi}{\partial \psi} = 0. \end{array} \right.$$

Because of the second equation,  $E_\varphi$  is independent of  $\psi$  and the Helmholtz equation is

$$\frac{\partial^2 E_\varphi}{\partial r^2} + \frac{1}{r} \frac{\partial E_\varphi}{\partial r} + \frac{\partial^2 E_\varphi}{\partial z^2} + (\kappa^2 - 1/r^2) E_\varphi = 0.$$

Separation equations and solutions, for  $\mathbf{E} = \mathbf{a}_\psi E_\psi(r, z)$ , are

$$\begin{aligned}\frac{d^2R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + (q^2 - 1/r^2) R = 0, & \quad \{24\} \quad R = A \mathcal{J}_1(q r) + B \mathcal{Y}_1(q r). \\ \frac{d^2Z}{dz^2} + (\kappa^2 - q^2) Z = 0, & \quad \{04\} \quad Z = A \sin(\kappa^2 - q^2)^{\frac{1}{2}} z + B \cos(\kappa^2 - q^2)^{\frac{1}{2}} z.\end{aligned}$$

For  $\mathbf{E} = \mathbf{a}_\psi E_\psi(r)$ ,

$$\frac{d^2E_\psi}{dr^2} + \frac{1}{r} \frac{dE_\psi}{dr} + (\kappa^2 - 1/r^2) E_\psi = 0, \quad \{24\} \quad E_\psi = A \mathcal{J}_1(\kappa r) + B \mathcal{Y}_1(\kappa r).$$

The differential equation in  $E_z$  can always be separated to give solutions identical with those for the scalar Helmholtz equation, Table 1.02. Evidently these results apply, irrespective of other components that  $\mathbf{E}$  may have.

**FIG. 1.03. ELLIPTIC-CYLINDER COORDINATES ( $\eta, \psi, z$ )**

$$g_{11} = g_{22} = g^1 = a^2(\cosh^2 \eta - \cos^2 \psi), \quad g_{33} = 1.$$

The vector Helmholtz equation is obtained from Eqs. (5.01), (5.02), (5.03):

$$\begin{aligned}& \frac{1}{a^2(\cosh^2 \eta - \cos^2 \psi)} \left\{ \frac{\partial^2 E_\eta}{\partial \eta^2} + \frac{\partial^2 E_\eta}{\partial \psi^2} \right\} + \frac{\partial^2 E_\eta}{\partial z^2} \\& + \frac{2}{a^2(\cosh^2 \eta - \cos^2 \psi)^2} \left\{ \cos \psi \sin \psi \frac{\partial E_\psi}{\partial \eta} - \cosh \eta \sinh \eta \frac{\partial E_\psi}{\partial \psi} \right\} \\& + \left\{ \kappa^2 - \frac{5(\cosh^2 \eta - \sin^2 \psi)}{4a^2(\cosh^2 \eta - \cos^2 \psi)^2} \right\} E_\eta = 0, \\& \frac{1}{a^2(\cosh^2 \eta - \cos^2 \psi)} \left\{ \frac{\partial^2 E_\psi}{\partial \eta^2} + \frac{\partial^2 E_\psi}{\partial \psi^2} \right\} + \frac{\partial^2 E_\psi}{\partial z^2} \\& + \frac{2}{a^2(\cosh^2 \eta - \cos^2 \psi)^2} \left\{ \cosh \eta \sinh \eta \frac{\partial E_\eta}{\partial \psi} - \cos \psi \sin \psi \frac{\partial E_\eta}{\partial \eta} \right\} \\& + \left\{ \kappa^2 - \frac{5(\cosh^2 \eta - \sin^2 \psi)}{4a^2(\cosh^2 \eta - \cos^2 \psi)^2} \right\} E_\psi = 0, \\& \frac{1}{a^2(\cosh^2 \eta - \cos^2 \psi)} \left\{ \frac{\partial^2 E_z}{\partial \eta^2} + \frac{\partial^2 E_z}{\partial \psi^2} \right\} + \frac{\partial^2 E_z}{\partial z^2} + \kappa^2 E_z = 0.\end{aligned}$$

Since the form of the equation for  $E_z$  is exactly the same as for the scalar Helmholtz equation in elliptic-cylinder coordinates, the solutions are given in Table 1.03.

**FIG. 1.04. PARABOLIC-CYLINDER COORDINATES ( $\mu, \nu, z$ )**

$$g_{11} = g_{22} = g^1 = \mu^2 + \nu^2, \quad g_{33} = 1.$$

The vector Helmholtz equation is obtained from Eqs. (5.01), (5.02), (5.03):

$$\begin{aligned}& \frac{1}{\mu^2 + \nu^2} \left\{ \frac{\partial^2 E_\mu}{\partial \mu^2} + \frac{\partial^2 E_\mu}{\partial \nu^2} \right\} + \frac{\partial^2 E_\mu}{\partial z^2} \\& + \frac{2}{(\mu^2 + \nu^2)^2} \left\{ \nu \frac{\partial E_\nu}{\partial \mu} - \mu \frac{\partial E_\nu}{\partial \nu} \right\} + \left\{ \kappa^2 - \frac{1}{(\mu^2 + \nu^2)^2} \right\} E_\mu = 0, \\& \frac{1}{\mu^2 + \nu^2} \left\{ \frac{\partial^2 E_\nu}{\partial \mu^2} + \frac{\partial^2 E_\nu}{\partial \nu^2} \right\} + \frac{\partial^2 E_\nu}{\partial z^2} \\& + \frac{2}{(\mu^2 + \nu^2)^2} \left\{ \mu \frac{\partial E_\mu}{\partial \nu} - \nu \frac{\partial E_\mu}{\partial \mu} \right\} + \left\{ \kappa^2 - \frac{1}{(\mu^2 + \nu^2)^2} \right\} E_\nu = 0, \\& \frac{1}{\mu^2 + \nu^2} \left\{ \frac{\partial^2 E_z}{\partial \mu^2} + \frac{\partial^2 E_z}{\partial \nu^2} \right\} + \frac{\partial^2 E_z}{\partial z^2} + \kappa^2 E_z = 0.\end{aligned}$$

Solutions for  $E_z$  are given in Table 1.04.

## TABLE 5.02. ROTATIONAL COORDINATES

FIG. 1.05. SPHERICAL COORDINATES  $(r, \theta, \psi)$ 

$$g_{11} = 1, \quad g_{22} = r^2, \quad g_{33} = r^2 \sin^2 \theta, \quad g^{\frac{1}{2}} = r^2 \sin \theta.$$

The vector Helmholtz equation has the three components:

$$\begin{aligned} \frac{\partial^2 E_r}{\partial r^2} + \frac{2}{r} \frac{\partial E_r}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_r}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial E_r}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_r}{\partial \psi^2} \\ - \frac{2}{r^2} \frac{\partial E_\theta}{\partial \theta} - \frac{2 \cot \theta}{r^2} E_\theta - \frac{2}{r^2 \sin \theta} \frac{\partial E_\psi}{\partial \psi} + (\kappa^2 - 2/r^2) E_r = 0, \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 E_\theta}{\partial r^2} + \frac{2}{r} \frac{\partial E_\theta}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_\theta}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial E_\theta}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_\theta}{\partial \psi^2} \\ + \frac{2}{r^2} \frac{\partial E_r}{\partial \theta} - \frac{2 \cot \theta}{r^2 \sin \theta} \frac{\partial E_\psi}{\partial \psi} + \left( \kappa^2 - \frac{1}{r^2 \sin^2 \theta} \right) E_\theta = 0, \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 E_\psi}{\partial r^2} + \frac{2}{r} \frac{\partial E_\psi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 E_\psi}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial E_\psi}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_\psi}{\partial \psi^2} \\ + \frac{2}{r^2 \sin \theta} \frac{\partial E_r}{\partial \psi} + \frac{2 \cot \theta}{r^2 \sin \theta} \frac{\partial E_\theta}{\partial \psi} + \left( \kappa^2 - \frac{1}{r^2 \sin^2 \theta} \right) E_\psi = 0. \end{aligned}$$

These equations can be written

$$\begin{cases} \nabla^2 E_r - \frac{2}{r^2} \frac{\partial E_\theta}{\partial \theta} - \frac{2 \cot \theta}{r^2} E_\theta - \frac{2}{r^2 \sin \theta} \frac{\partial E_\psi}{\partial \psi} + (\kappa^2 - 2/r^2) E_r = 0, \\ \nabla^2 E_\theta + \frac{2}{r^2} \frac{\partial E_r}{\partial \theta} - \frac{2 \cot \theta}{r^2 \sin \theta} \frac{\partial E_\psi}{\partial \psi} + \left( \kappa^2 - \frac{1}{r^2 \sin^2 \theta} \right) E_\theta = 0, \\ \nabla^2 E_\psi + \frac{2}{r^2 \sin \theta} \frac{\partial E_r}{\partial \psi} + \frac{2 \cot \theta}{r^2 \sin \theta} \frac{\partial E_\theta}{\partial \psi} + \left( \kappa^2 - \frac{1}{r^2 \sin^2 \theta} \right) E_\psi = 0. \end{cases}$$

## Solutions

For  $\mathbf{E} = \mathbf{a}_r E_r$ ,

$$\begin{cases} \nabla^2 E_r + (\kappa^2 - 2/r^2) E_r = 0, \\ \frac{\partial E_r}{\partial \theta} = 0, \quad \frac{\partial E_r}{\partial \psi} = 0. \end{cases}$$

Thus, if  $\mathbf{E}$  is in the radial direction, it must be a function of  $r$  alone, and

$$\frac{d^2 E_r}{dr^2} + \frac{2}{r} \frac{d E_r}{dr} + [\kappa^2 - 2/r^2] E_r = 0, \quad \{24\} \quad E_r = r^{-\frac{1}{2}} [A J_{\frac{1}{2}}(\kappa r) + B J_{-\frac{1}{2}}(\kappa r)].$$

For  $\mathbf{E} = \mathbf{a}_\theta E_\theta$ ,

$$\begin{cases} \nabla^2 E_\theta + \left( \kappa^2 - \frac{1}{r^2 \sin^2 \theta} \right) E_\theta = 0, \\ \frac{\partial E_\theta}{\partial \theta} + \cot \theta E_\theta = 0, \quad \frac{\partial E_\theta}{\partial \psi} = 0. \end{cases}$$

Thus if  $\mathbf{E}$  is in the  $\theta$ -direction, it must depend on both  $r$  and  $\theta$ . Separation equations and solutions are

$$\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{d R}{dr} + \kappa^2 R = 0, \quad \{04\} \quad R = \frac{1}{r} [A \sin \kappa r + B \cos \kappa r].$$

$$\frac{d \Theta}{d \theta} + (\cot \theta) \Theta = 0, \quad \Theta = \frac{A}{\sin \theta}.$$

For  $\mathbf{E} = \mathbf{a}_\psi E_\psi$ ,

$$\begin{cases} \nabla^2 E_\psi + \left( \kappa^2 - \frac{1}{r^2 \sin^2 \theta} \right) E_\psi = 0, \\ \frac{\partial E_\psi}{\partial \psi} = 0. \end{cases}$$

Thus, if  $\mathbf{E}$  is in the  $\psi$ -direction, the field must be axially symmetric,  $\mathbf{E} = \mathbf{a}_\psi E_\psi(r, \theta)$ . If  $E_\psi = R(r) \Theta(\theta)$ , separation equations and solutions are

$$\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \left[ \kappa^2 - \frac{p(p+1)}{r^2} \right] R = 0,$$

$$\{24\} \quad R = r^{-\frac{1}{2}} [A J_{p+\frac{1}{2}}(\kappa r) + B J_{-(p+\frac{1}{2})}(\kappa r)].$$

$$\frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left[ p(p+1) - \frac{1}{\sin^2 \theta} \right] \Theta = 0,$$

$$\{222\} \quad \Theta = A P_p^1(\cos \theta) + B Q_p^1(\cos \theta).$$

FIG. 1.06. PROLATE SPHEROIDAL COORDINATES ( $\eta, \theta, \psi$ )

$$g_{11} = g_{22} = a^2 (\sinh^2 \eta + \sin^2 \theta), \quad g_{33} = a^2 \sinh^2 \eta \sin^2 \theta,$$

$$g^{\frac{1}{2}} = a^2 (\sinh^2 \eta + \sin^2 \theta) \sinh \eta \sin \theta.$$

For  $\mathbf{E}$  independent of  $\psi$ , the  $\psi$ -component of the vector Helmholtz equations is

$$\begin{aligned} \frac{\partial^2 E_\psi}{\partial \eta^2} + \coth \eta \frac{\partial E_\psi}{\partial \eta} + \frac{\partial^2 E_\psi}{\partial \theta^2} + \cot \theta \frac{\partial E_\psi}{\partial \theta} \\ + \left[ \kappa^2 a^2 (\sinh^2 \eta + \sin^2 \theta) - \left( \frac{1}{\sinh^2 \eta} + \frac{1}{\sin^2 \theta} \right) \right] E_\psi = 0. \end{aligned}$$

For  $E_\psi = H(\eta) \cdot \Theta(\theta)$ , separation equations are

$$\begin{cases} \frac{d^2 H}{d\eta^2} + \coth \eta \frac{dH}{d\eta} + \left[ \kappa^2 a^2 \sinh^2 \eta - p(p+1) - \frac{1}{\sinh^2 \eta} \right] H = 0, \\ \frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left[ \kappa^2 a^2 \sin^2 \theta + p(p+1) - \frac{1}{\sin^2 \theta} \right] \Theta = 0. \end{cases}$$

Solutions are

$$H = A P_p^1(\kappa a, \cosh \eta) + B Q_p^1(\kappa a, \cosh \eta),$$

$$\Theta = A P_p^1(\kappa a, \cos \theta) + B Q_p^1(\kappa a, \cos \theta).$$

FIG. 1.07. OBLATE SPHEROIDAL COORDINATES ( $\eta, \theta, \psi$ )

$$g_{11} = g_{22} = a^2 (\cosh^2 \eta - \sin^2 \theta), \quad g_{33} = a^2 \cosh^2 \eta \sin^2 \theta,$$

$$g^{\frac{1}{2}} = a^3 (\cosh^2 \eta - \sin^2 \theta) \cosh \eta \sin \theta.$$

For  $\mathbf{E}$  independent of  $\psi$ , the  $\psi$ -component of the vector Helmholtz equation is

$$\begin{aligned} \frac{\partial^2 E_\psi}{\partial \eta^2} + \tanh \eta \frac{\partial E_\psi}{\partial \eta} + \frac{\partial^2 E_\psi}{\partial \theta^2} + \cot \theta \frac{\partial E_\psi}{\partial \theta} \\ + \left[ \kappa^2 a^2 (\cosh^2 \eta - \sin^2 \theta) + \left( \frac{1}{\cosh^2 \eta} - \frac{1}{\sin^2 \theta} \right) \right] E_\psi = 0. \end{aligned}$$

For  $E_\psi = H(\eta) \cdot \Theta(\theta)$ , separation equations are

$$\begin{cases} \frac{d^2H}{d\eta^2} + \tanh \eta \frac{dH}{d\eta} + \left[ \kappa^2 a^2 \cosh^2 \eta - p(p+1) + \frac{1}{\cosh^2 \eta} \right] H = 0, \\ \frac{d^2\Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left[ -\kappa^2 a^2 \sin^2 \theta + p(p+1) - \frac{1}{\sin^2 \theta} \right] \Theta = 0. \end{cases}$$

Solutions are

$$\begin{aligned} H &= A \mathcal{P}_p^1(i \kappa a, i \sinh \eta) + B \mathcal{Q}_p^1(i \kappa a, i \sinh \eta), \\ \Theta &= A \mathcal{P}_p^1(i \kappa a, \cos \theta) + B \mathcal{Q}_p^1(i \kappa a, \cos \theta). \end{aligned}$$

### FIG. 1.08. PARABOLIC COORDINATES ( $\mu, \nu, \psi$ )

$$g_{11} = g_{22} = \mu^2 + \nu^2, \quad g_{33} = \mu^2 \nu^2, \quad g^3 = \mu \nu (\mu^2 + \nu^2).$$

For  $\mathbf{E}$  independent of  $\psi$ , the  $\psi$ -component of the vector Helmholtz equation is

$$\frac{\partial^2 E_\psi}{\partial \mu^2} + \frac{1}{\mu} \frac{\partial E_\psi}{\partial \mu} + \frac{\partial^2 E_\psi}{\partial \nu^2} + \frac{1}{\nu} \frac{\partial E_\psi}{\partial \nu} + \left[ \kappa^2 (\mu^2 + \nu^2) - \left( \frac{1}{\mu^2} + \frac{1}{\nu^2} \right) \right] E_\psi = 0.$$

For  $E_\psi = M(\mu) \cdot N(\nu)$ , separation equations are

$$\begin{cases} \frac{d^2M}{d\mu^2} + \frac{1}{\mu} \frac{dM}{d\mu} + [\kappa^2 \mu^2 - q^2 - 1/\mu^2] M = 0, \\ \frac{d^2N}{d\nu^2} + \frac{1}{\nu} \frac{dN}{d\nu} + [\kappa^2 \nu^2 + q^2 - 1/\nu^2] N = 0. \end{cases}$$

Solutions are

$$\begin{aligned} M &= A \mathcal{J}_1(\kappa, q, i \mu) + B \mathcal{Y}_1(\kappa, q, i \mu), \\ N &= A \mathcal{J}_1(\kappa, q, \nu) + B \mathcal{Y}_1(\kappa, q, \nu). \end{aligned}$$

## Section VI

# DIFFERENTIAL EQUATIONS

Previous sections have treated the separation of the Laplace and Helmholtz equations in 40 coordinate systems. In this section, all the separation equations are tabulated in a systematic manner. Each equation is designated in terms of its singularities in the complex plane, and the general solutions of the differential equations are listed.

A comprehensive view of the separation equations of mathematical physics seems to have been first attempted by FELIX KLEIN [21] and MAXIME BÔCHER [22] in 1894. Their method was to consider the various separation equations as degenerate cases of a “generalized Lamé equation” with 5 singularities. Unfortunately, this scheme does not seem to cover all the necessary equations. Their work was extended by INCE [23], who developed a specification in terms of singularities.

To be satisfactory, any such specification must provide

- (a) Different specifications for all equations whose solutions involve different functions,
- (b) The same specification for all equations whose solutions are the same.

Since the Klein-Bôcher-Ince method does not satisfy these criteria [24], we shall employ a different method of specification.

## 6.01 BÔCHER EQUATIONS

It is convenient to consider a fairly general form of ordinary differential equation of the second order, called the *Bôcher equation*. All the foregoing separation equations can be written as Bôcher equations or can be obtained from Bôcher equations by transformation of independent or dependent variable. The Bôcher equation is

$$\frac{d^2Z}{dz^2} + P(z) \frac{dZ}{dz} + Q(z) Z = 0, \quad (6.01)$$

where

$$P(z) = \frac{1}{2} \left[ \frac{m_1}{z - a_1} + \frac{m_2}{z - a_2} + \cdots + \frac{m_{n-1}}{z - a_{n-1}} \right],$$

$$Q(z) = \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \cdots + \bar{A}_l z^l}{(z - a_1)^{m_1} (z - a_2)^{m_2} \cdots (z - a_{n-1})^{m_{n-1}}} \right],$$

and where  $m_i$ ,  $n$ , and  $l$  are non-negative integers.

Bôcher equations can be classified in terms of the singularities of  $P(z)$  and  $Q(z)$ . Evidently, Eq. (6.01) has no essential singularities, though both  $P(z)$  and

$Q(z)$  have poles at  $z = a_1, a_2, \dots, a_{n-1}$ . The *order of a pole* of the differential equation may be defined as the larger of the two integers representing the order of the pole of  $P$  and the order of the pole of  $Q$  at  $z = a_i$ . Usually the poles of  $Q$  are of higher (or equal) order compared with those of  $P$ . The integers  $m_i$  are then taken as the orders of the poles of the equation.

A special case occurs where  $(z - a_i)$  is a factor of the numerator of  $Q$ . This particular pole is then reduced in order for  $Q$ , though it remains unchanged for  $P$ . Another degenerate case occurs when the numerator of  $Q$  is zero. The poles of the equation are then fixed by  $P$ . Usually, however, the orders of the poles of the differential equation are  $m_1, m_2, \dots, m_{n-1}$ .

The above discussion applies to the singularities in the finite  $z$ -plane. Generally there is also a pole at infinity. Its order is obtained by taking the larger of the two integers representing the orders of

$$[2z - z^2 P(z)] \quad \text{and} \quad z^4 Q(z)$$

as  $z \rightarrow \infty$ .

## 6.02 SPECIFICATION

A Bôcher equation may be specified by writing a sequence of integers representing the order of the poles of the differential equation:

$$\{m_1, m_2, m_3, \dots, m_n\}.$$

The final integer  $m_n$  in this sequence holds a privileged position, since it refers to the pole at  $z \rightarrow \infty$ . The other integers indicate the orders of the poles in the finite  $z$ -plane, and they may be arranged in any convenient order [25].

The most complicated separation equation found in the preceding sections has four singularities ( $n = 4$ ) and is written

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{2}{z-a_2} + \frac{2}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2 + \bar{A}_3 z^3}{(z-a_1)(z-a_2)^2(z-a_3)^2} \right] Z = 0. \quad (6.02)$$

Evidently, there are three singularities in the finite plane: one first-order pole and two second-order poles. There is also a second-order pole at infinity. Thus the specification is  $\{1222\}$ .

A Bôcher equation with three singularities is

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{2}{z-a_1} + \frac{2}{z-a_2} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_2 z^2}{(z-a_1)^2(z-a_2)^2} \right] Z = 0. \quad (6.03)$$

If  $a_1 = 1$ ,  $a_2 = -1$ ,  $\bar{A}_0 = 4[p(p+1) - q^2]$ , and  $\bar{A}_2 = -4p(p+1)$ , then

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{2}{z-1} + \frac{2}{z+1} \right] \frac{dZ}{dz} + \left[ \frac{p(p+1) - q^2 - p(p+1)z^2}{(z-1)^2(z+1)^2} \right] Z = 0.$$

Second-order poles occur at  $z = \pm 1$ . At infinity,

$$[2z - z^2 P(z)] \quad \text{is of order 0,}$$

$$z^4 Q(z) \quad \text{is of order 2,}$$

so the pole at  $\infty$  is of order 2 and the equation is designated as {222}. It is usually written

$$(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} - \left[ p(p+1) + \frac{q^2}{z^2 - 1} \right] Z = 0 \quad (6.04)$$

which is *Legendre's equation*.

An example of a differential equation with two singularities is

$$\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{2}{z - a_1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_2 z^2}{(z - a_1)^2} \right] Z = 0. \quad (6.05)$$

Evidently the specification is {24}. If  $\bar{A}_0 = -4p^2$  and  $\bar{A}_2 = 4q^2$ ,

$$\frac{d^2 Z}{dz^2} + \frac{1}{z - a_1} \frac{dZ}{dz} + \left[ \frac{q^2 z^2 - p^2}{(z - a_1)^2} \right] Z = 0 \quad (6.05 \text{ a})$$

and if  $a_1 = 0$ , we have the familiar *Bessel equation*,

$$\frac{d^2 Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (q^2 - p^2/z^2) Z = 0. \quad (6.05 \text{ b})$$

Equations (6.05), (6.05 a), and (6.05 b) are all designated as {24}. But if  $p = 0$  and  $a_1 = 0$  in Eq. (6.05 a), the  $z$ 's cancel in  $Q$ , leaving

$$\frac{d^2 Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + q^2 Z = 0. \quad (6.06)$$

If the cancellation is not performed, the equation apparently remains a {24}; but as written in (6.06), it is {14}. The ordinary Bôcher {14} is

$$\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z - a_1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_2 z}{z - a_1} \right] Z = 0,$$

which differs from Eq. (6.06) by the  $\frac{1}{2}$  in the second term. To prevent ambiguity, we designate Eq. (6.06) as {14D}, indicating that it is not the ordinary Bôcher {14} but is a degenerate form of a higher type of Bôcher equation.

For an equation with one singularity, the specification must still contain two integers to distinguish between a pole at infinity and a pole in the finite  $z$ -plane. For instance, the simple differential equation

$$\frac{d^2 Z}{dz^2} = 0 \quad (6.07)$$

has no singularity in the finite plane and is specified as {01}. But the transformation  $z = \zeta^{-1}$  gives the differential equation

$$\frac{d^2 Z}{d\zeta^2} + \frac{2}{\zeta} \frac{dZ}{d\zeta} = 0,$$

which may be designated as {10}. Lists of differential equations and their specifications are given in Tables 6.01 and 6.02.

### 6.03 TRANSFORMATIONS

Some of the separation equations of mathematical physics do not appear in Bôcher form; but all of them can be changed into Bôcher equations by suitable transformations of the independent (or the dependent) variable. For  $n \geq 2$ , each

separation equation has a unique specification, obtained by examination of its Bôcher form. As an example, a separation equation in spherical coordinates is

$$\frac{d^2Z}{d\xi^2} + \cot \xi \frac{dZ}{d\xi} + p(p+1)Z = 0,$$

which is not in Bôcher form and which appears to have an infinite number of singularities. But the transformation  $z = \cos \xi$  gives

$$(z^2 - 1) \frac{d^2Z}{dz^2} + 2z \frac{dZ}{dz} - p(p+1)Z = 0.$$

Evidently both equations have the designation {112}.

The only cases where our method of designation leads to ambiguity are in the simplest examples [25], such as {01} and {04}. For instance, the separation equation

$$\frac{d^2Z}{d\xi^2} + \frac{1}{\xi} \frac{dZ}{d\xi} - \frac{p^2Z}{\xi} = 0$$

is a Bôcher equation and might be designated as {22}. But the transformation  $z = -i \ln \xi$  gives the elementary Bôcher equation

$$\frac{d^2Z}{dz^2} + p^2Z = 0,$$

whose designation is {04}. In these simple equations, it seems preferable to arbitrarily employ the same designation for all forms that can be obtained by functional transformations (see Table 6.02).

## 6.04 TABLES

Table 6.01 shows that, of all the differential equations obtained by separation of the Laplace and Helmholtz equations in 40 coordinate systems, there are only 9 distinct types, with some degenerate cases [26]. Table 6.02 gives canonical forms of the Bôcher equation for the separation equations needed in this work. Table 6.03 lists all the separation equations and shows how they are obtained from Table 6.02 by functional transformations. Also given are the general solutions of the differential equations. Details of the functions are treated in Section VII.

In only two cases is there an apparent violation of our uniqueness criterion:

(a) Solutions of {06} are listed as

$$Z = z^{\frac{1}{2}} [A \mathcal{J}_{\frac{1}{4}}(qz^2/2) + B \mathcal{I}_{-\frac{1}{4}}(qz^2/2)],$$

$$Z = A \mathcal{W}_e(p, qz) + B \mathcal{W}_o(p, qz),$$

$$Z = z^{\frac{1}{2}} [A \mathcal{J}_{\frac{1}{2}}(\varkappa, qz) + B \mathcal{I}_{-\frac{1}{2}}(\varkappa, qz)];$$

(b) Solutions of {113} are listed as Baer functions and as Mathieu functions.

It might seem that the specification of the differential equation should be different when it leads to a Bessel function  $\mathcal{J}_{\frac{1}{4}}$  and when it leads to a Weber function  $\mathcal{W}$ . It can be shown, however, that the above Bessel functions are special cases of the Weber functions. Similarly, the Mathieu functions are obtainable from the Baer functions when the singularities of the latter are moved to  $a_1 = 0$ ,  $a_2 = 1$ . Thus our specifications {06} and {113} are unambiguous.

**TABLE 6.01. CLASSIFICATION OF BÖCHER EQUATIONS**

obtained by separation of Laplace and Helmholtz equations in 40 coordinate systems.

| No.<br>Singularities $n$ | Name  | Original                               | Degenerate cases   |
|--------------------------|---|--|--|
| 1                        | Elementary<br>Weber equation                              | $\{04\}$<br>$\{06\}$                   | $\{04\}$   |
| 2                        | Elementary<br>Bessel wave equation                        | $\{42\}$<br>$\{26\}$                   | $\{22D\}$<br>$\{14D\}, \{16D\}, \{24\}$                  |
| 3                        | Baer wave equation<br>Legendre wave equation              | $\{114\}$<br>$\{224\}$                 | $\{113\}$<br>$\{112D\}, \{114D\}, \{220\},$<br>$\{222\}$ |
| 4                        | Lamé wave equation<br>Wangerin equation<br>Heine equation | $\{1113\}$<br>$\{1122\}$<br>$\{1222\}$ | $\{1111\}, \{1112\}$                                     |

$D$  indicates a degenerate form in which a cancellation of  $z$  occurs in numerator and denominator of  $Q(z)$ .

**TABLE 6.02. CANONICAL EQUATIONS**

occurring in field theory.

| Basic form             | Degenerate<br>form | Equation   |
|------------------------|--------------------|--|
| <b>One singularity</b> |                    |  |
| $\{04\}$               |                    | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{m_1}{z - a_1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0}{(z - a_1)^{m_1}} \right] Z = 0,$ $m_1 = 0, \quad \bar{A}_0 = 4\dot{p}^2; \text{ or}$ $\frac{d^2Z}{dz^2} + \dot{p}^2 Z = 0.$  |
|                        | $\{04\}$           | $\frac{d^2Z}{dz^2} = 0, \quad \{04\} \text{ with } \dot{p} = 0.$   |
| $\{06\}$               |                    | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{m_1}{z - a_1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_2 z^2}{(z - a_1)^{m_1}} \right] Z = 0,$ with $m_1 = 0$ . If $\bar{A}_0 = 4M, \bar{A}_2 = 4N$ , the foregoing equation becomes $\frac{d^2Z}{dz^2} + [M + Nz^2] Z = 0.$ |

Table 6.02. Continuation

| Basic form               | Degenerate form | Equation   |
|--------------------------|-----------------|--|
| {06}                     |                 | If $M = 0$ and $N = q^2$ ,<br>$\frac{d^2Z}{dz^2} + q^2 z^2 Z = 0.$   |
| {06}                     |                 | If $M = q^2(p + \frac{1}{2})$ and $N = -q^4/4$ ,<br>$\frac{d^2Z}{dz^2} + [q^2(p + \frac{1}{2}) - q^4 z^2/4] Z = 0.$<br>(Weber equation)  |
| {06}                     |                 | If $M = q^2$ and $N = \kappa^2$ ,<br>$\frac{d^2Z}{dz^2} + [q^2 + \kappa^2 z^2] Z = 0.$   |
| <b>Two singularities</b> |                 |  |
| {42}                     |                 | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{4}{z - a_1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_2 z^2}{(z - a_1)^4} \right] Z = 0.$<br>If $a_1 = 0$ and $\bar{A}_2 = -4p(p + 1)$ , the above equation reduces to  |
|                          | {22D}           | $\frac{d^2Z}{dz^2} + \frac{2}{z} \frac{dZ}{dz} - \frac{p(p + 1)}{z^2} Z = 0.$  |
| {26}                     |                 | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{2}{z - a_1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_2 z^2 + \bar{A}_4 z^4}{(z - a_1)^2} \right] Z = 0.$<br>If $a_1 = 0$ , $\bar{A}_0 = -4p^2$ , $\bar{A}_2 = 4q^2$ , $\bar{A}_4 = 4\kappa^2$ ,<br>$\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 + q^2 - p^2/z^2) Z = 0.$<br>(Bessel wave equation) |
| {24}                     |                 | $\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (q^2 - p^2/z^2) Z = 0,$<br>{26} with $\kappa = 0$ .<br>(Bessel equation)  |
| {16D}                    |                 | $\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 + q^2) Z = 0.$<br>{26} with $p = 0$ .   |
| {14D}                    |                 | $\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + q^2 Z = 0,$<br>{26} with $p = 0$ and $\kappa = 0$ .   |

Table 6.02. Continuation

| Basic form                 | Degenerate form | Equation  |
|----------------------------|-----------------|---|
| <b>Three singularities</b> |                 |   |
| $\{114\}$                  |                 | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} \right] \frac{dZ}{dz}$ $+ \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z-a_1)(z-a_2)} \right] Z = 0.$ <p>If <math>\bar{A}_0 = -4q^2</math>, <math>\bar{A}_1 = -4p^2</math>, <math>\bar{A}_2 = 4\kappa^2</math>,</p> $(z-a_1)(z-a_2) \frac{d^2Z}{dz^2} + \frac{1}{2} [2z - (a_1+a_2)] \frac{dZ}{dz}$ $+ [\kappa^2 z^2 - p^2 z - q^2] Z = 0.$ <p>(Baer wave equation)</p>   |
| $\{113\}$                  |                 | <p>If <math>\kappa = 0</math>,</p> $(z-a_1)(z-a_2) \frac{d^2Z}{dz^2} + \frac{1}{2} [2z - (a_1+a_2)] \frac{dZ}{dz}$ $- [p^2 z + q^2] Z = 0.$ <p>(Baer equation)</p>  |
| $\{113\}$                  |                 | <p>In general, Bôcher <math>\{113\}</math> is</p> $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} \right] \frac{dZ}{dz}$ $+ \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z}{(z-a_1)(z-a_2)} \right] Z = 0.$ <p>If <math>a_1 = 0</math>, <math>a_2 = +1</math>, <math>\bar{A}_0 = -(2q + \lambda)</math>, <math>\bar{A}_1 = 4q</math>, then</p> $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{1}{z-1} \right] \frac{dZ}{dz}$ $+ \frac{1}{4} \left[ \frac{4qz - (2q + \lambda)}{z(z-1)} \right] Z = 0.$ |
| $\{113\}$                  |                 | <p>Substitution of <math>z = \cos^2 \xi</math> gives</p> $\frac{d^2Z}{d\xi^2} + (\lambda - 2q \cos 2\xi) Z = 0.$ <p>(Mathieu equation)</p>  |
| $\{224\}$                  |                 | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{2}{z-a_1} + \frac{2}{z-a_2} \right] \frac{dZ}{dz}$ $+ \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2 + \bar{A}_3 z^3 + \bar{A}_4 z^4}{(z-a_1)^2(z-a_2)^2} \right] Z = 0.$   |

Table 6.02. Continuation

| Basic form                | Degenerate form | Equation   |
|---------------------------|-----------------|--|
|                           |                 | If $a_1 = 1$ , $a_2 = -1$ , $\bar{A}_0 = 4[\kappa^2 a^2 + p(p+1) - q^2]$ ,<br>$\bar{A}_2 = -4[2\kappa^2 a^2 + p(p+1)]$ , $\bar{A}_1 = \bar{A}_3 = 0$ , $\bar{A}_4 = 4\kappa^2 a^2$ ,   |
|                           |                 | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz}$<br>$+ [\kappa^2 a^2(z^2 - 1) - p(p+1) - \frac{q^2}{z^2 - 1}] Z = 0.$   |
|                           |                 | (Legendre wave equation)   |
| {222}                     |                 | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz}$<br>$- [p(p+1) + \frac{q^2}{z^2 - 1}] Z = 0,$   |
|                           |                 | Legendre wave equation {224} with $\kappa = 0$ .   |
|                           |                 | (Legendre equation)  |
| {220}                     |                 | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} - \frac{q^2}{z^2 - 1} Z = 0,$   |
|                           |                 | {224} with $\kappa = 0$ and $p = 0$ .  |
| {114D}                    |                 | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz}$<br>$+ [\kappa^2 a^2(z^2 - 1) - p(p+1)] Z = 0,$   |
|                           |                 | {224} with $q = 0$ .   |
| {112D}                    |                 | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} - p(p+1) Z = 0,$  |
|                           |                 | {224} with $\kappa = 0$ and $q = 0$ .  |
| <b>Four singularities</b> |                 |  |
| {1113}                    |                 | $\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z - a_1} + \frac{1}{z - a_2} + \frac{1}{z - a_3} \right] \frac{dZ}{dz}$<br>$+ \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z - a_1)(z - a_2)(z - a_3)} \right] Z = 0.$ |
|                           |                 | If $a_1 = 0$ , $\bar{A}_0 = (a_2^2 + a_3^2)q$ , $\bar{A}_1 = -p(p+1)$ , $\bar{A}_2 = \kappa^2$ ,   |
|                           |                 | $\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{1}{z - a_2} + \frac{1}{z - a_3} \right] \frac{dZ}{dz}$<br>$+ \frac{1}{4} \left[ \frac{(a_2^2 + a_3^2)q - p(p+1)z + \kappa^2 z^2}{z(z - a_2)(z - a_3)} \right] Z = 0.$             |
|                           |                 | (Lamé wave equation)   |

Table 6.02. Continuation

| Basic form       | Degenerate form  | Equation   |
|------------------|------------------|--|
|                  | $\{1\ 1\ 1\ 1\}$ | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{1}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0}{(z-a_1)(z-a_2)(z-a_3)} \right] Z = 0.$   |
|                  | $\{1\ 1\ 1\ 2\}$ | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{1}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z}{(z-a_1)(z-a_2)(z-a_3)} \right] Z = 0.$ <p style="text-align: center;">(Lamé equation)</p>                                      |
| $\{1\ 1\ 2\ 2\}$ |                  | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{2}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z-a_1)(z-a_2)(z-a_3)^2} \right] Z = 0.$ <p style="text-align: center;">(Wangerin equation)</p>                |
| $\{1\ 2\ 2\ 2\}$ |                  | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{2}{z-a_2} + \frac{2}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2 + \bar{A}_3 z^3}{(z-a_1)(z-a_2)^2(z-a_3)^2} \right] Z = 0.$ <p style="text-align: center;">(Heine equation)</p> |

TABLE 6.03. THE SEPARATION EQUATIONS

of field theory, for Laplace and Helmholtz equations in 40 coordinate systems.

| Standard designation   | Transformation | Differential equation and solution  |
|------------------------|----------------|---|
| <b>One singularity</b> |                |   |
|                        |                | <b>Canonical form</b>   |
| $\{0\ 1\}$             |                | $\frac{d^2Z}{dz^2} = 0, \quad Z = A + Bz.$  |
| <b>Other forms</b>     |                |   |
|                        | $z = 1/\xi$    | $\frac{d^2Z}{d\xi^2} + \frac{2}{\xi} \frac{dZ}{d\xi} = 0, \quad Z = A + B/\xi.$     |
|                        | $z = \ln \xi$  | $\frac{d^2Z}{d\xi^2} + \frac{1}{\xi} \frac{dZ}{d\xi} = 0, \quad Z = A + B \ln \xi.$ |

Table 6.03. Continuation

| Standard designation | Transformation  | Differential equation and solution   |
|----------------------|---|--|
|                      | $z = \ln \cot(\zeta/2)$   | $\frac{d^2 Z}{d\zeta^2} + \cot \zeta \frac{dZ}{d\zeta} = 0,$<br>$Z = A + B \ln \cot(\zeta/2).$   |
|                      | $z = \ln \coth(\zeta/2)$  | $\frac{d^2 Z}{d\zeta^2} + \coth \zeta \frac{dZ}{d\zeta} = 0,$<br>$Z = A + B \ln \coth(\zeta/2).$   |
|                      | $z = \cot^{-1}(\sinh \zeta)$  | $\frac{d^2 Z}{d\zeta^2} + \tanh \zeta \frac{dZ}{d\zeta} = 0,$<br>$Z = A + B \cot^{-1}(\sinh \zeta).$   |
|                      | $z = \operatorname{sn}^{-1}\left(\frac{\zeta}{b}, \frac{b}{c}\right)$ | $\frac{d}{d\zeta} \left[ (b^2 - \zeta^2)^{\frac{1}{2}} (c^2 - \zeta^2)^{\frac{1}{2}} \frac{dZ}{d\zeta} \right] = 0$<br>or $\frac{d}{d\zeta} \left[ (\zeta^2 - b^2)^{\frac{1}{2}} (\zeta^2 - c^2)^{\frac{1}{2}} \frac{dZ}{d\zeta} \right] = 0$<br>or $\frac{d}{d\zeta} \left[ (b^2 - \zeta^2)^{\frac{1}{2}} (\zeta^2 - c^2)^{\frac{1}{2}} \frac{dZ}{d\zeta} \right] = 0,$<br>$Z = A + B \operatorname{sn}^{-1}\left(\frac{\zeta}{b}, \frac{b}{c}\right).$ |
|                      | $z = \sin^{-1}\left[\frac{2\zeta - (b+c)}{b-c}\right]$                | $\frac{d}{d\zeta} \left[ (b - \zeta)^{\frac{1}{2}} (c - \zeta)^{\frac{1}{2}} \frac{dZ}{d\zeta} \right] = 0$<br>or $\frac{d}{d\zeta} \left[ (\zeta - b)^{\frac{1}{2}} (\zeta - c)^{\frac{1}{2}} \frac{dZ}{d\zeta} \right] = 0$<br>or $\frac{d}{d\zeta} \left[ (b - \zeta)^{\frac{1}{2}} (\zeta - c)^{\frac{1}{2}} \frac{dZ}{d\zeta} \right] = 0,$<br>$Z = A + B \sin^{-1}\left[\frac{2\zeta - (b+c)}{b-c}\right].$  |
| {04}                 |   | $\frac{d^2 Z}{dz^2} + p^2 Z = 0,$<br>$Z = A \sin pz + B \cos pz.$  |
|                      | $z = i\zeta$  | $\frac{d^2 Z}{d\zeta^2} - p^2 Z = 0,$<br>$Z = A e^{pz} + B e^{-pz}.$   |
|                      | $z = -i \ln \zeta$  | $\frac{d^2 Z}{d\zeta^2} + \frac{1}{\zeta} \frac{dZ}{d\zeta} - \frac{p^2 Z}{\zeta^2} = 0,$<br>$Z = A \zeta^p + B \zeta^{-p}.$   |
|                      | $X = Z/z$   | $\frac{d^2 X}{dz^2} + \frac{2}{z} \frac{dX}{dz} + p^2 X = 0,$<br>$X = \frac{1}{z} [A \sin pz + B \cos pz].$  |

Table 6.03. Continuation

| Standard designation | Transformation | Differential equation and solution  |
|----------------------|----------------|---|
| {06}                 |                | $\frac{d^2 Z}{dz^2} + [M + Nz^2] Z = 0.$ <p>If <math>M = 0</math> and <math>N = q^2</math>,</p> $\frac{d^2 Z}{dz^2} + q^2 z^2 Z = 0,$ $Z = z^{\frac{1}{2}} [A \mathcal{J}_{\frac{1}{4}}(qz^2/2) + B \mathcal{I}_{-\frac{1}{4}}(qz^2/2)].$ |
| {06}                 |                | <p>If <math>M = q^2(p + \frac{1}{2})</math> and <math>N = -q^4/4</math>,</p> $\frac{d^2 Z}{dz^2} + [q^2(p + \frac{1}{2}) - q^4 z^2/4] Z = 0,$ $Z = A \mathcal{W}_e(p, qz) + B \mathcal{W}_o(p, qz).$ <p>(Weber equation)</p>              |
|                      | $z = i\zeta$   | $\frac{d^2 Z}{d\zeta^2} - [q^2(p + \frac{1}{2}) + q^4 \zeta^2/4] Z = 0,$ $Z = A \mathcal{W}_e(p, iq\zeta) + B \mathcal{W}_o(p, iq\zeta).$   |
| {06}                 |                | <p>If <math>M = q^2</math> and <math>N = \kappa^2</math>,</p> $\frac{d^2 Z}{dz^2} + [q^2 + \kappa^2 z^2] Z = 0,$ $Z = z^{\frac{1}{2}} [A \mathcal{J}_{\frac{1}{2}}(\kappa, qz) + B \mathcal{I}_{-\frac{1}{2}}(\kappa, qz)].$              |
|                      | $z = i\zeta$   | $\frac{d^2 Z}{d\zeta^2} - [q^2 - \kappa^2 \zeta^2] Z = 0,$ $Z = \sqrt{i\zeta} [A \mathcal{J}_{\frac{1}{2}}(\kappa, iq\zeta) + B \mathcal{I}_{-\frac{1}{2}}(\kappa, iq\zeta)].$  |
| Two singularities    |                |   |
| {22D}                |                | $\frac{d^2 Z}{dz^2} + \frac{2}{z} \frac{dZ}{dz} - \frac{p(p+1)}{z^2} Z = 0,$ $Z = A z^p + B z^{-(p+1)}.$  |
| {14D}                |                | $\frac{d^2 Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + q^2 Z = 0,$ $Z = A \mathcal{J}_0(qz) + B \mathcal{Y}_0(qz).$  |
|                      | $z = i\zeta$   | $\frac{d^2 Z}{d\zeta^2} + \frac{1}{\zeta} \frac{dZ}{d\zeta} - q^2 Z = 0,$ $Z = A \mathcal{J}_0(iq\zeta) + B \mathcal{Y}_0(iq\zeta).$  |
| {16D}                |                | $\frac{d^2 Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 + q^2) Z = 0,$ $Z = A \mathcal{J}_0(\kappa, qz) + B \mathcal{Y}_0(\kappa, qz).$   |

Table 6.03. Continuation

| Standard designation | Transformation                                    | Differential equation and solution   |
|----------------------|---|--|
|                      | $z = i\zeta$                                      | $\frac{d^2Z}{d\zeta^2} + \frac{1}{\zeta} \frac{dZ}{d\zeta} + (\kappa^2\zeta^2 - q^2)Z = 0,$ $Z = A\mathcal{J}_0(\kappa, iq\zeta) + B\mathcal{Y}_0(\kappa, iq\zeta).$   |
| {24}                 |   | $\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (q^2 - p^2/z^2)Z = 0,$ $Z = A\mathcal{J}_p(qz) + B\mathcal{J}_{-p}(qz)$ or $Z = A\mathcal{J}_n(qz) + B\mathcal{Y}_n(qz).$ <p>(Bessel equation)</p>  |
|                      | $z = i\zeta$                                      | $\frac{d^2Z}{d\zeta^2} + \frac{1}{\zeta} \frac{dZ}{d\zeta} - (q^2 + p^2/\zeta^2)Z = 0,$ $Z = A\mathcal{J}_p(iq\zeta) + B\mathcal{J}_{-p}(iq\zeta)$ or $Z = A\mathcal{J}_n(iq\zeta) + B\mathcal{Y}_n(iq\zeta).$   |
|                      | $Z = X z^{\frac{1}{2}},$<br>$s = p - \frac{1}{2}$ | $\frac{d^2X}{dz^2} + \frac{2}{z} \frac{dX}{dz} + \left[ q^2 - \frac{s(s+1)}{z^2} \right] X = 0,$ $X = z^{-\frac{1}{2}} [A\mathcal{J}_{s+\frac{1}{2}}(qz) + B\mathcal{J}_{-(s+\frac{1}{2})}(qz)].$  |
| {26}                 |   | $\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 + q^2 - p^2/z^2)Z = 0,$ $Z = A\mathcal{J}_p(\kappa, q, z) + B\mathcal{J}_{-p}(\kappa, q, z)$ or $Z = A\mathcal{J}_n(\kappa, q, z) + B\mathcal{Y}_n(\kappa, q, z).$ <p>(Bessel wave equation)</p>          |
|                      | $z = i\zeta$                                      | $\frac{d^2Z}{d\zeta^2} + \frac{1}{\zeta} \frac{dZ}{d\zeta} + (\kappa^2\zeta^2 - q^2 - p^2/\zeta^2)Z = 0,$ $Z = A\mathcal{J}_p(\kappa, q, i\zeta) + B\mathcal{J}_{-p}(\kappa, q, i\zeta)$ or $Z = A\mathcal{J}_n(\kappa, q, i\zeta) + B\mathcal{Y}_n(\kappa, q, i\zeta).$ |
| Three singularities  |   |  |
| {112D}               |   | $(z^2 - 1) \frac{d^2Z}{dz^2} + 2z \frac{dZ}{dz} - p(p+1)Z = 0,$ $Z = A\mathcal{P}_p(z) + B\mathcal{Q}_p(z).$   |
|                      | $z = \cos \zeta$                                  | $\frac{d^2Z}{d\zeta^2} + \cot \zeta \frac{dZ}{d\zeta} + p(p+1)Z = 0,$ $Z = A\mathcal{P}_p(\cos \zeta) + B\mathcal{Q}_p(\cos \zeta).$   |

Table 6.03. Continuation

| Standard designation | Transformation      | Differential equation and solution   |
|----------------------|---------------------|--|
|                      | $z = \cosh \zeta$   | $\frac{d^2Z}{d\zeta^2} + \coth \zeta \frac{dZ}{d\zeta} - p(p+1)Z = 0,$ $Z = A \mathcal{P}_p(\cosh \zeta) + B \mathcal{Q}_p(\cosh \zeta).$<br>$\frac{d^2Z}{d\zeta^2} + \coth \zeta \frac{dZ}{d\zeta} - (p^2 - \frac{1}{4})Z = 0,$ $Z = A \mathcal{P}_{p-\frac{1}{2}}(\cosh \zeta) + B \mathcal{Q}_{p-\frac{1}{2}}(\cosh \zeta).$<br>$\frac{d^2Z}{d\zeta^2} + \coth \zeta \frac{dZ}{d\zeta} + Z/4 = 0,$ $Z = A \mathcal{P}_{-\frac{1}{2}}(\cosh \zeta) + B \mathcal{Q}_{-\frac{1}{2}}(\cosh \zeta).$   |
|                      | $z = i \sinh \zeta$ | $\frac{d^2Z}{d\zeta^2} + \tanh \zeta \frac{dZ}{d\zeta} - p(p+1)Z = 0,$ $Z = A \mathcal{P}_p(i \sinh \zeta) + B \mathcal{Q}_p(i \sinh \zeta).$  |
| {113}                |                     | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{1}{z-1} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{4qz - (2q+\lambda)}{z(z-1)} \right] Z = 0.$<br>$\frac{d^2Z}{d\zeta^2} + (\lambda - 2q \cos 2\zeta) Z = 0,$ $Z = A \text{ce}_m(\zeta, q) + B \text{fe}_m(\zeta, q)$ <p style="text-align: center;">or</p> $Z = A \text{se}_m(\zeta, q) + B \text{ge}_m(\zeta, q).$<br>$\frac{d^2Z}{d\zeta^2} + (\lambda + 2q \cos 2\zeta) Z = 0,$ $Z = A \text{ce}_m(\zeta, -q) + B \text{fe}_m(\zeta, -q)$ <p style="text-align: center;">or</p> $Z = A \text{se}_m(\zeta, -q) + B \text{ge}_m(\zeta, -q).$<br>$\frac{d^2Z}{d\zeta^2} - (\lambda - 2q \cosh 2\zeta) Z = 0,$ $Z = A \text{ce}_m(i\zeta, q) + B \text{fe}_m(i\zeta, q)$ <p style="text-align: center;">or</p> $Z = A \text{se}_m(i\zeta, q) + B \text{ge}_m(i\zeta, q).$<br>$\frac{d^2Z}{d\zeta^2} - (\lambda + 2q \cosh 2\zeta) Z = 0,$ $Z = A \text{ce}_m(i\zeta, -q) + B \text{fe}_m(i\zeta, -q)$ <p style="text-align: center;">or</p> $Z = A \text{se}_m(i\zeta, -q) + B \text{ge}_m(i\zeta, -q).$ |

Table 6.03. Continuation

| Standard designation | Transformation | Differential equation and solution  |
|----------------------|----------------|---|
| {113}                |                | $(z - a_1)(z - a_2) \frac{d^2 Z}{dz^2} + \frac{1}{2}[2z - (a_1 + a_2)] \frac{dZ}{dz} - [p(p+1)z + q(a_1 + a_2)] Z = 0,$ $Z = A\mathcal{B}_p^q(z) + B\mathcal{C}_p^q(z).$ <p style="text-align: center;">(Baer equation)</p>   |
| {114}                |                | $(z - a_1)(z - a_2) \frac{d^2 Z}{dz^2} + \frac{1}{2}[2z - (a_1 + a_2)] \frac{dZ}{dz} + [\kappa^2 z^2 - p(p+1)z - q(a_1 + a_2)] Z = 0,$ $Z = A\mathcal{B}_p^q(\kappa z, z) + B\mathcal{C}_p^q(\kappa z, z).$ <p style="text-align: center;">(Baer wave equation)</p>   |
| {114D}               |                | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} + [\kappa^2 a^2(z^2 - 1) - p(p+1)] Z = 0,$ $Z = A\mathcal{P}_p(\kappa a, z) + B\mathcal{Q}_p(\kappa a, z).$<br>$(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} - [\kappa^2 a^2(z^2 - 1) + p(p+1)] Z = 0,$ $Z = A\mathcal{P}_p(i\kappa a, z) + B\mathcal{Q}_p(i\kappa a, z).$<br>$\frac{d^2 Z}{d\zeta^2} + \cot \zeta \frac{dZ}{d\zeta} + [\kappa^2 a^2 \sin^2 \zeta + p(p+1)] Z = 0,$ $Z = A\mathcal{P}_p(\kappa a, \cos \zeta) + B\mathcal{Q}_p(\kappa a, \cos \zeta).$<br>$\frac{d^2 Z}{d\zeta^2} + \cot \zeta \frac{dZ}{d\zeta} + [-\kappa^2 a^2 \sin^2 \zeta + p(p+1)] Z = 0,$ $Z = A\mathcal{P}_p(i\kappa a, \cos \zeta) + B\mathcal{Q}_p(i\kappa a, \cos \zeta).$<br>$\frac{d^2 Z}{d\zeta^2} + \coth \zeta \frac{dZ}{d\zeta} + [\kappa^2 a^2 \sinh^2 \zeta - p(p+1)] Z = 0,$ $Z = A\mathcal{P}_p(\kappa a, \cosh \zeta) + B\mathcal{Q}_p(\kappa a, \cosh \zeta).$ |

Table 6.03. Continuation

| Standard designation | Transformation    | Differential equation and solution  |
|----------------------|-------------------|---|
|                      | $z = i \sinh \xi$ | $\frac{d^2 Z}{d\xi^2} + \tanh \xi \frac{dZ}{d\xi}$ $+ [x^2 a^2 \cosh^2 \xi - p(p+1)] Z = 0,$ $Z = A \mathcal{P}_p(i \alpha a, i \sinh \xi) + B \mathcal{Q}_p(i \alpha a, i \sinh \xi).$                                   |
| {220}                |                   | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} - \left( \frac{q^2}{z^2 - 1} \right) Z = 0,$ $Z = A \mathcal{P}_0^q(z) + B \mathcal{Q}_0^q(z).$  |
|                      | $z = \cos \xi$    | $\frac{d^2 Z}{d\xi^2} + \cot \xi \frac{dZ}{d\xi} - \left( \frac{q^2}{\sin^2 \xi} \right) Z = 0,$ $Z = A \mathcal{P}_0^q(\cos \xi) + B \mathcal{Q}_0^q(\cos \xi).$   |
| {222}                |                   | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz}$ $- [p(p+1) + \frac{q^2}{z^2 - 1}] Z = 0,$ $Z = A \mathcal{P}_p^q(z) + B \mathcal{Q}_p^q(z).$ <p>(Legendre equation)</p>   |
|                      | $z = \cos \xi$    | $\frac{d^2 Z}{d\xi^2} + \cot \xi \frac{dZ}{d\xi}$ $+ \left[ p(p+1) - \frac{q^2}{\sin^2 \xi} \right] Z = 0,$ $Z = A \mathcal{P}_p^q(\cos \xi) + B \mathcal{Q}_p^q(\cos \xi).$  |
|                      | $z = \cosh \xi$   | $\frac{d^2 Z}{d\xi^2} + \coth \xi \frac{dZ}{d\xi}$ $- \left[ p(p+1) + \frac{q^2}{\sinh^2 \xi} \right] Z = 0,$ $Z = A \mathcal{P}_p^q(\cosh \xi) + B \mathcal{Q}_p^q(\cosh \xi).$  |
|                      |                   | $\frac{d^2 Z}{d\xi^2} + \coth \xi \frac{dZ}{d\xi}$ $- \left[ (p^2 - \frac{1}{4}) + \frac{q^2}{\sinh^2 \xi} \right] Z = 0,$ $Z = A \mathcal{P}_{p-\frac{1}{2}}^q(\cosh \xi) + B \mathcal{Q}_{p-\frac{1}{2}}^q(\cosh \xi).$ |
|                      | $z = i \sinh \xi$ | $\frac{d^2 Z}{d\xi^2} + \tanh \xi \frac{dZ}{d\xi}$ $+ \left[ -p(p+1) + \frac{q^2}{\cosh^2 \xi} \right] Z = 0,$ $Z = A \mathcal{P}_p^q(i \sinh \xi) + B \mathcal{Q}_p^q(i \sinh \xi).$                                     |

Table 6.03. Continuation

| Standard designation | Transformation    | Differential equation and solution  |
|----------------------|-------------------|---|
| {224}                |                   | $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} + [\kappa^2 a^2 (z^2 - 1) - p(p+1) - \frac{q^2}{z^2 - 1}] Z = 0.$ $Z = A \mathcal{P}_p^q(\kappa a, z) + B \mathcal{Q}_p^q(\kappa a, z).$ <p>(Legendre wave equation)</p> $(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} - [\kappa^2 a^2 (z^2 - 1) + p(p+1) + \frac{q^2}{z^2 - 1}] Z = 0,$ $Z = A \mathcal{P}_p^q(i\kappa a, z) + B \mathcal{Q}_p^q(i\kappa a, z).$ |
|                      | $z = \cos \xi$    | $\frac{d^2 Z}{d\xi^2} + \cot \xi \frac{dZ}{d\xi} + [\kappa^2 a^2 \sin^2 \xi + p(p+1) - \frac{q^2}{\sin^2 \xi}] Z = 0,$ $Z = A \mathcal{P}_p^q(\kappa a, \cos \xi) + B \mathcal{Q}_p^q(\kappa a, \cos \xi).$   |
|                      | $z = \cos \xi$    | $\frac{d^2 Z}{d\xi^2} + \cot \xi \frac{dZ}{d\xi} + [-\kappa^2 a^2 \sin^2 \xi + p(p+1) - \frac{q^2}{\sin^2 \xi}] Z = 0,$ $Z = A \mathcal{P}_p^q(i\kappa a, \cos \xi) + B \mathcal{Q}_p^q(i\kappa a, \cos \xi).$  |
|                      | $z = \cosh \xi$   | $\frac{d^2 Z}{d\xi^2} + \coth \xi \frac{dZ}{d\xi} + [\kappa^2 a^2 \sinh^2 \xi - p(p+1) - \frac{q^2}{\sinh^2 \xi}] Z = 0,$ $Z = A \mathcal{P}_p^q(\kappa a, \cosh \xi) + B \mathcal{Q}_p^q(\kappa a, \cosh \xi).$  |
|                      | $z = i \sinh \xi$ | $\frac{d^2 Z}{d\xi^2} + \tanh \xi \frac{dZ}{d\xi} + [\kappa^2 a^2 \cosh^2 \xi - p(p+1) + \frac{q^2}{\cosh^2 \xi}] Z = 0,$ $Z = A \mathcal{P}_p^q(i\kappa a, i \sinh \xi) + B \mathcal{Q}_p^q(i\kappa a, i \sinh \xi).$  |
| Four singularities   |                   |   |
| {1111}               |                   | $\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{1}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0}{(z-a_1)(z-a_2)(z-a_3)} \right] Z = 0.$ <p>If <math>a_1 = 0</math> and <math>\bar{A}_0 = (a_2^2 + a_3^2) q</math>,</p> $Z = A \mathcal{E}_0^q(z^{\frac{1}{2}}) + B \mathcal{F}_0^q(z^{\frac{1}{2}}).$   |

Table 6.03. Continuation

| Standard designation | Transformation | Differential equation and solution   |
|----------------------|----------------|--|
|                      | $z = \zeta^2$  | $\begin{aligned} & (\zeta^2 - a_2^2)(\zeta^2 - a_3^2) \frac{d^2 Z}{d\zeta^2} + \zeta [2\zeta^2 - (a_2^2 + a_3^2)] \\ & \times \frac{dZ}{d\zeta} + q^2(a_2^2 + a_3^2) Z = 0, \\ & Z = A \mathcal{E}_0^q(\zeta) + B \mathcal{F}_0^q(\zeta). \end{aligned}$   |
| {1112}               |                | $\begin{aligned} & \frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{1}{z-a_3} \right] \frac{dZ}{dz} \\ & + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z}{(z-a_1)(z-a_2)(z-a_3)} \right] Z = 0. \end{aligned}$ <p>If <math>a_1 = 0</math>, <math>\bar{A}_0 = (a_2^2 + a_3^2) q</math>, and<br/> <math>\bar{A}_1 = -p(p+1)</math>,</p> $Z = A \mathcal{E}_p^q(z^{\frac{1}{2}}) + B \mathcal{F}_p^q(z^{\frac{1}{2}}).$   |
|                      | $z = \zeta^2$  | $\begin{aligned} & (\zeta^2 - a_2^2)(\zeta^2 - a_3^2) \frac{d^2 Z}{d\zeta^2} \\ & + \zeta [2\zeta^2 - (a_2^2 + a_3^2)] \frac{dZ}{d\zeta} \\ & + [(a_2^2 + a_3^2)q - p(p+1)\zeta^2] Z = 0, \\ & Z = A \mathcal{E}_p^q(\zeta) + B \mathcal{F}_p^q(\zeta). \end{aligned}$ <p>(Lamé equation)</p>  |
| {1113}               |                | $\begin{aligned} & \frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{1}{z-a_3} \right] \frac{dZ}{dz} \\ & + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z-a_1)(z-a_2)(z-a_3)} \right] Z = 0. \end{aligned}$ <p>If <math>a_1 = 0</math>, <math>\bar{A}_0 = (a_2^2 + a_3^2)q</math>, <math>\bar{A}_1 = -p(p+1)</math>,<br/> and <math>\bar{A}_2 = \kappa^2</math>,</p> $Z = A \mathcal{E}_p^q(\kappa, z^{\frac{1}{2}}) + B \mathcal{F}_p^q(\kappa, z^{\frac{1}{2}}).$ <p>(Lamé wave equation)</p> |
|                      | $z = \zeta^2$  | $\begin{aligned} & (\zeta^2 - a_2^2)(\zeta^2 - a_3^2) \frac{d^2 Z}{d\zeta^2} \\ & + \zeta [2\zeta^2 - (a_2^2 + a_3^2)] \frac{dZ}{d\zeta} + [\kappa^2 \zeta^4 \\ & - p(p+1)\zeta^2 + (a_2^2 + a_3^2)q] Z = 0, \\ & Z = A \mathcal{E}_p^q(\kappa, \zeta) + B \mathcal{F}_p^q(\kappa, \zeta). \end{aligned}$  |

Table 6.03. Continuation

| Standard designation | Transformation | Differential equation and solution  |
|----------------------|----------------|---|
| $\{1122\}$           |                | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{1}{z-a_2} + \frac{2}{z-a_3} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{(z-a_1)(z-a_2)(z-a_3)^2} \right] Z = 0.$ <p>If <math>a_1 = 1, a_2 = 1/k^2, a_3 = 0, \bar{A}_0 = -\alpha_3/k^2, \bar{A}_1 = -\alpha_2/k^2, \bar{A}_2 = 1 - \alpha_3,</math></p> $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-1} + \frac{1}{z-1/k^2} + \frac{2}{z} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{-\alpha_3/k^2 - \alpha_2 z/k^2 + (1 - \alpha_3) z^2}{z^2(z-1)(z-1/k^2)} \right] \times Z = 0,$ $Z = A \mathcal{S}_p^q(k, z) + B \mathcal{T}_p^q(k, z).$ <p>(Wangerin equation)</p> $z = \operatorname{sn}^2 \zeta$ $\frac{d^2Z}{d\zeta^2} + \frac{\operatorname{cn} \zeta \operatorname{dn} \zeta}{\operatorname{sn} \zeta} \frac{dZ}{d\zeta} + \left[ k^2 \operatorname{sn}^2 \zeta - \alpha_2 - \alpha_3 \left( k^2 \operatorname{sn}^2 \zeta + \frac{1}{\operatorname{sn}^2 \zeta} \right) \right] Z = 0.$ $Z = A \mathcal{S}_p^q(k, \operatorname{sn}^2 \zeta) + B \mathcal{T}_p^q(k, \operatorname{sn}^2 \zeta).$ $z = \operatorname{cn}^2 \zeta$ $\frac{d^2Z}{d\zeta^2} - \frac{\operatorname{sn} \zeta \operatorname{dn} \zeta}{\operatorname{cn} \zeta} \frac{dZ}{d\zeta} + \left[ k^2 \operatorname{sn}^2 \zeta - \alpha_2 + \alpha_3 \left( k^2 \operatorname{cn}^2 \zeta - \frac{k'^2}{\operatorname{cn}^2 \zeta} \right) \right] Z = 0,$ <p>If <math>a_1 = 1, a_2 = -(k'/k)^2, a_3 = 0, \bar{A}_0 = (k'/k)^2, \bar{A}_1 = (\alpha_2 - k^2)/k^2, \bar{A}_2 = \alpha_3 - 1,</math></p> $Z = A \mathcal{S}_p^q(k, \operatorname{cn}^2 \zeta) + B \mathcal{T}_p^q(k, \operatorname{cn}^2 \zeta).$ $z = \operatorname{cn}^2 \zeta$ $\frac{d^2Z}{d\zeta^2} - \frac{\operatorname{sn} \zeta \operatorname{dn} \zeta}{\operatorname{cn} \zeta} \frac{dZ}{d\zeta} + \left[ -\operatorname{dn}^2 \zeta + \alpha_2 + \alpha_3 \left( k'^2 \operatorname{cn}^2 \zeta - \frac{k^2}{\operatorname{cn}^2 \zeta} \right) \right] Z = 0,$ <p>If <math>a_1 = 1, a_2 = -(k'/k)^2, a_3 = 0, \bar{A}_0 = \alpha_3, \bar{A}_1 = (k'^2 - \alpha_2)/k^2, \bar{A}_2 = 1 - \alpha_3 (k'/k)^2,</math></p> $Z = A \mathcal{S}_p^q(k, \operatorname{cn}^2 \zeta) + B \mathcal{T}_p^q(k, \operatorname{cn}^2 \zeta).$ |

Table 6.03. Continuation

| Standard designation | Transformation                  | Differential equation and solution   |
|----------------------|---------------------------------|--|
|                      | $z = \operatorname{dn}^2 \zeta$ | $\frac{d^2 Z}{d\zeta^2} - \frac{k'^2 \operatorname{sn} \zeta \operatorname{cn} \zeta}{\operatorname{dn} \zeta} \frac{dZ}{d\zeta} + \left[ -\operatorname{dn}^2 \zeta + \alpha_2 + \alpha_3 \left( \operatorname{dn}^2 \zeta + \frac{k^2}{\operatorname{dn}^2 \zeta} \right) \right] Z = 0,$ <p>If <math>\alpha_1 = 1, \alpha_2 = k^2, \alpha_3 = 0, \bar{A}_0 = -\alpha_3 k^2, \bar{A}_1 = -\alpha_2, \bar{A}_3 = 1 - \alpha_3</math> in the original {1122} equation</p> $Z = A \mathcal{U}_p^q(k, \operatorname{dn}^2 \zeta) + B \mathcal{T}_p^q(k, \operatorname{dn}^2 \zeta).$   |
| {1222}               |                                 | $\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-a_1} + \frac{2}{z-a_2} + \frac{2}{z-a_3} \right] \frac{dZ}{dz}$ $+ \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2 + \bar{A}_3 z^3}{(z-a_1)(z-a_2)^2(z-a_3)^2} \right]$ $\times Z = 0,$ $Z = A \mathcal{U}_p^q(k, z) + B \mathcal{V}_p^q(k, z).$ <p>(Heine equation)</p> <p>Let <math>a_1 = 0, a_2 = 1, a_3 = 1/k^2, \bar{A}_0 = -\alpha_2/k^2, \bar{A}_1 = (\alpha_2 + 2) + \alpha_2/k^2 - \alpha_3 k'^4/k^2, \bar{A}_2 = (\alpha_2 + 2) + 2k^2, \bar{A}_3 = 2k^4.</math></p>  |
|                      | $z = \operatorname{sn}^2 \zeta$ | $\frac{d^2 Z}{d\zeta^2} - \frac{\operatorname{sn} \zeta (\operatorname{dn}^2 \zeta + k^2 \operatorname{cn}^2 \zeta)}{\operatorname{cn} \zeta \operatorname{dn} \zeta} \frac{dZ}{d\zeta}$ $+ \left[ 2k^2 \operatorname{sn}^2 \zeta - \alpha_2 - \alpha_3 \frac{k'^4 \operatorname{sn}^2 \zeta}{\operatorname{cn}^2 \zeta \operatorname{dn}^2 \zeta} \right]$ $\times Z = 0,$ $Z = A \mathcal{U}_p^q(k, \operatorname{sn}^2 \zeta) + B \mathcal{V}_p^q(k, \operatorname{sn}^2 \zeta).$ <p>Let <math>a_1 = 0, a_2 = 1, a_3 = k^2, \bar{A}_0 = -\alpha_2 k^2, \bar{A}_1 = (\alpha_2 - \alpha_3) + k^2(\alpha_2 + 2), \bar{A}_2 = -(\alpha_2 + 2) - 2k^2, \bar{A}_3 = 2.</math></p> |
|                      | $z = \operatorname{dn}^2 \zeta$ | $\frac{d^2 Z}{d\zeta^2} + \frac{\operatorname{dn} \zeta (\operatorname{cn}^2 \zeta - \operatorname{sn}^2 \zeta)}{\operatorname{sn} \zeta \operatorname{cn} \zeta} \frac{dZ}{d\zeta}$ $+ \left[ -2 \operatorname{dn}^2 \zeta + \alpha_2 + \alpha_3 \frac{\operatorname{dn}^2 \zeta}{\operatorname{sn}^2 \zeta \operatorname{cn}^2 \zeta} \right]$ $\times Z = 0,$ $Z = A \mathcal{U}_p^q(k, \operatorname{dn}^2 \zeta) + B \mathcal{V}_p^q(k, \operatorname{dn}^2 \zeta).$  |

## Section VII

### FUNCTIONS

The purpose of this section is to provide a summary of the mathematical functions obtained as solutions of the differential equations of field theory. The summary is necessarily incomplete, since many of the functions have never been thoroughly investigated. In particular, almost nothing is known about the properties of the various *wave functions*. Recent tabulation of spheroidal wave functions is a beginning in this direction, but much remains to be done.

Even with the functions that have been studied, the question of *notation* is a troublesome one. With Mathieu functions, for instance, there are almost as many notations as there are investigators. We have tried in such cases to adopt a notation that is modern and logical.

In particular, there seems to be no advantage in employing different symbols for the same function of real and imaginary arguments. Bessel functions of the first kind, for example, are here denoted by  $\mathcal{J}_p(z)$  for the complex argument  $z$ , and are written  $\mathcal{J}_p(iy)$ , not  $I_p(y)$ , for imaginary arguments. Similarly, Mathieu functions of the first kind are written  $ce_m(q, iy)$ ,  $se_m(q, iy)$  rather than  $Ce_m(q, y)$ ,  $Se_m(q, y)$ . Where an additional parameter enters, as in transition from Laplace to Helmholtz equations, it seems best to retain the basic letter for the function, as when  $\mathcal{J}_p(z)$  changes to  $\mathcal{J}_p(\kappa, z)$ .

#### 7.01 FUNCTIONS

Many of the differential equations of Section VI are satisfied by elementary functions. The properties of these functions are well-known and will not be considered here. But other differential equations of field theory require Bessel, Legendre, and other functions for their solution. We shall attempt to tabulate useful information on these functions. For additional data, see the references listed in the Bibliography, Section VIII. The Bibliography also refers to numerical tables of the various functions, in so far as such tables are known to the authors.

The functions considered are as follows:

| Differential equation  | Functions  | Differential equation   | Functions  |
|------------------------|--|-------------------------|--|
| <b>Weber functions</b> |  | <b>Bessel functions</b> |  |
| $\{06\}$               | $\mathcal{W}_e(p, qz), \quad \mathcal{W}_e(p, iqz),$<br>$\mathcal{W}_o(p, qz), \quad \mathcal{W}_o(p, iqz).$ | $\{06\}$                | $\mathcal{J}_{\frac{1}{2}}(\kappa, q, z), \quad \mathcal{J}_{\frac{1}{2}}(\kappa, q, iz),$<br>$\mathcal{J}_{-\frac{1}{2}}(\kappa, q, z), \quad \mathcal{J}_{-\frac{1}{2}}(\kappa, q, iz).$ |

| Differential equation | Functions  | Differential equation     | Functions  |
|-----------------------|--|---------------------------|--|
| {06}                  | $\mathcal{J}_\frac{1}{4}(qz)$ , $\mathcal{J}_{-\frac{1}{4}}(qz)$ .         | {114}                     | $\mathcal{P}_p(\kappa a, z)$ , $\mathcal{P}_p(i\kappa a, z)$ ,     |
| {14}                  | $\mathcal{J}_0(qz)$ , $\mathcal{J}_0(iqz)$ ,                               |                           | $\mathcal{Q}_p(\kappa a, z)$ , $\mathcal{Q}_p(i\kappa a, z)$ ,     |
|                       | $\mathcal{Y}_0(qz)$ , $\mathcal{Y}_0(iqz)$ .                               |                           | $\mathcal{P}_p(i\kappa a, iz)$ ,                                   |
| {16}                  | $\mathcal{J}_0(\kappa, q, z)$ , $\mathcal{J}_0(\kappa, q, iz)$ ,           | {220}                     | $\mathcal{P}_0^q(z)$ ,   |
|                       | $\mathcal{Y}_0(\kappa, q, z)$ , $\mathcal{Y}_0(\kappa, q, iz)$ .           |                           | $\mathcal{Q}_0^q(z)$ .   |
| {24}                  | $\mathcal{J}_p(qz)$ , $\mathcal{J}_p(iqz)$ ,                               | {222}                     | $\mathcal{P}_p^q(z)$ , $P_n^m(z)$ ,                                |
|                       | $\mathcal{J}_{-p}(qz)$ , $\mathcal{J}_{-p}(iqz)$ .                         |                           | $\mathcal{Q}_p^q(z)$ , $Q_n^m(z)$ ,                                |
|                       | $\mathcal{Y}_n(qz)$ , $\mathcal{Y}_n(iqz)$ .                               |                           | $\mathcal{P}_{p-\frac{1}{2}}^q(z)$ ,                               |
| {24}                  | $\mathcal{J}_{s+\frac{1}{2}}(qz)$ , $\mathcal{J}_{-(s+\frac{1}{2})}(qz)$ . |                           | $\mathcal{Q}_{p-\frac{1}{2}}^q(z)$ .                               |
| {26}                  | $\mathcal{J}_p(\kappa, q, z)$ , $\mathcal{J}_p(\kappa, q, iz)$ ,           | {224}                     | $\mathcal{P}_p^q(\kappa a, z)$ , $\mathcal{P}_p^q(i\kappa a, z)$ , |
|                       | $\mathcal{J}_{-p}(\kappa, q, z)$ , $\mathcal{J}_{-p}(\kappa, q, iz)$ ,     |                           | $\mathcal{Q}_p^q(\kappa a, z)$ , $\mathcal{Q}_p^q(i\kappa a, z)$ , |
|                       | $\mathcal{Y}_n(\kappa, q, z)$ , $\mathcal{Y}_n(\kappa, q, iz)$ .           |                           | $\mathcal{P}_p^q(i\kappa a, iz)$ ,                                 |
|                       | <b>Baer functions</b>  |                           |  |
| {113}                 | $\mathcal{B}_p^q(z)$ ,   | <b>Lamé functions</b>     |  |
|                       | $\mathcal{C}_p^q(z)$ ,   |                           |  |
| {114}                 | $\mathcal{B}_0^q(\kappa, z)$ ,   | {1111}                    | $E_0^q(z)$ ,   |
|                       | $\mathcal{C}_0^q(\kappa, z)$ .   |                           | $\mathcal{F}_0^q(z)$ ,   |
|                       | <b>Mathieu functions</b>   |                           |  |
| {113}                 | $ce_m(q, z)$ , $ce_m(q, iz)$ ,   | {1112}                    | $E_n^q(z)$ ,   |
|                       | $fe_m(q, z)$ , $fe_m(q, iz)$ ,   |                           | $\mathcal{F}_n^q(z)$ ,   |
|                       | $se_m(q, z)$ , $se_m(q, iz)$ ,   | {1113}                    | (Polynomials.)   |
|                       | $ge_m(q, z)$ , $ge_m(q, iz)$ .   |                           | Applicable in<br>only a few<br>special cases.)                     |
|                       | <b>Legendre functions</b>  |                           |  |
| {112}                 | $\mathcal{P}_p(z)$ , $P_n(z)$ ,  | <b>Wangerin functions</b> |  |
|                       | (Polynomials)  |                           |  |
|                       | $\mathcal{Q}_p(z)$ , $Q_n(z)$ ,  | {1122}                    | $\mathcal{S}_p^q(k, z)$ ,  |
|                       | $\mathcal{P}_{p-\frac{1}{2}}(z)$ ,   |                           | $\mathcal{T}_p^q(k, z)$ .  |
|                       | $\mathcal{Q}_{p-\frac{1}{2}}(z)$ ,   | <b>Heine functions</b>    |  |
|                       | $\mathcal{P}_{-\frac{1}{2}}(z)$ ,  | {1222}                    | $\mathcal{U}_p^q(k, z)$ ,  |
|                       | $\mathcal{Q}_{-\frac{1}{2}}(z)$ .  |                           | $\mathcal{V}_p^q(k, z)$ .  |

## 7.02 SERIES SOLUTIONS

We are interested in the solution of equations of the Bôcher type:

$$\frac{d^2Z}{dz^2} + P(z) \frac{dZ}{dz} + Q(z) Z = 0, \quad (7.01)$$

where  $P(z)$  and  $Q(z)$  are given in §6.01. The simple equations are usually satisfied by elementary functions; the more complicated equations are handled by use of infinite series.

Suppose that the solution is to be obtained about an *ordinary point*,  $z = z_0$ , where  $P(z)$  and  $Q(z)$  are analytic. Assume a solution of the form

$$Z = \sum_{n=0}^{\infty} C_n (z - z_0)^n. \quad (7.02)$$

This series is substituted into the differential equation, and the coefficients  $C_n$  are evaluated. If  $P(z)$  and  $Q(z)$  are polynomials in  $(z - z_0)$ , they fit nicely into the assumed power series. If  $P$  and  $Q$  are other functions, they are analytic at  $z_0$  and can be expanded in power series in  $(z - z_0)$ . In either case, the solution of Eq. (7.01) is obtained by equating the coefficient of each power of  $(z - z_0)$  to zero. The series solution converges within the circle whose center is at  $z_0$  and whose radius extends to the nearest singularity of the differential equation.

If the equation has poles in the finite  $z$ -plane, one may find it convenient to expand about a *singular point*  $z_0$ . Assume a solution

$$Z = (z - z_0)^\beta \sum_{j=0}^{\infty} C_j (z - z_0)^j. \quad (7.03)$$

Equation (7.01) may be written

$$(z - z_0)^2 \frac{d^2Z}{dz^2} + (z - z_0) [(z - z_0) P(z)] \frac{dZ}{dz} + [(z - z_0)^2 Q(z)] Z = 0. \quad (7.01\text{a})$$

If  $[(z - z_0) P(z)]$  and  $[(z - z_0)^2 Q(z)]$  are analytic at  $z_0$ , this singularity is said to be a *regular singularity*. Then

$$\left. \begin{aligned} [(z - z_0) P(z)] &= \sum_{i=0}^{\infty} A_i (z - z_0)^i, \\ [(z - z_0)^2 Q(z)] &= \sum_{i=0}^{\infty} B_i (z - z_0)^i. \end{aligned} \right\} \quad (7.04)$$

Substitution of Eqs. (7.03) and (7.04) into (7.01a) gives

$$\left. \begin{aligned} C_0 f_0(\beta) &= 0, \\ C_1 f_0(\beta + 1) + C_0 f_1(\beta) &= 0, \\ C_2 f_0(\beta + 2) + C_1 f_1(\beta + 1) + C_0 f_2(\beta) &= 0, \\ \dots &\dots \\ C_j f_0(\beta + j) + C_{j-1} f_1(\beta + j - 1) + C_{j-2} f_2(\beta + j - 2) + \dots \\ &\quad + C_2 f_{j-2}(\beta + 2) + C_1 f_{j-1}(\beta + 1) + C_0 f_j(\beta) &= 0, \\ \dots &\dots \end{aligned} \right\} \quad (7.05)$$

where

$$\left. \begin{aligned} f_0(\beta) &= \beta^2 + (A_0 - 1)\beta + B_0, \\ f_j(\beta) &= \beta A_j + B_j, \quad j = 1, 2, 3, \dots \end{aligned} \right\} \quad (7.06)$$

For a non-trivial solution, we have the *indicial equation*,

$$\beta^2 + (A_0 - 1)\beta + B_0 = 0. \quad (7.07)$$

Take the roots of the indicial equation as  $b_1$  and  $b_2$ , with  $b_1 > b_2$ . Then

$$f_0(\beta) = (\beta - b_1)(\beta - b_2). \quad (7.08)$$

The case of equal roots will be considered later (Section 7.03).

A solution of Eq. (7.01) requires first the evaluation of the roots  $b_1$  and  $b_2$ . The  $f$ 's are then obtained from Eq. (7.06), and the coefficients  $C_n$  are found from Eq. (7.05). If the roots are distinct and do not differ by an integer, this procedure yields the two series required for the general solution of the second-order differential equation. Eq. (7.05) expresses each coefficient in terms of the preceding ones. Or any coefficient may be obtained directly by use of the determinant  $\Delta_j(\beta)$ :

$$C_j(\beta) = \frac{(-1)^j C_0 \Delta_j(\beta)}{f_0(\beta+1) \cdot f_0(\beta+2) \cdots f_0(\beta+j)}, \quad (7.09)$$

where  $\Delta_0 = 1$  and, for  $j \geq 1$ ,

$$\Delta_j(\beta) = \begin{vmatrix} f_1(\beta) & f_0(\beta+1) & 0 & 0 & 0 & \dots & 0 \\ f_2(\beta) & f_1(\beta+1) & f_0(\beta+2) & 0 & 0 & \dots & 0 \\ f_3(\beta) & f_2(\beta+1) & f_1(\beta+2) & f_0(\beta+3) & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ f_{j-1}(\beta) & f_{j-2}(\beta+1) & f_{j-3}(\beta+2) & \dots & f_1(\beta+j-2) & f_0(\beta+j-1) \\ f_j(\beta) & \dots & \dots & \dots & f_2(\beta+j-2) & f_1(\beta+j-1) \end{vmatrix}.$$

According to Eq. (7.08), the denominator of Eq. (7.09) may be written

$$[(\beta - b_1 + 1)(\beta - b_2 + 1)][(\beta - b_1 + 2)(\beta - b_2 + 2)] \cdots [(\beta - b_1 + j)(\beta - b_2 + j)] \\ = \prod_{l=1}^j (\beta - b_1 + l) \cdot \prod_{l=1}^j (\beta - b_2 + l) = \frac{\Gamma[(\beta - b_1) + j + 1] \cdot \Gamma[(\beta - b_2) + j + 1]}{\Gamma[(\beta - b_1) + 1] \cdot \Gamma[(\beta - b_2) + 1]}.$$

Evidently,

$$\Delta_0 = 1,$$

$$\Delta_1(\beta) = f_1(\beta),$$

$$\Delta_2(\beta) = \begin{vmatrix} f_1(\beta) & f_0(\beta+1) \\ f_2(\beta) & f_1(\beta+1) \end{vmatrix},$$

$$\Delta_3(\beta) = \begin{vmatrix} f_1(\beta) & f_0(\beta+1) & 0 \\ f_2(\beta) & f_1(\beta+1) & f_0(\beta+2) \\ f_3(\beta) & f_2(\beta+1) & f_1(\beta+2) \end{vmatrix}, \quad \text{etc.}$$

If  $b_1 - b_2 = k$  is not an integer, the denominator is

$$\frac{j! \Gamma(j+1+k)}{\Gamma(1+k)} \text{ if } \beta = b_1 \quad \text{and} \quad \frac{j! \Gamma(j+1-k)}{\Gamma(1-k)} \text{ if } \beta = b_2,$$

and

$$\left. \begin{aligned} C_j(b_1) &= \frac{(-1)^j C_0 \Gamma(1+k) \Delta_j(b_1)}{j! \Gamma(j+1+k)}, \\ C_j(b_2) &= \frac{(-1)^j C_0 \Gamma(1-k) \Delta_j(b_2)}{j! \Gamma(j+1-k)}. \end{aligned} \right\} \quad (7.10)$$

If the roots differ by an integer, however,  $C_j$  may become infinite at  $\beta \rightarrow b_2$ , since the denominator is then

$$[(1-k)(2-k)(3-k)\dots(k-k)\dots(j-k)]j!$$

Thus, when the roots differ by an integer,

$$\left. \begin{aligned} C_j(b_2) &= \frac{(k-j-1)! C_0 \Delta_j(b_2)}{(k-1)! j!}, & j < k; \\ C_j(b_2) &= \frac{(-1)^{j-k+1} C_0}{(k-1)! (j-k)! j!} \left[ \frac{\Delta_j(\beta)}{\beta - b_2} \right]_{\beta \rightarrow b_2}, & j \geq k. \end{aligned} \right\} \quad (7.10a)$$

The determinant may possibly contain the factor  $(\beta - b_2)$ ; in which case,  $C_j(b_2)$  remains finite. Generally, however, a new quantity is introduced:

$$D_j(\beta) = (\beta - b_2) \cdot C_j(\beta) = \frac{(-1)^j C_0 \Delta_j(\beta)}{\prod_{l=1}^{k-1} (\beta - b_1 + l) \prod_{l=k+1}^j (\beta - b_1 + l) \prod_{l=1}^j (\beta - b_2 + l)}. \quad (7.11)$$

When  $\beta \rightarrow b_2$ , Eq. (7.11) reduces to

$$D_j(b_2) = \frac{(-1)^{j-k+1} C_0 \Delta_j(b_2)}{(k-1)! (j-k)! j!}. \quad (7.12)$$

Froms Eqs. (7.05) and (7.10),

$$\frac{C_{k+j}(b_2)}{C_k(b_2)} = \frac{C_j(b_1)}{C_0} = \frac{(-1)^j k! \Delta_j(b_1)}{j! (k+j)!}.$$

But, according to Eq. (7.12),

$$\frac{C_{k+j}(b_2)}{C_k(b_2)} = \frac{(-1)^j k! \Delta_{k+j}(b_2)}{j! (k+j)! \Delta_k(b_2)}.$$

Thus,

$$\Delta_{k+j}(b_2) = \Delta_k(b_2) \cdot \Delta_j(b_1) \quad \text{if } k \text{ is an integer.} \quad (7.13)$$

Also needed are derivatives with respect to  $\beta$ . According to Eq. (7.09),

$$\left. \begin{aligned} \frac{\partial C_j}{\partial \beta} &= \frac{(-1)^j C_0}{\prod_{l=1}^j (\beta - b_1 + l) \cdot \prod_{l=1}^j (\beta - b_2 + l)} \\ &\times \left\{ \frac{\partial \Delta_j}{\partial \beta} - \Delta_j(\beta) \left[ \sum_{l=1}^j \frac{1}{\beta - b_1 + l} + \sum_{l=1}^j \frac{1}{\beta - b_2 + l} \right] \right\}. \end{aligned} \right\} \quad (7.14)$$

If  $\beta = b_1$ ,

$$\left[ \frac{\partial C_j}{\partial \beta} \right]_{\beta=b_1} = \frac{(-1)^j C_0 \Gamma(k+1)}{j! \Gamma(k+j+1)} \left\{ \left[ \frac{\partial \Delta_j}{\partial \beta} \right]_{\beta=b_1} - \Delta_j(b_1) \left[ \sum_{l=1}^j \frac{1}{l} + \sum_{l=1}^j \frac{1}{k+l} \right] \right\}. \quad (7.15)$$

Differentiation of Eq. (7.11) gives, when  $k$  is an integer,

$$\left. \begin{aligned} \frac{\partial D_j}{\partial \beta} &= \frac{(-1)^j C_0}{\prod_{l=1}^{k-1} (\beta - b_1 + l) \prod_{l=k+1}^j (\beta - b_1 + l) \prod_{l=1}^j (\beta - b_2 + l)} \\ &\times \left\{ \frac{\partial \Delta_j}{\partial \beta} - \Delta_j(\beta) \left[ \sum_{l=1}^{k-1} \frac{1}{\beta - b_1 + l} + \sum_{l=k+1}^j \frac{1}{\beta - b_1 + l} + \sum_{l=1}^j \frac{1}{\beta - b_2 + l} \right] \right\}. \end{aligned} \right\} \quad (7.16)$$

If  $\beta \rightarrow b_2$  and  $j \geq k$ , Eq. (7.16) becomes

$$\left. \begin{aligned} \left[ \frac{\partial D_j}{\partial \beta} \right]_{\beta \rightarrow b_2} &= \frac{(-1)^{j-k+1} C_0}{(k-1)! (j-k)! j!} \\ &\times \left\{ \left[ \frac{\partial A_j}{\partial \beta} \right]_{\beta \rightarrow b_2} - A_j(b_2) \left[ - \sum_{l=1}^{k-1} \frac{1}{l} + \sum_{l=1}^{j-k} \frac{1}{l} + \sum_{l=1}^k \frac{1}{l} \right] \right\}. \end{aligned} \right\} \quad (7.17)$$

### 7.03 THE FROBENIUS METHOD [27]

A solution of a Bôcher equation is to be obtained as a series expansion about a *regular singularity*. Evidently there are three possibilities [28] with respect to the roots of the indicial equation:

- I. The roots are distinct and do not differ by an integer.
- II. The roots are identical.
- III. The roots differ by an integer.

**Case I.** In I, the procedure of § 7.02 yields two distinct series:

$$\left. \begin{aligned} Z_1 &= C_0 \Gamma(1+k) \cdot (z-z_0)^{b_1} \sum_{j=0}^{\infty} \frac{(-1)^j A_j(b_1)}{j! \Gamma(j+1+k)} (z-z_0)^j, \\ Z_2 &= C_0 \Gamma(1-k) \cdot (z-z_0)^{b_2} \sum_{j=0}^{\infty} \frac{(-1)^j A_j(b_2)}{j! \Gamma(j+1-k)} (z-z_0)^j. \end{aligned} \right\} \quad (7.18)$$

The general solution of Eq. (7.01) is therefore

$$Z = A Z_1(b_1, z) + B Z_2(b_2, z).$$

But in II and III, only the first solution is obtained in this way. A new second solution, independent of  $Z_1$  is found by differentiation [28].

**Case II.** Consider the case of equal roots. The first solution is given by Eq. (7.18) with  $k=0$ :

$$Z_1 = C_0 (z-z_0)^{b_1} \sum_{j=0}^{\infty} \frac{(-1)^j A_j(b_1)}{(j!)^2} (z-z_0)^j. \quad (7.18a)$$

For the second solution, assume

$$Z(\beta, z) = C_0 (z-z_0)^\beta + \sum_{j=1}^{\infty} C_j(\beta) \cdot (z-z_0)^{\beta+j}, \quad (7.19)$$

where  $\beta$  is considered to be a variable and where  $C_0$  is the constant of § 7.02. Thus  $Z(\beta, z)$  satisfies the recursion formula, Eq. (7.05), but does not satisfy the indicial equation (7.07). Introduce the operator  $L$ :

$$L \equiv (z-z_0)^2 \frac{\partial^2}{\partial z^2} + (z-z_0) [(z-z_0) P(z)] \frac{\partial}{\partial z} + [(z-z_0)^2 Q(z)].$$

If  $\beta \neq b_1$ , Eq. (7.08) gives (for equal roots)

$$f_0(\beta) = (\beta - b_1)^2.$$

Thus,

$$L[Z(\beta, z)] = C_0(\beta - b_1)^2(z - z_0)^\beta \quad (7.20)$$

and

$$\frac{\partial}{\partial \beta} L(Z) = 2C_0(\beta - b_1)(z - z_0)^\beta + C_0(\beta - b_1)^2(z - z_0)^\beta \ln(z - z_0).$$

In general,  $\partial Z / \partial \beta$  does not constitute a solution of the differential equation. But if  $\beta \rightarrow b_1$ , the second solution is

$$Z_2 = \left[ \frac{\partial Z}{\partial \beta} \right]_{\beta \rightarrow b_1}. \quad (7.21)$$

Differentiation of Eq. (7.19) gives

$$\frac{\partial Z}{\partial \beta} = C_0(z - z_0)^\beta \ln(z - z_0) + \sum_{j=1}^{\infty} \frac{\partial C_j}{\partial \beta} \cdot (z - z_0)^{\beta+j} + \sum_{j=1}^{\infty} C_j(\beta) \cdot (z - z_0)^{\beta+j} \ln(z - z_0).$$

Thus, according to Eq. (7.15),

$$\begin{aligned} \left[ \frac{\partial Z}{\partial \beta} \right]_{\beta \rightarrow b_1} &= \left[ \sum_{j=0}^{\infty} C_j(b_1) \cdot (z - z_0)^{b_1+j} \right] \ln(z - z_0) \\ &\quad + C_0 \sum_{j=1}^{\infty} \frac{(-1)^j (z - z_0)^{b_1+j}}{(j!)^2} \left\{ \left[ \frac{\partial A_j}{\partial \beta} \right]_{\beta \rightarrow b_1} - 2A_j(b_1) \sum_{l=1}^j \frac{1}{l} \right\}. \end{aligned}$$

But the first bracket is  $Z_1$ , by Eq. (7.03), so the second solution is

$$\begin{aligned} Z_2 &= Z_1 \ln(z - z_0) \\ &\quad + C_0(z - z_0)^{b_1} \sum_{j=1}^{\infty} \frac{(-1)^j (z - z_0)^j}{(j!)^2} \left\{ \left[ \frac{\partial A_j}{\partial \beta} \right]_{\beta \rightarrow b_1} - 2A_j(b_1) \sum_{l=1}^j \frac{1}{l} \right\}. \end{aligned} \quad (7.22)$$

**Case III.** For III, the roots of the indicial equation are distinct but differ by an integer. The first solution is

$$Z_1 = C_0 k! (z - z_0)^{b_1} \sum_{j=0}^{\infty} \frac{(-1)^j A_j(b_1)}{j!(k+j)!} (z - z_0)^j. \quad (7.18b)$$

To obtain an independent second solution, let

$$\bar{Z}(\beta, z) = (\beta - b_2) \left[ C_0(z - z_0)^\beta + \sum_{j=1}^{k-1} C_j(\beta) \cdot (z - z_0)^{\beta+j} \right] + \sum_{j=k}^{\infty} D_j(\beta) \cdot (z - z_0)^{\beta+j}, \quad (7.23)$$

where  $D_j(\beta) = (\beta - b_2) \cdot a_j(\beta)$  as in § 7.02. A relation similar to Eq. (7.20) is

$$L[\bar{Z}(\beta, z)] = C_0(\beta - b_2)^2(\beta - b_1)(z - z_0)^\beta. \quad (7.24)$$

Also,

$$L\left(\frac{\partial \bar{Z}}{\partial \beta}\right) = \frac{\partial}{\partial \beta} [C_0(\beta - b_2)^2(\beta - b_1)(z - z_0)^\beta].$$

But

$$\begin{aligned} L\left(\frac{\partial \bar{Z}}{\partial \beta}\right) &= 2C_0(\beta - b_2)(\beta - b_1)(z - z_0)^\beta + C_0(\beta - b_2)^2(z - z_0)^\beta \\ &\quad + C_0(\beta - b_2)^2(\beta - b_1)(z - z_0)^\beta \ln(z - z_0). \end{aligned}$$

The entire expression vanishes when  $\beta = b_2$ . Thus a second solution is

$$Z_2 = \left[ \frac{\partial \bar{Z}}{\partial \beta} \right]_{\beta \rightarrow b_2}. \quad (7.25)$$

By differentiation of Eq. (7.23), we obtain

$$\left. \begin{aligned} \frac{\partial \bar{Z}}{\partial \beta} &= \sum_{j=0}^{k-1} C_j(\beta) \cdot (z - z_0)^{\beta+j} + (\beta - b_2) \left[ C_0(z - z_0)^\beta \cdot \ln(z - z_0) \right. \\ &\quad \left. + \sum_{j=1}^{k-1} \left\{ \frac{\partial C_j}{\partial \beta} \cdot (z - z_0)^{\beta+j} + C_j(\beta) \cdot (z - z_0)^{\beta+j} \cdot \ln(z - z_0) \right\} \right] \\ &\quad + \sum_{j=k}^{\infty} \left\{ \frac{\partial D_j}{\partial \beta} \cdot (z - z_0)^{\beta+j} + D_j(\beta) \cdot (z - z_0)^{\beta+j} \cdot \ln(z - z_0) \right\}. \end{aligned} \right\} \quad (7.26)$$

As  $\beta \rightarrow b_2$ , the first term becomes

$$\sum_{j=0}^{k-1} C_j(b_2) \cdot (z - z_0)^{b_2+j}.$$

All terms in the square bracket are finite, so  $(\beta - b_2)$  reduces this part of Eq. (7.26) to zero. There remain the summations containing  $D_j(\beta)$  and its derivative [29]. Substitution of Eqs. (7.10a), (7.12), and (7.17) gives the second solution,

$$\begin{aligned} Z_2 &= \frac{C_0(z - z_0)^{b_2}}{(k-1)!} \sum_{j=0}^{k-1} \frac{(k-j-1)! \Delta_j(b_2)}{j!} (z - z_0)^j \\ &\quad + \frac{C_0(z - z_0)^{b_2}}{(k-1)!} \ln(z - z_0) \sum_{j=k}^{\infty} \frac{(-1)^{j-k+1} \Delta_j(b_2) \cdot (z - z_0)^j}{(j-k)! j!} \\ &\quad + \frac{C_0(z - z_0)^{b_2}}{(k-1)!} \sum_{j=k}^{\infty} \frac{(-1)^{j-k+1} (z - z_0)^j}{(j-k)! j!} \\ &\quad \times \left\{ \left[ \frac{\partial \Delta_j}{\partial \beta} \right]_{\beta \rightarrow b_2} - \Delta_j(b_2) \left[ \sum_{l=1}^{j-k} \frac{1}{l} + \sum_{l=1}^j \frac{1}{l} - \sum_{l=1}^{k-1} \frac{1}{l} \right] \right\}. \end{aligned}$$

The second summation may be written

$$\frac{C_0(z - z_0)^{b_2+k}}{(k-1)!} \sum_{n=0}^{\infty} \frac{(-1)^{n+1} \Delta_{n+k}(b_2) \cdot (z - z_0)^n}{n! (n+k)!},$$

or, by Eq. (7.13),

$$\frac{C_0(z - z_0)^{b_1} \Delta_k(b_2)}{(k-1)!} \sum_{n=0}^{\infty} \frac{(-1)^{n+1} \Delta_n(b_1) \cdot (z - z_0)^n}{n! (n+k)!}.$$

Comparison with Eq. (7.18) shows that

$$\left. \begin{aligned} Z_2 &= - \frac{\Delta_k(b_2)}{k! (k-1)!} Z_1 \ln(z - z_0) \\ &\quad + \frac{C_0(z - z_0)^{b_2}}{(k-1)!} \sum_{j=0}^{k-1} \frac{(k-j-1)! \Delta_j(b_2)}{j!} (z - z_0)^j \\ &\quad + \frac{C_0(z - z_0)^{b_2}}{(k-1)!} \sum_{j=k}^{\infty} \frac{(-1)^{j-k+1} (z - z_0)^j}{(j-k)! j!} \\ &\quad \times \left\{ \left[ \frac{\partial \Delta_j}{\partial \beta} \right]_{\beta \rightarrow b_2} - \Delta_j(b_2) \left[ \sum_{l=1}^{j-k} \frac{1}{l} + \sum_{l=1}^j \frac{1}{l} - \sum_{l=1}^{k-1} \frac{1}{l} \right] \right\}. \end{aligned} \right\} \quad (7.27)$$

## 7.04 AN EXAMPLE, CASE I

As an example of the methods of obtaining a series solution, consider the *Bessel wave equation* {26}:

$$\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 + q^2 - p^2/z^2) Z = 0. \quad (7.28)$$

The equation has a second-order pole at the origin ( $z = 0$ ) and a sixth-order pole at infinity. An expansion will be obtained about the origin, which is a regular singularity.

As in § 7.02, let

$$Z = z^\beta \sum_{j=0}^{\infty} C_j z^j.$$

Then

$$[z P(z)] = 1, \quad A_0 = 1, \quad A_i = 0 \quad \text{for } i > 0;$$

$$[z^2 Q(z)] = \kappa^2 z^4 + q^2 z^2 - p^2,$$

$$B_0 = -p^2, \quad B_2 = q^2, \quad B_4 = \kappa^2,$$

with all other  $B$ 's equal to zero. The indicial equation is

$$\beta^2 - p^2 = 0,$$

so  $b_1 = +p$ ,  $b_2 = -p$ , and

$$t_0(\beta) = (\beta - p)(\beta + p).$$

From Eq. (7.06),  $f_2 = q^2$ ,  $f_4 = \kappa^2$ , and all other  $f$ 's are zero. Thus,

$$C_j(\beta) = \frac{(-1)^j C_0 \Delta_j(\beta)}{[1][2]\dots[j]},$$

where

$$\Delta_j(\beta) = \begin{vmatrix} 0 & [1] & 0 & 0 & \dots & \dots & \dots & 0 \\ q^2 & 0 & [2] & 0 & \dots & \dots & \dots & 0 \\ 0 & q^2 & 0 & [3] & \dots & \dots & \dots & 0 \\ \kappa^2 & 0 & q^2 & 0 & \dots & \dots & \dots & 0 \\ \dots & \dots \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & [j-1] \\ 0 & 0 & 0 & 0 & \dots & \kappa^2 & 0 & q^2 & 0 \end{vmatrix}, \quad (7.29)$$

and

$$[j] = [(\beta + j)^2 - p^2].$$

This determinant vanishes for odd values of  $j$ . For even values, the determinant may be simplified by expanding in terms of the elements [1], [2], etc. Then

$$C_{2m}(\beta) = \frac{(-1)^m C_0 \Delta'_m(\beta)}{[2][4]\dots[2m]},$$

where

$$\Delta'_m(\beta) = \begin{vmatrix} q^2 & [2] & 0 & 0 & \dots & \dots & \dots & 0 \\ \kappa^2 & q^2 & [4] & 0 & \dots & \dots & \dots & 0 \\ 0 & \kappa^2 & q^2 & [6] & \dots & \dots & \dots & 0 \\ 0 & 0 & \kappa^2 & q^2 & \dots & \dots & \dots & 0 \\ \dots & \dots \\ 0 & 0 & 0 & 0 & \dots & \kappa^2 & q^2 & [2(m-1)] \\ 0 & 0 & \dots & \dots & 0 & \kappa^2 & q^2 & \end{vmatrix}, \quad (7.30)$$

with

$$m = j/2.$$

The principal diagonal consists of elements  $q^2$ , the adjacent diagonals contain  $\varkappa^2$  and functions of  $\beta$ , and all other elements are zero.

For the first root,  $\beta = b_1 = +\beta$ , and

$$[2] = [(\beta + 2)^2 - \beta^2] = 4(\beta + 1),$$

$$[4] = [(\beta + 4)^2 - \beta^2] = 8(\beta + 2),$$

· ·

$$[2m] = [(\beta + 2m)^2 - \beta^2] = 4m(\beta + m),$$

$$C_{2m}(\beta) = \frac{(-1)^m C_0 \Gamma(\beta + 1) \Delta'_m(\beta)}{2^{2m} m! \Gamma(\beta + m + 1)}.$$

Therefore,

$$C_2 = -\frac{C_0 (q/2)^2}{1! (\beta + 1)},$$

$$C_4 = \frac{C_0 (q/2)^4}{2! (\beta + 1) (\beta + 2)} [1 - 4(\beta + 1) \varkappa^2/q^4],$$

$$C_6 = -\frac{C_0 (q/2)^6}{3! (\beta + 1) (\beta + 2) (\beta + 3)} [1 - 4(3\beta + 5) \varkappa^2/q^4], \dots$$

For convenience, let

$$C_0 = \frac{(q/2)^\beta}{\Gamma(\beta + 1)}.$$

The first solution of Eq. (7.28) is then defined as a *Bessel wave function*  $\mathcal{J}_\beta(\varkappa, q, z)$ :

$$\left. \begin{aligned} Z_1 &= \mathcal{J}_\beta(\varkappa, q, z) = \frac{(qz/2)^\beta}{\Gamma(\beta + 1)} \left\{ 1 - \frac{(qz/2)^2}{1! (\beta + 1)} + \frac{(qz/2)^4 [1 - 4(\beta + 1) \varkappa^2/q^4]}{2! (\beta + 1) (\beta + 2)} \right. \\ &\quad \left. - \frac{(qz/2)^6 [1 - 4(3\beta + 5) \varkappa^2/q^4]}{3! (\beta + 1) (\beta + 2) (\beta + 3)} + \dots \right\} \\ &= (qz/2)^\beta \sum_{m=0}^{\infty} \frac{(-1)^m \Delta'_m(\beta) \cdot (z/2)^{2m}}{m! \Gamma(m + \beta + 1)} \end{aligned} \right\} \quad (7.31)$$

with  $\Delta_0 = 1$ ,  $\Delta_1 = q^2$ . Similarly, for the other root of the indicial equation,  $\beta = -\beta$  and

$$\mathcal{J}_{-\beta}(\varkappa, q, z) = (qz/2)^{-\beta} \sum_{m=0}^{\infty} \frac{(-1)^m \Delta'_m(-\beta) \cdot (z/2)^{2m}}{m! \Gamma(m - \beta + 1)}. \quad (7.32)$$

The complete solution of Eq. (7.28), for non-integral values of  $\beta$ , is

$$Z = A \mathcal{J}_\beta(\varkappa, q, z) + B \mathcal{J}_{-\beta}(\varkappa, q, z). \quad (7.33)$$

If  $\varkappa = 0$ , Eq. (7.31) reverts to the ordinary Bessel function of the first kind,

$$\left. \begin{aligned} \mathcal{J}_\beta(qz) &= \frac{(qz/2)^\beta}{\Gamma(\beta + 1)} \left\{ 1 - \frac{(qz/2)^2}{1! (\beta + 1)} \right. \\ &\quad \left. + \frac{(qz/2)^4}{2! (\beta + 1) (\beta + 2)} - \frac{(qz/2)^6}{3! (\beta + 1) (\beta + 2) (\beta + 3)} + \dots \right\} \\ &= \sum_{m=0}^{\infty} \frac{(-1)^m (qz/2)^{\beta+2m}}{m! \Gamma(m + \beta + 1)} \end{aligned} \right\} \quad (7.34)$$

## 7.05 EXAMPLE, CASE II

With Bessel wave functions, the only case of equal roots occurs when  $b_1 = b_2 = 0$ . Then  $C_0 = 1$  and the first solution is

$$\left. \begin{aligned} Z_1 = J_0(\kappa, q, z) &= 1 - \frac{(qz/2)^2}{(1!)^2} + \frac{(qz/2)^4}{(2!)^2} [1 - 4\kappa^2/q^4] \\ &\quad - \frac{(qz/2)^6}{(3!)^2} [1 - 20\kappa^2/q^4] + \dots \\ &= \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(0) \cdot (z/2)^{2m}}{(m!)^2}. \end{aligned} \right\} \quad (7.31 \text{a})$$

Since all odd determinants are zero, Eq. (7.22) becomes

$$Z_2 = Z_1 \ln z + \sum_{m=1}^{\infty} \frac{z^{2m}}{(2m)!} \left\{ \left[ \frac{\partial A_{2m}}{\partial \beta} \right]_{\beta \rightarrow 0} - 2A_m(0) \sum_{l=1}^{2m} \frac{1}{l} \right\}. \quad (7.22 \text{a})$$

But

$$\begin{aligned} A_{2m}(\beta) &= (-1)^m [1][3]\dots[2m-1] A'_m(\beta), \\ \frac{\partial A_{2m}}{\partial \beta} &= (-1)^m [1][3]\dots[2m-1] \left\{ \frac{\partial A'_m}{\partial \beta} + 2A'_m(\beta) \sum_{l=1}^m \frac{1}{\beta + 2l-1} \right\}, \\ \frac{\partial A'_m}{\partial \beta} &= - \left\{ \frac{\partial [2]}{\partial \beta} M_{12}(\beta) + \frac{\partial [4]}{\partial \beta} M_{23}(\beta) + \dots + \frac{\partial [2(m-1)]}{\partial \beta} M_{(m-1)m}(\beta) \right\} \\ &= -2 \{ (\beta+2) M_{12}(\beta) + (\beta+4) M_{23}(\beta) + \dots + [\beta+2(m-1)] M_{(m-1)m}(\beta) \}. \end{aligned}$$

Thus, the second solution may be written

$$Z_2 = Z_1 \ln z - \sum_{m=1}^{\infty} \frac{(-1)^m (z/2)^{2m}}{(m!)^2} \left\{ 4 \sum_{l=1}^{m-1} l M_{l(l+1)}(0) + A'_m(0) \sum_{l=1}^m \frac{1}{l} \right\}, \quad (7.35)$$

where  $M_{l(l+1)}$  is the minor of the corresponding element in the determinant  $A'_m$ .

Bessel functions of the second kind, however, are usually written in a slightly different form from the above. Let

$$\mathcal{Y}_0(\kappa, q, z) = \frac{2}{\pi} [Z_2 + [\gamma + \ln(q/2)] J_0(\kappa, q, z)]. \quad (7.36)$$

Substitution of Eq. (7.35) gives

$$\left. \begin{aligned} \mathcal{Y}_0(\kappa, q, z) &= \frac{2}{\pi} \left\{ [\gamma + \ln(qz/2)] J_0(\kappa, q, z) \right. \\ &\quad \left. - \sum_{m=1}^{\infty} \frac{(-1)^m (z/2)^{2m}}{(m!)^2} \left[ 4 \sum_{l=1}^{m-1} l M_{l(l+1)}(0) + A'_m(0) \sum_{l=1}^m \frac{1}{l} \right] \right\} \right\} \quad (7.37)$$

For the special case of  $\kappa = 0$ , Eq. (7.37) reduces to

$$\mathcal{Y}_0(qz) = \frac{2}{\pi} \left\{ [\gamma + \ln(qz/2)] J_0(qz) - \sum_{m=1}^{\infty} \frac{(-1)^m (qz/2)^{2m}}{(m!)^2} \sum_{l=1}^m \frac{1}{l} \right\}, \quad (7.37 \text{a})$$

in accordance with the usual formulation of NIELSEN [30].

### 7.06 EXAMPLE, CASE III

If  $p = n$ , an integer not equal to zero, the second solution of the Bessel wave equation is given by Eq. (7.27) with  $b_1 = n$ ,  $b_2 = -n$ ,  $k = 2n$ ,  $j = 2m$ :

$$\left. \begin{aligned} Z_2 &= -\frac{\Delta_{2n}(-n)}{(2n)! (2n-1)!} Z_1 \ln z \\ &+ \frac{C_0 z^{-n}}{(2n-1)!} \sum_{m=0}^{n-1} \frac{(2n-2m-1)! \Delta_{2m}(-n)}{(2m)!} z^{2m} \\ &+ \frac{C_0 z^{-n}}{(2n-1)!} \sum_{m=n}^{\infty} \frac{(-1)^{2m-2n-1} z^{2m}}{(2m-2n)! (2m)!} \\ &\times \left[ \left[ \frac{\partial \Delta_{2m}}{\partial \beta} \right]_{\beta \rightarrow -n} - \Delta_{2m}(-n) \left[ \sum_{l=1}^{2m-2n} \frac{1}{l} + \sum_{l=1}^{2m} \frac{1}{l} - \sum_{l=1}^{2n-1} \frac{1}{l} \right] \right] \end{aligned} \right\} \quad (7.27a)$$

But

$$\frac{\Delta_{2n}(-n)}{(2n)! (2n-1)!} = \frac{\Delta'_n(-n)}{2^{2n-1} n! (n-1)!}. \quad (7.38)$$

For  $m < n$ ,

$$\frac{(2n-2m-1)! \Delta_{2m}(-n)}{(2m)!} = \frac{(2n)! (n-m-1)! \Delta'_m(-n)}{2^{2m+1} m! n!}. \quad (7.39)$$

For  $m > n$ ,

$$\Delta_{2m}(-n) = (-1)^{m+n} \frac{(2m)! (2n)! (2m-2n)!}{2^{2m} m! n! (m-n)!} \Delta'_m(-n). \quad (7.40)$$

Also,

$$\begin{aligned} \frac{\partial \Delta_{2m}}{\partial \beta} &= (-1)^{m+n} \frac{(2m)! (2n)! (2m-2n)!}{2^{2m} m! n! (m-n)!} \\ &\times \left\{ \frac{\partial \Delta'_m}{\partial \beta} + 2 \Delta'_m(\beta) \left[ \frac{\beta+1}{[1]} + \frac{\beta+3}{[3]} + \dots + \frac{\beta+2m-1}{[2m-1]} \right] \right\}, \\ \left[ \frac{\partial \Delta'_m}{\partial \beta} \right]_{\beta \rightarrow -n} &= 2 \sum_{l=1}^{m-1} (n-2l) M_{l(l+1)}, \end{aligned}$$

where  $M_{l(l+1)}$  is the minor of the corresponding term in the determinant  $\Delta'_m$ .

Substitution of these relations into Eq. (7.27a) gives

$$\begin{aligned} Z_2 &= -\frac{\Delta'_n(-n)}{2^{2n-1} n! (n-1)!} Z_1 \ln z + \frac{C_0 z^{-n}}{(n-1)!} \sum_{m=0}^{n-1} \frac{(n-m-1)! \Delta'_m(-n) \cdot (z/2)^{2m}}{m!} \\ &+ \frac{C_0 z^{-n}}{(n-1)!} \sum_{m=n}^{\infty} \frac{(-1)^{m-n-1} (z/2)^{2m}}{m! (m-n)!} \\ &\times \left\{ 4 \sum_{l=1}^{m-1} (n-2l) M_{l(l+1)} - \Delta'_m(-n) \left[ \sum_{l=1}^{m-n} \frac{1}{l} + \sum_{l=1}^m \frac{1}{l} - \sum_{l=1}^{n-1} \frac{1}{l} \right] \right\}. \end{aligned}$$

A change of the second summation so that it begins at  $m=0$ , and use of the relation,

$$\Delta'_{m+n}(b_2) = \Delta'_m(b_2) \cdot \Delta'_n(b_1),$$

gives

$$\left. \begin{aligned}
 Z_2 = & -\frac{1}{2^{2n-1} n! (n-1)!} \\
 & \times \left\{ \left[ \ln(qz/2) - \ln(q/2) + 2^{2n-1} \sum_{l=1}^{n-1} \frac{1}{l} \right] \Delta'_n(-n) \cdot \mathcal{I}_n(\varkappa, q, z) \right. \\
 & - \frac{1}{2} (q/2)^n 2^{2n} z^{-n} \sum_{m=0}^{n-1} \frac{(n-m-1)! \Delta'_m(-n) \cdot (z/2)^{2m}}{m!} \\
 & + \frac{1}{2} (qz/2)^n \sum_{m=0}^{\infty} \frac{(-1)^m (z/2)^{2m}}{m! (m+n)!} \\
 & \left. \times \left\{ 4 \sum_{l=1}^{m+n-1} (n-2l) M_{l(l+1)} - \Delta'_m(n) \Delta'_n(-n) \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \right\} \right\}. 
 \end{aligned} \right\} \quad (7.41)$$

To make this solution more nearly like the familiar Bessel function, we introduce the *Bessel wave function of the second kind*, defined by

$$\mathcal{Y}_n(\varkappa, q, z) = -\frac{2^{2n} n! (n-1)!}{\pi q^{2n}} Z_2 + \frac{2 \mathcal{I}_n(\varkappa, q, z)}{\pi q^{2n}} \left[ \gamma + \ln(q/2) - 2^{2n-1} \sum_{l=1}^{n-1} \frac{1}{l} \right].$$

Substitution of Eq. (7.41) results in the second solution,

$$\left. \begin{aligned}
 \mathcal{Y}_n(\varkappa, q, z) = & \frac{2}{\pi q^{2n}} \left\{ [\gamma + \ln(qz/2)] \Delta'_n(-n) \mathcal{I}_n(\varkappa, q, z) \right. \\
 & - \frac{1}{2} (q/2)^n 2^{2n} z^{-n} \sum_{m=0}^{n-1} \frac{(n-m-1)! \Delta'_m(-n) \cdot (z/2)^{2m}}{m!} \\
 & + \frac{1}{2} (qz/2)^n \sum_{m=0}^{\infty} \frac{(-1)^m (z/2)^{2m}}{m! (m+n)!} \\
 & \left. \times \left\{ 4 \sum_{l=1}^{m+n-1} (n-2l) M_{l(l+1)} - \Delta'_m(n) \Delta'_n(-n) \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \right\} \right\}.
 \end{aligned} \right\} \quad (7.42)$$

If  $\varkappa = 0$ , Eq. (7.42) reduces to the Bessel function as written by NIELSEN [30]:

$$\left. \begin{aligned}
 \mathcal{Y}_n(qz) = & \frac{2}{\pi} \left\{ [\gamma + \ln(qz/2)] \mathcal{J}_n(qz) - \frac{1}{2} (qz/2)^{-n} \sum_{m=0}^{n-1} \frac{(n-m-1)! (qz/2)^{2m}}{m!} \right. \\
 & - \frac{1}{2} (qz/2)^n \sum_{m=0}^{\infty} \frac{(-1)^m (qz/2)^{2m}}{m! (m+n)!} \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \left. \right\}.
 \end{aligned} \right\} \quad (7.42a)$$

## 7.07 ORTHOGONALITY

Solution of a Bôcher equation of field theory yields two functions  $Z_1(\varkappa, \phi, q, z)$  and  $Z_2(\varkappa, \phi, q, z)$  where  $\phi$  and  $q$  are separation constants. A series of these functions is then employed to fit the boundary conditions and thus to obtain a unique solution of a field problem. The evaluation of the coefficients in the series is greatly facilitated if the functions are *orthogonal* on the given interval.

The question of orthogonality could be investigated separately for each function. But it is advantageous to consider the general subject of orthogonality for all the functions obtained in field theory. The STURM-LIOUVILLE [31] treatment is presented here in slightly modified form.

The Bôcher equation is

$$\frac{d^2Z}{dz^2} + P(z) \frac{dZ}{dz} + Q(z) Z = 0, \quad (7.43)$$

with general boundary conditions

$$\left. \begin{aligned} z = a, \quad k_1 Z'_i(a) + k_2 Z_i(a) &= 0; \\ z = b, \quad k_3 Z'_i(b) + k_4 Z_i(b) &= 0; \end{aligned} \right\} \quad (7.44)$$

where  $Z'_i(a) = (dZ/dz)_{z=a}$ ,  $i = 1, 2$ , and the  $k$ 's are constants. These general conditions include DIRICHLET, NEUMANN and mixed conditions.

Introduce a new quantity  $u$ :

$$u(z) = K e^{\int P(z) dz}$$

where  $K$  is an arbitrary constant, or

$$\frac{1}{u} \frac{du}{dz} = P(z). \quad (7.45)$$

Also let

$$Q(z) = \frac{1}{u(z)} [v(z) + f(\lambda) w(z)]. \quad (7.46)$$

Here  $\lambda$  is an *eigenvalue*, and  $w(z)$  is a *weighting function*. If the differential equation contains only one separation constant, say  $p$ , then  $\lambda$  is a function of  $p$  only, and the various eigenvalues  $\lambda_n$  are expressed in terms of the allowable values  $p_n$  that fit the boundary conditions. In this case, the quantities  $u$ ,  $v$ , and  $w$  do not contain  $p$ . If there are two separation constants,  $p$  and  $q$ , boundary conditions determine which is to be associated with the eigenvalues and which is to be regarded as a constant. The latter may appear in  $v$  and  $w$ . Substitution of Eqs. (7.45) and (7.46) into Eq. (7.43) gives

$$\frac{d^2Z}{dz^2} + \left( \frac{1}{u} \frac{du}{dz} \right) \frac{dZ}{dz} + \frac{1}{u} [v(z) + f(\lambda) w(z)] Z = 0$$

or

$$\frac{d}{dz} \left[ u \frac{dZ}{dz} \right] + [v + f(\lambda) w] Z = 0. \quad (7.47)$$

Eqs. (7.47) and (7.44) are said to constitute a *Sturm-Liouville system* [31].

It is easily proved that the solutions of a Sturm-Liouville system are *orthogonal on an arbitrary interval*  $(a, b)$ . Take  $Z_m$  and  $Z_n$  as any two distinct solutions of Eq. (7.43), corresponding to the distinct eigenvalues  $\lambda_m$  and  $\lambda_n$ . Then

$$\left\{ \begin{aligned} \frac{d}{dz} \left[ u \frac{dZ_m}{dz} \right] + [v + f(\lambda_m) w] Z_m &= 0, \\ \frac{d}{dz} \left[ u \frac{dZ_n}{dz} \right] + [v + f(\lambda_n) w] Z_n &= 0. \end{aligned} \right.$$

Multiplication by  $Z_n$  and  $Z_m$ :

$$\begin{cases} Z_n \frac{d}{dz} \left[ u \frac{dZ_m}{dz} \right] + [v + f(\lambda_m) w] Z_m Z_n = 0, \\ Z_m \frac{d}{dz} \left[ u \frac{dZ_n}{dz} \right] + [v + f(\lambda_n) w] Z_n Z_m = 0, \end{cases}$$

and subtraction gives

$$[f(\lambda_m) - f(\lambda_n)] w Z_m Z_n = \frac{d}{dz} [u Z'_n Z_m - u Z'_m Z_n]. \quad (7.48)$$

Integration over the arbitrary interval  $(a, b)$  yields

$$\begin{aligned} [f(\lambda_m) - f(\lambda_n)] \int_a^b w Z_m Z_n dz &= u(b) [Z'_n(b) Z_m(b) - Z'_m(b) Z_n(b)] \\ &\quad - u(a) [Z'_n(a) Z_m(a) - Z'_m(a) Z_n(a)]. \end{aligned}$$

Boundary conditions, Eq. (7.44), are now substituted, reducing the bracketed quantities to zero. Thus,

$$\int_a^b w Z_m Z_n dz = 0, \quad (7.49)$$

which is the *condition for orthogonality on the interval  $(a, b)$  with respect to the weighting function  $w(z)$* . As indicated by Table 7.01, all the ordinary differential equations of this book, with general boundary conditions of field theory, constitute Sturm-Liouville systems. Consequently, the solutions of these equations can be used to build up orthogonal series of eigenfunctions.

Given a Sturm-Liouville system and an arbitrary function  $f(z)$  on the interval  $(a, b)$ . This function can be expressed in terms of the eigenfunctions  $Z_n$  obtained as solutions of the differential equation:

$$f(z) = \sum_{n=0}^{\infty} A_n Z_n(z).$$

Because of orthogonality, with respect to weighting function  $w(z)$ , on interval  $(a, b)$ ,

$$A_n = \frac{1}{N_n} \int_a^b w(z) f(z) Z_n(z) dz,$$

where the norm is

$$N_n = \int_a^b w(z) [Z_n(z)]^2 dz.$$

There are, of course, restrictions on the functions  $f(z)$  that can be handled in this way. But these restrictions are so light that they can be ignored in practically all physical applications. Accordingly, no further attention will be paid to such restrictions. The fitting of boundary values by this method has had a great many practical applications [32] in physics and engineering; and the additional coordinate systems now available should allow a still wider field of usefulness.

TABLE 7.01. ORTHOGONALITY

| Equation   | $u$     | $v$                                      | $w$          | $f(\lambda)$                | $\lambda$  |
|--|---------|--|--------------|-----------------------------|------------|
| $\{04\}$<br>$\frac{d^2Z}{dz^2} + p^2 Z = 0.$   | 1       | 0  | 1            | $\lambda^2$                 | $p$        |
| Eigenfunctions: $\sin pz$ , $\cos pz$ .  |         |  |              |                             |            |
| $\{06\}$ Weber<br>$\frac{d^2Z}{dz^2} + [q^2(p + \frac{1}{2}) - q^4 z^2/4] Z = 0,$<br>$\mathcal{W}_o(p, qz)$ , $\mathcal{W}_o(p, qz)$ .                                 | $1/q^2$ | $-q^2 z^2/4$                             | 1            | $\lambda + \frac{1}{2}$     | $p$        |
| $\{24\}$ Bessel<br>$\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (q^2 - p^2/z^2) Z = 0,$<br>$\mathcal{J}_p(qz)$ , $\mathcal{J}_{-p}(qz)$ .                          | $z$     | $-p^2/z$                                 | $z$          | $\lambda^2$                 | $q$        |
| $\{26\}$ Bessel wave<br>$\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (x^2 z^2 + q^2 - p^2/z^2) Z = 0,$<br>$\mathcal{I}_p(x, q, z)$ , $\mathcal{I}_{-p}(x, q, z)$ . | $z$     | $(x^2 z^3 - p^2/z)$<br>$(x^2 z^2 + q^2)$ | $z$<br>$1/z$ | $\lambda^2$<br>$-\lambda^2$ | $q$<br>$p$ |

Table 7.01. Orthogonality

|   |   |                              |                                   |           |                       |     |
|---|---|------------------------------|-----------------------------------|-----------|-----------------------|-----|
| $\{113\}$ Baer                                  | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-b} + \frac{1}{z-c} \right] \frac{dZ}{dz}$                                      | $[(z-b)(z-c)]^{\frac{1}{2}}$ | $-q(b+c)/u$                       | $(b+c)/u$ | $-\lambda(\lambda+1)$ | $p$ |
|   | $-\left[ \frac{p(p+1)z+q(b+c)}{(z-b)(z-c)} \right] Z = 0,$  | $[(z-b)(z-c)]^{\frac{1}{2}}$ | $-\phi(p+1)z/u$                   | $(b+c)/u$ | $-\lambda$            | $q$ |
| $\mathcal{B}_p^q(z), \quad \mathcal{C}_p^q(z).$ |   |                              |                                   |           |                       |     |
|   |   |                              |                                   |           |                       |     |
| $\{114\}$ Baer wave                             | $\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-b} + \frac{1}{z-c} \right] \frac{dZ}{dz}$                                      | $[(z-b)(z-c)]^{\frac{1}{2}}$ | $\frac{1}{u} [x^2 z^2 - q(b+c)]$  | $(b+c)/u$ | $-\lambda(\lambda+1)$ | $p$ |
|   | $+\left[ \frac{x^2 z^2 - p(p+1)z - q(b+c)}{(z-b)(z-c)} \right] Z = 0,$  | $[(z-b)(z-c)]^{\frac{1}{2}}$ | $\frac{1}{u} [x^2 z^2 - p(p+1)z]$ | $(b+c)/u$ | $-\lambda$            | $q$ |
| $\{222\}$ Legendre                              |   |                              |                                   |           |                       |     |
|   |   |                              |                                   |           |                       |     |
| $\{12*$   | $\frac{d^2Z}{dz^2} + \left( \frac{2z}{z^2-1} \right) \frac{dZ}{dz} - \left[ \frac{p(p+1)(z^2-1) + q^2}{(z^2-1)^2} \right] Z = 0,$ | $z^2 - 1$                    | $-q^2/u$                          | $1$       | $-\lambda(\lambda+1)$ | $p$ |
|   | $\mathcal{B}_p^q(z), \quad \mathcal{D}_p^q(z).$   | $z^2 - 1$                    | $-\phi(p+1)$                      | $1/u$     | $-\lambda^2$          | $q$ |

Table 7.01. Continuation

| Equation  | $u$  | $v$   | $w$                      | $f(\lambda)$                         | $\lambda$  |
|---|--|---|--------------------------|--------------------------------------|------------|
| $\{224\}$ Legendre wave   | $z^2 - 1$  | $\frac{1}{u} [\kappa^2 a^2 u^2 - q^2]$  | 1                        | $-\lambda(\lambda + 1)$              | $p$        |
| $\frac{d^2 Z}{dz^2} + \left( \frac{2z}{z^2 - 1} \right) \frac{dZ}{dz}$<br>+ $\left[ \frac{\kappa^2 a^2 (z^2 - 1)^2 - p(p + 1)(z^2 - 1) - q^2}{(z^2 - 1)^2} \right] Z = 0,$<br>$\mathcal{P}_p^q(xa, z), \quad \mathcal{D}_p^q(xa, z).$   | $z^2 - 1$  | $[\kappa^2 a^2 u - p(p + 1)]$   | $1/u$                    | $-a^2$                               | $q$        |
| $\{1112\}$ Lamé   | $[z(z - b)(z - c)]^{\frac{1}{2}}$<br>$[z(z - b)(z - c)]^{\frac{1}{2}}$ | $(b^2 + c^2)q/u$<br>$-p(p + 1)z/u$  | $z/u$<br>$(b^2 + c^2)/u$ | $-\lambda(\lambda + 1)$<br>$\lambda$ | $p$<br>$q$ |
| $\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{1}{z - b} + \frac{1}{z - c} \right] \frac{dZ}{dz}$<br>+ $\left[ \frac{(b^2 + c^2)q - p(p + 1)z}{z(z - b)(z - c)} \right] Z = 0,$<br>$\mathcal{C}_p^q(z^{\frac{1}{2}}), \quad \mathcal{F}_p^q(z^{\frac{1}{2}}).$                      |  |   |                          |                                      |            |
| $\{1113\}$ Lamé wave  | $[z(z - b)(z - c)]^{\frac{1}{2}}$<br>$[z(z - b)(z - c)]^{\frac{1}{2}}$ | $\frac{1}{u} [\kappa^2 z^2 + (b^2 + c^2)q]$<br>$\frac{1}{u} [\kappa^2 z^2 - p(p + 1)z]$ | $z/u$<br>$(b^2 + c^2)/u$ | $-\lambda(\lambda + 1)$<br>$\lambda$ | $p$<br>$q$ |
| $\frac{d^2 Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{1}{z - b} + \frac{1}{z - c} \right] \frac{dZ}{dz}$<br>+ $\left[ \frac{\kappa^2 z^2 - p(p + 1)z + (b^2 + c^2)q}{z(z - b)(z - c)} \right] Z = 0,$<br>$\mathcal{C}_p^q(x, z^{\frac{1}{2}}), \quad \mathcal{F}_p^q(x, z^{\frac{1}{2}}).$ |  |   |                          |                                      |            |

|  |  |
|--|--|
| $\{1122\}$ Wangerin<br>$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-1} + \frac{1}{z-c} + \frac{2}{z} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{-q^2c - p^2cz + (1-q^2)z^2}{z^2(z-1)(z-c)} \right] Z = 0,$ $\mathcal{W}_p^q(k, z), \quad \mathcal{T}_p^q(k, z).$ <p>Or</p> $u = z[(z-1)(z-c)]^{\frac{1}{2}}, \quad f(\lambda) = -\lambda^2, \quad \lambda = p.$ $w(z) = \frac{cz}{4u}, \quad v(z) = \frac{1}{4u}[-q^2c + (1-q^2)z^2].$ $f(\lambda) = -\lambda^2, \quad \lambda = q.$ $w(z) = \frac{1}{4u}[z^2 + c], \quad v = \frac{z}{4u}[z - p^2c].$ | $\{1222\}$ Heine<br>$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{2}{z-1} + \frac{2}{z-c} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{-p^2c^2 + [p^2c(1+c) - q^2(1-c)^2 + 2c]z}{z(z-1)^2} \right. \\ \left. - \frac{[p^2c + 2(1+c)]z^2 + 2z^3}{\times(z-c)^2} \right] Z = 0,$ $\mathcal{W}_p^q(k, z), \quad \mathcal{V}_p^q(k, z).$ <p>Or</p> $u = z^{\frac{1}{2}}(z-1)(z-c), \quad f(\lambda) = -\lambda^2, \quad \lambda = p.$ $w(z) = \frac{c}{4u}[c - (1+c)z + z^2],$ $v(z) = \frac{z}{4u}[-q^2(1-c)^2 + 2c - 2(1+c)z + 2z^2].$ $f(\lambda) = -\lambda^2, \quad \lambda = q. \quad w(z) = \frac{z}{4u}(1-c)^2,$ $v(z) = \frac{1}{4u}[-p^2c^2 + [p^2c(1+c) + 2c]z - [p^2c + 2(1+c)]z^2 + 2z^3].$ |
|--|--|

### 7.08 WEBER FUNCTIONS

As the basic form of the {06} equation, we have, according to Table 6.02,

$$\frac{d^2Z}{dz^2} + [M + Nz^2] Z = 0. \quad (7.50)$$

If  $M = q^2(p + \frac{1}{2})$  and  $N = -q^4/4$ , Eq. (7.50) becomes

$$\frac{d^2Z}{dz^2} + [q^2(p + \frac{1}{2}) - q^4z^2/4] Z = 0, \quad (7.51)$$

which is *Weber's equation* [33].

Expansion in an infinite series about the origin gives

$$\left. \begin{aligned} \mathcal{W}_e(p, qz) &= e^{-(qz)^2/4} \left\{ 1 - \frac{p}{2!} (qz)^2 + \frac{p}{4!} (p-2)(qz)^4 \right. \\ &\quad \left. - \frac{p}{6!} (p-2)(p-4)(qz)^6 + \dots \right\} \\ &= e^{-(qz)^2/4} \left\{ 1 + \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} p(p-2)(p-4)\dots(p-2[n-1])(qz)^{2n} \right\}, \end{aligned} \right\} \quad (7.52)$$

$$\left. \begin{aligned} \mathcal{W}_o(p, qz) &= e^{-(qz)^2/4} (qz) \left\{ 1 - \frac{1}{3!} (p-1)(qz)^2 + \frac{1}{5!} (p-1)(p-3)(qz)^4 \right. \\ &\quad \left. - \frac{1}{7!} (p-1)(p-3)(p-5)(qz)^6 + \dots \right\} \\ &= e^{-(qz)^2/4} (qz) \left\{ 1 + \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n+1)!} (p-1)(p-3)\dots(p-[2n-1])(qz)^{2n} \right\}. \end{aligned} \right\} \quad (7.53)$$

Both series are absolutely convergent for the finite  $z$ -plane. The first function is even, the second odd. They may be called *Weber functions*, and the general solution of Eq. (7.51) may be written

$$Z = A\mathcal{W}_e(p, qz) + B\mathcal{W}_o(p, qz).$$

For the special case of  $p$  an *even integer*,

$$\mathcal{W}_e(p, qz) = (-2)^{p/2} \frac{(p/2)!}{p!} e^{-(qz)^2/4} H_p(qz). \quad (7.54)$$

If  $p$  is an *odd integer*,

$$\mathcal{W}_o(p, qz) = (-2)^{(p-1)/2} \frac{[(p-1)/2]!}{p!} e^{-(qz)^2/4} H_p(qz), \quad (7.55)$$

where  $H_p(qz)$  are the *Hermite polynomials* [34]:

$$H_0 = 1,$$

$$H_1 = qz,$$

$$H_2 = (qz)^2 - 1,$$

$$H_3 = (qz)^3 - 3(qz),$$

$$H_4 = (qz)^4 - 6(qz)^2 + 3,$$

$$H_5 = (qz)^5 - 10(qz)^3 + 15(qz),$$

$$H_6 = (qz)^6 - 15(qz)^4 + 45(qz)^2 - 15, \dots$$

Except for these special cases, the series does not terminate and solutions are given by the infinite series of Eqs. (7.52) and (7.53).

For the alternative form of the Weber equation,

$$\frac{d^2Z}{dz^2} - [q^2(p + \frac{1}{2}) + q^4 z^2/4] Z = 0,$$

the solution is

$$Z = A \mathcal{W}_e(p, iqz) + B \mathcal{W}_o(p, iqz).$$

Here

$$\left. \begin{aligned} \mathcal{W}_e(p, iqz) &= e^{(qz)^2/4} \left\{ 1 + \frac{p}{2!} (qz)^2 + \frac{p}{4!} (p-2)(qz)^4 \right. \\ &\quad \left. + \frac{p}{6!} (p-2)(p-4)(qz)^6 + \dots \right\} \\ &= e^{(qz)^2/4} \left\{ 1 + \sum_{n=1}^{\infty} \frac{1}{(2n)!} p(p-2)(p-4)\dots(p-2[n-1])(qz)^{2n} \right\}, \end{aligned} \right\} \quad (7.56)$$

$$\left. \begin{aligned} \mathcal{W}_o(p, iqz) &= e^{(qz)^2/4} (iqz) \left\{ 1 + \frac{1}{3!} (p-1)(qz)^2 + \frac{1}{5!} (p-1)(p-3)(qz)^4 \right. \\ &\quad \left. + \frac{1}{7!} (p-1)(p-3)(p-5)(qz)^6 + \dots \right\} \\ &= e^{(qz)^2/4} (iqz) \left\{ 1 + \sum_{n=1}^{\infty} \frac{1}{(2n+1)!} (p-1)(p-3)\dots(p-[2n-1])(qz)^{2n} \right\}. \end{aligned} \right\} \quad (7.57)$$

WHITTAKER [35] expresses the solution of the Weber equation in degenerate forms of the confluent hypergeometric function, which he denotes [36] by  $D$ :

$$Z = A D_p(qz) + B D_{-(p+1)}(iqz).$$

There are some advantages, however, in the use of the simple even and odd functions of Eqs. (7.52) and (7.53).

### Special values

$$\begin{aligned} \mathcal{W}_e(p, 0) &= 1, & \mathcal{W}_o(p, 0) &= 0; \\ \mathcal{W}_e(p, \pm\infty) &= 0, & \mathcal{W}_o(p, \pm\infty) &= 0. \end{aligned}$$

### Asymptotic expansions

$$\begin{aligned} \mathcal{W}_e(p, qz) &\sim (-2)^{p/2} \frac{\Gamma(p/2 + 1)}{\Gamma(p + 1)} e^{-(qz)^2/4} (qz)^p \left[ 1 - \frac{p(p-1)}{2(qz)^2} \right. \\ &\quad \left. + \frac{p(p-1)(p-2)(p-3)}{2 \cdot 4 (qz)^4} - \frac{p(p-1)(p-2)(p-3)(p-4)(p-5)}{2 \cdot 4 \cdot 6 (qz)^6} + \dots \right], \end{aligned}$$

$$\begin{aligned} \mathcal{W}_o(p, qz) &\sim (-2)^{(p-1)/2} \frac{\Gamma(\frac{p+1}{2})}{\Gamma(p+1)} e^{-(qz)^2/4} (qz)^p \left[ 1 - \frac{p(p-1)}{2(qz)^2} \right. \\ &\quad \left. + \frac{p(p-1)(p-2)(p-3)}{2 \cdot 4 (qz)^4} - \frac{p(p-1)(p-2)(p-3)(p-4)(p-5)}{2 \cdot 4 \cdot 6 (qz)^6} + \dots \right]. \end{aligned}$$

## Contour integrals

$$\begin{aligned}\mathcal{W}_e(\phi, qz) &= \frac{2^{\phi/2} e^{-(qz)^2/4}}{i \sin \phi \pi \Gamma(-\phi/2)} \int_{-\infty}^{0+} \cos qzt e^{-t^2/2} (-t)^{-\phi-1} dt, \\ \mathcal{W}_o(\phi, qz) &= \frac{2^{(\phi-1)/2} e^{-(qz)^2/4}}{i \sin \phi \pi \Gamma\left(\frac{1-\phi}{2}\right)} \int_{-\infty}^{0+} \sin qzt e^{-t^2/2} (-t)^{-\phi-1} dt.\end{aligned}$$

**Recurrence formulas.** For either the even or the odd functions,

$$\begin{aligned}\mathcal{W}(\phi + 1, qz) &= (qz) \mathcal{W}(\phi, qz) - \phi \mathcal{W}(\phi - 1, qz), \\ \frac{d\mathcal{W}(\phi, qz)}{dz} &= \phi \mathcal{W}(\phi - 1, qz) - \frac{qz}{2} \mathcal{W}(\phi, qz).\end{aligned}$$

**Orthogonality.** Weber functions are orthogonal on an arbitrary interval  $(a, b)$  if boundary conditions are of the form

$$\begin{cases} k_1 \mathcal{W}(\phi, qa) + k_2 \mathcal{W}'(\phi, qa) = 0, \\ k_3 \mathcal{W}(\phi, qb) + k_4 \mathcal{W}'(\phi, qb) = 0. \end{cases}$$

We have merely a special case of a Sturm-Liouville system (§ 7.07), with  $w = 1$  and  $\lambda = \phi$ . Therefore,

$$\int_a^b \mathcal{W}(\phi_m, qz) \mathcal{W}(\phi_n, qz) dz = 0, \quad m \neq n,$$

for either the odd or even functions.

An arbitrary function  $f(z)$  can be expanded in terms of Weber functions on any interval  $(a, b)$ :

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{W}(\phi_m, qz),$$

where

$$\begin{aligned}A_m &= \frac{1}{N_m} \int_a^b f(z) \mathcal{W}(\phi_m, qz) dz, \\ N_m &= \int_a^b [\mathcal{W}(\phi_m, qz)]^2 dz.\end{aligned}$$

Here all conditions apply to  $\mathcal{W}_e$ -functions, or  $\mathcal{W}_o$ -functions, or linear combinations of them.

On the infinite interval, since  $\mathcal{W}(\phi, \pm \infty) = 0$ ,

$$\int_{-\infty}^{\infty} \mathcal{W}(\phi_m, qz) \mathcal{W}(\phi_n, qz) dz = 0, \quad m \neq n.$$

Thus

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{W}(\phi_m, qz),$$

$$A_m = \frac{1}{N_m} \int_{-\infty}^{\infty} f(z) \mathcal{W}(\phi_m, qz) dz,$$

$$N_m = \int_{-\infty}^{\infty} [\mathcal{W}(\phi_m, qz)]^2 dz.$$

In particular if  $\phi = m$ ,

$$N_m = (2\pi)^{\frac{1}{2}} m!$$

## 7.09 BESSEL FUNCTIONS

**The wave equation.** The Bessel wave equation {26} is

$$\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 + q^2 - p^2/z^2) Z = 0. \quad (7.28)$$

The series solution was developed in § 7.04. The general solution of Eq. (7.28) may be written, for  $p \neq$  integer,

$$Z = A\mathcal{J}_p(\kappa, q, z) + B\mathcal{I}_{-p}(\kappa, q, z), \quad (7.33)$$

where the Bessel wave functions are

$$\mathcal{J}_p(\kappa, q, z) = (qz/2)^p \sum_{m=0}^{\infty} \frac{(-1)^m \Delta'_m(p) \cdot (z/2)^{2m}}{m! \Gamma(m+p+1)}, \quad (7.31)$$

$$\mathcal{I}_{-p}(\kappa, q, z) = (qz/2)^{-p} \sum_{m=0}^{\infty} \frac{(-1)^m \Delta'_m(-p) \cdot (z/2)^{2m}}{m! \Gamma(m-p+1)}, \quad (7.32)$$

and where the determinant  $\Delta'_m$  is given by Eq. (7.30). These series are valid everywhere in the finite complex plane.

If the differential equation is written

$$\frac{d^2Z}{dz^2} + \frac{1}{z} \frac{dZ}{dz} + (\kappa^2 z^2 - q^2 - p^2/z^2) Z = 0,$$

the general solution (for  $p \neq$  integer) is

$$Z = A\mathcal{J}_p(\kappa, q, iz) + B\mathcal{I}_{-p}(\kappa, q, iz).$$

If  $p = n$ , an integer,  $\mathcal{I}_{-p}$  is no longer independent of  $\mathcal{J}_p$  and the general solution of Eq. (7.28) is

$$Z = A\mathcal{J}_n(\kappa, q, z) + B\mathcal{Y}_n(\kappa, q, z). \quad (7.33a)$$

The Bessel wave function of the second kind may be written

$$\mathcal{Y}_n(\kappa, q, z) = \frac{2}{\pi q^{2n}} \left\{ [\gamma + \ln(qz/2)] \Delta'_n(-n) \mathcal{J}_n(\kappa, q, z) \right. \\ \left. - \frac{1}{2} (q/2)^n 2^{2n} z^{-n} \sum_{m=0}^{n-1} \frac{(n-m-1)! \Delta'_m(-n) \cdot (z/2)^{2m}}{m!} \right. \\ \left. + \frac{1}{2} (qz/2)^n \sum_{m=0}^{\infty} \frac{(-1)^m (z/2)^{2m}}{m! (m+n)!} \right. \\ \left. \times \left\{ 4 \sum_{l=1}^{m+n-1} (n-2l) M_{l(l+1)} - \Delta'_m(n) \Delta'_n(-n) \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \right\} \right\}. \quad (7.42)$$

For  $n = 0$ ,

$$\mathcal{Y}_0(\kappa, q, z) = \frac{2}{\pi} \left\{ [\gamma + \ln(qz/2)] \mathcal{J}_0(\kappa, q, z) \right. \\ \left. - \sum_{m=1}^{\infty} \frac{(-1)^m (z/2)^{2m}}{(m!)^2} \left[ 4 \sum_{l=1}^{m-1} l M_{l(l+1)}(0) + \Delta'_m(0) \sum_{l=1}^m \frac{1}{l} \right] \right\}. \quad (7.37)$$

**Degenerate cases.** The other Bessel functions [37] listed in § 7.01 are all degenerate forms of the foregoing. All series converge within a circle of infinite radius in the  $z$ -plane.

$$\mathcal{J}_p(qz) = \sum_{m=0}^{\infty} \frac{(-1)^m (qz/2)^{p+2m}}{m! \Gamma(m+p+1)}. \quad (7.34)$$

$$\mathcal{J}_{-p}(qz) = \sum_{m=0}^{\infty} \frac{(-1)^m (qz/2)^{-p+2m}}{m! \Gamma(m-p+1)}. \quad (7.34a)$$

These functions are shown in Fig. 7.01.

$$\begin{aligned} \mathcal{Y}_n(qz) = & \frac{2}{\pi} \left\{ [\gamma + \ln(qz/2)] \mathcal{J}_n(qz) - \frac{1}{2} (qz/2)^{-n} \sum_{m=0}^{n-1} \frac{(n-m-1)! (qz/2)^{2m}}{m!} \right. \\ & \left. - \frac{1}{2} (qz/2)^n \sum_{m=0}^{\infty} \frac{(-1)^m (qz/2)^{2m}}{m! (m+n)!} \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \right\}. \end{aligned} \quad (7.42a)$$

$$\mathcal{J}_p(iqz) = \sum_{m=0}^{\infty} \frac{(-1)^m (iqz/2)^{p+2m}}{m! \Gamma(m+p+1)}. \quad (7.34b)$$

$$\mathcal{J}_{-p}(iqz) = \sum_{m=0}^{\infty} \frac{(-1)^m (iqz/2)^{-p+2m}}{m! \Gamma(m-p+1)}. \quad (7.34c)$$

$$\begin{aligned} \mathcal{Y}_n(iqz) = & \frac{2}{\pi} \left\{ [\gamma + \ln(iqz/2)] \mathcal{J}_n(iqz) \right. \\ & - \frac{1}{2} (iqz/2)^{-n} \sum_{m=0}^{n-1} \frac{(n-m-1)! (iqz/2)^{2m}}{m!} \\ & \left. - \frac{1}{2} (iqz/2)^n \sum_{m=0}^{\infty} \frac{(-1)^m (iqz/2)^{2m}}{m! (m+n)!} \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \right\}. \end{aligned} \quad (7.42b)$$

$$\mathcal{J}_0(qz) = 1 - \frac{(qz/2)^2}{(1!)^2} + \frac{(qz/2)^4}{(2!)^2} - \frac{(qz/2)^6}{(3!)^2} + \cdots + (-1)^n \frac{(qz/2)^{2n}}{(n!)^2} + \cdots.$$

A plot of this function is given in Fig. 7.02.

$$\begin{aligned} \mathcal{Y}_0(qz) = & \frac{2}{\pi} [\gamma + \ln(qz/2)] \mathcal{J}_0(qz) \\ & + \frac{2}{\pi} \left\{ \frac{(qz/2)^2}{(1!)^2} - \frac{(qz/2)^4}{(2!)^2} (1 + \frac{1}{2}) + \frac{(qz/2)^6}{(3!)^2} (1 + \frac{1}{2} + \frac{1}{3}) - \cdots \right\}. \end{aligned}$$

See Fig. 7.03 for the  $\mathcal{Y}_0$ -function and Fig. 7.04 for  $\mathcal{J}_{\frac{1}{2}}$  and  $\mathcal{J}_{\frac{3}{2}}$ .

$$\mathcal{J}_0(iqz) = 1 + \frac{(qz/2)^2}{(1!)^2} + \frac{(qz/2)^4}{(2!)^2} + \frac{(qz/2)^6}{(3!)^2} + \cdots.$$

$$\begin{aligned} \mathcal{Y}_0(iqz) = & \frac{2}{\pi} \left[ \gamma + \ln(iqz/2) + i \frac{\pi}{2} \right] \mathcal{J}_0(iqz) \\ & - \frac{2}{\pi} \left\{ \frac{(qz/2)^2}{(1!)^2} + \frac{(qz/2)^2}{(2!)^2} (1 + \frac{1}{2}) + \frac{(qz/2)^6}{(3!)^2} (1 + \frac{1}{2} + \frac{1}{3}) + \cdots \right\}. \end{aligned}$$

$$\mathcal{J}_0(\varkappa, q, z) = 1 - \frac{(qz/2)^2}{(1!)^2} + \frac{(qz/2)^4}{(2!)^2} [1 - 4\varkappa^2/q^4] - \frac{(qz/2)^6}{(3!)^2} [1 - 20\varkappa^2/q^4] + \cdots.$$




Fig. 7.01 a and b. Bessel functions of the first kind, as functions of  $x$  and  $p$ . Note that for positive  $p$ , the Bessel functions are bounded; but for negative  $p$ , the functions become infinite at  $x=0$ . (From JAHNKE and EMDE, Funktionentafeln, by courtesy of the publishers, B. G. Teubner, Leipzig, and Dover Publications, New York City)






Fig. 7.02. Bessel functions  $J_0(x)$  and  $J_1(x)$ , showing the first few zeros

$$\begin{aligned}\mathcal{Y}_0(\kappa, q, z) = \frac{2}{\pi} & \left\{ [\gamma + \ln(qz/2)] \mathcal{I}_0(\kappa, q, z) \right. \\ & + \frac{(qz/2)^2}{(1!)^2} - \frac{(qz/2)^4}{(2!)^2} [4\kappa^2/q^4 + (1 - 4\kappa^2/q^4)(1 + \frac{1}{2})] \\ & \left. + \frac{(qz/2)^6}{(3!)^2} [12\kappa^2/q^4 + (1 - 20\kappa^2/q^4)(1 + \frac{1}{2} + \frac{1}{3})] - \dots \right\}.\end{aligned}$$

Fig. 7.03. Bessel functions  $\mathcal{Y}_0(x)$  and  $\mathcal{Y}_1(x)$ Fig. 7.04. Bessel functions  $\mathcal{J}_{1/2}(x)$  and  $\mathcal{J}_{3/2}(x)$ 

$$\mathcal{J}_0(\kappa, q, iz) = 1 + \frac{(qz/2)^2}{(1!)^2} + \frac{(qz/2)^4}{(2!)^2} [1 - 4\kappa^2/q^4] + \frac{(qz/2)^6}{(3!)^2} [1 - 20\kappa^2/q^4] + \dots.$$

$$\begin{aligned}\mathcal{Y}_0(\kappa, q, iz) = \frac{2}{\pi} & \left\{ [\gamma + \ln(izqz/2)] \mathcal{I}_0(\kappa, q, iz) \right. \\ & - \frac{(qz/2)^2}{(1!)^2} - \frac{(qz/2)^4}{(2!)^2} [4\kappa^2/q^4 + (1 - 4\kappa^2/q^4)(1 + \frac{1}{2})] \\ & \left. - \frac{(qz/2)^6}{(3!)^2} [12\kappa^2/q^4 + (1 - 20\kappa^2/q^4)(1 + \frac{1}{2} + \frac{1}{3})] - \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_p(\kappa, q, iz) = \frac{(iqz/2)^p}{\Gamma(p+1)} & \left\{ 1 + \frac{(qz/2)^2}{1!(p+1)} + \frac{(qz/2)^4[1 - 4(p+1)\kappa^2/q^4]}{2!(p+1)(p+2)} \right. \\ & + \frac{(qz/2)^6[1 - 4(3p+5)\kappa^2/q^4]}{3!(p+1)(p+2)(p+3)} + \dots \left. \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{-\rho}(\kappa, q, iz) &= \frac{(iqz/2)^{-\rho}}{\Gamma(-\rho+1)} \left\{ 1 - \frac{(qz/2)^2}{1!(\rho-1)} + \frac{(qz/2)^4 [1+4(\rho-1)\kappa^2/q^4]}{2!(\rho-1)(\rho-2)} \right. \\ &\quad \left. - \frac{(qz/2)^6 [1+4(3\rho-5)\kappa^2/q^4]}{3!(\rho-1)(\rho-2)(\rho-3)} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{Y}_n(\kappa, q, iz) &= \frac{2}{\pi q^{2n}} \left\{ [\gamma + \ln(iqz/2)] A'_n(-n) \mathcal{J}_n(\kappa, q, iz) \right. \\ &\quad - \frac{1}{2}(q/2)^n 2^{2n} (iz)^{-n} \sum_{m=0}^{n-1} \frac{(-1)^m (n-m-1)! A'_m(-n) (z/2)^{2m}}{m!} \\ &\quad + \frac{1}{2}(iqz/2)^n \sum_{m=0}^{\infty} \frac{(z/2)^{2m}}{m! (m+n)!} \\ &\quad \times \left. \left\{ 4 \sum_{l=1}^{m+n-1} (n-2l) M_{l(l+1)} - A'_m(n) A'_n(-n) \left[ \sum_{l=1}^m \frac{1}{l} + \sum_{l=1}^{m+n} \frac{1}{l} \right] \right\} \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{\frac{1}{2}}(\kappa, q, z) &= \frac{(qz/2)^{\frac{1}{2}}}{\Gamma(\frac{3}{2})} \\ &\times \left\{ 1 - \frac{(qz/2)^2}{1!(\frac{3}{2})} + \frac{(qz/2)^4 [1-3\kappa^2/q^4]}{2!(\frac{3}{2})(\frac{5}{2})} - \frac{(qz/2)^6 [1-13\kappa^2/q^4]}{3!(\frac{3}{2})(\frac{5}{2})(\frac{7}{2})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{-\frac{1}{2}}(\kappa, q, z) &= \frac{(qz/2)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})} \\ &\times \left\{ 1 - \frac{(qz/2)^2}{1!(\frac{1}{2})} + \frac{(qz/2)^4 [1-\kappa^2/q^4]}{2!(\frac{1}{2})(\frac{3}{2})} - \frac{(qz/2)^6 [1-7\kappa^2/q^4]}{3!(\frac{1}{2})(\frac{3}{2})(\frac{5}{2})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{\frac{1}{4}}(\kappa, q, iz) &= \frac{(iqz/2)^{\frac{1}{4}}}{\Gamma(\frac{3}{2})} \\ &\times \left\{ 1 + \frac{(qz/2)^2}{1!(\frac{3}{2})} + \frac{(qz/2)^4 [1-3\kappa^2/q^4]}{2!(\frac{3}{2})(\frac{5}{2})} + \frac{(qz/2)^6 [1-13\kappa^2/q^4]}{3!(\frac{3}{2})(\frac{5}{2})(\frac{7}{2})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{-\frac{1}{4}}(\kappa, q, iz) &= \frac{(iqz/2)^{-\frac{1}{4}}}{\Gamma(\frac{1}{2})} \\ &\times \left\{ 1 + \frac{(qz/2)^2}{1!(\frac{1}{2})} + \frac{(qz/2)^4 [1-\kappa^2/q^4]}{2!(\frac{1}{2})(\frac{3}{2})} + \frac{(qz/2)^6 [1-7\kappa^2/q^4]}{3!(\frac{1}{2})(\frac{3}{2})(\frac{5}{2})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{\frac{5}{4}}(qz) &= \frac{(qz/2)^{\frac{5}{4}}}{\Gamma(\frac{5}{4})} \\ &\times \left\{ 1 - \frac{(qz/2)^2}{1!(\frac{5}{4})} + \frac{(qz/2)^4}{2!(\frac{5}{4})(\frac{9}{4})} - \frac{(qz/2)^6}{3!(\frac{5}{4})(\frac{9}{4})(\frac{13}{4})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{-\frac{5}{4}}(qz) &= \frac{(qz/2)^{-\frac{5}{4}}}{\Gamma(\frac{3}{4})} \\ &\times \left\{ 1 - \frac{(qz/2)^2}{1!(\frac{3}{4})} + \frac{(qz/2)^4}{2!(\frac{3}{4})(\frac{7}{4})} - \frac{(qz/2)^6}{3!(\frac{3}{4})(\frac{7}{4})(\frac{11}{4})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{s+\frac{1}{2}}(qz) &= \frac{(qz/2)^{s+\frac{1}{2}}}{\Gamma(s+\frac{3}{2})} \left\{ 1 - \frac{(qz/2)^2}{1!(s+\frac{3}{2})} + \frac{(qz/2)^4}{2!(s+\frac{3}{2})(s+\frac{5}{2})} \right. \\ &\quad \left. - \frac{(qz/2)^6}{3!(s+\frac{3}{2})(s+\frac{5}{2})(s+\frac{7}{2})} + \dots \right\}.\end{aligned}$$

$$\begin{aligned}\mathcal{J}_{-(s+\frac{1}{2})}(qz) &= \frac{(qz/2)^{-(s+\frac{1}{2})}}{\Gamma(-s+\frac{1}{2})} \left\{ 1 + \frac{(qz/2)^2}{1!(s-\frac{1}{2})} + \frac{(qz/2)^4}{2!(s-\frac{1}{2})(s-\frac{3}{2})} \right. \\ &\quad \left. + \frac{(qz/2)^6}{3!(s-\frac{1}{2})(s-\frac{3}{2})(s-\frac{5}{2})} + \dots \right\}.\end{aligned}$$

*Hankel functions* (Bessel functions of the third kind) are linear combinations of Bessel functions of the first and second kinds:

$$\left. \begin{aligned} \mathcal{H}_p^{(1)}(qz) &= \mathcal{I}_p(qz) + i\mathcal{Y}_p(qz), \\ \mathcal{H}_p^{(2)}(qz) &= \mathcal{I}_p(qz) - i\mathcal{Y}_p(qz). \end{aligned} \right\} \quad (7.58)$$

The general solution of Eq. (7.28) may be written

$$Z = A\mathcal{H}_p^{(1)}(qz) + B\mathcal{H}_p^{(2)}(qz). \quad (7.59)$$

Other relations among Bessel functions may be written

$$\begin{aligned} \mathcal{I}_p(qz) &= \frac{1}{2} [\mathcal{H}_p^{(1)}(qz) + \mathcal{H}_p^{(2)}(qz)], \\ \mathcal{J}_{-p}(qz) &= \frac{1}{2} [e^{ip\pi}\mathcal{H}_p^{(1)}(qz) + e^{-ip\pi}\mathcal{H}_p^{(2)}(qz)], \\ \mathcal{Y}_p(qz) &= \frac{1}{\sin p\pi} [\mathcal{I}_p(qz) \cos p\pi - \mathcal{J}_{-p}(qz)] \\ &= \frac{1}{2i} [\mathcal{H}_p^{(1)}(qz) - \mathcal{H}_p^{(2)}(qz)], \\ \mathcal{Y}_{-p}(qz) &= \frac{1}{\sin p\pi} [\mathcal{I}_p(qz) - \mathcal{J}_{-p}(qz) \cos p\pi] \\ &= \frac{1}{2i} [e^{ip\pi}\mathcal{H}_p^{(1)}(qz) - e^{-ip\pi}\mathcal{H}_p^{(2)}(qz)], \\ \mathcal{H}_p^{(1)}(qz) &= \frac{1}{i \sin p\pi} [\mathcal{J}_{-p}(qz) - e^{-ip\pi}\mathcal{I}_p(qz)], \\ \mathcal{H}_p^{(2)}(qz) &= \frac{1}{i \sin p\pi} [e^{ip\pi}\mathcal{J}_p(qz) - \mathcal{J}_{-p}(qz)]. \end{aligned}$$

### Bessel functions as definite integrals

$$\begin{aligned} \mathcal{I}_p(qz) &= \frac{(qz/2)^p}{\Gamma(p+\frac{1}{2}) \Gamma(\frac{1}{2})} \int_0^\pi \cos(qz \cos u) \sin^{2p} u \, du, \\ \mathcal{J}_p(qz) &= \frac{2(qz/2)^p}{\Gamma(p+\frac{1}{2}) \Gamma(\frac{1}{2})} \int_0^1 (1-t^2)^{p-\frac{1}{2}} \cos(qzt) \, dt, \\ \mathcal{J}_p(qz) &= \frac{(qz/2)^p}{\Gamma(p+\frac{1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 (1-t^2)^{p-\frac{1}{2}} \cos(qzt) \, dt, \\ \mathcal{J}_p(qz) &= \frac{(qz/2)^p}{\Gamma(p+\frac{1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 (1-t^2)^{p-\frac{1}{2}} e^{izt} \, dt, \\ \mathcal{J}_p(qz) &= \frac{(qz/2)^p}{\Gamma(p+\frac{1}{2}) \Gamma(\frac{1}{2})} \int_0^\pi e^{iqz \cos \theta} \sin^{2p} \theta \, d\theta, \\ \mathcal{J}_{n+\frac{1}{2}}(qz) &= (-i)^n \left( \frac{qz}{2\pi} \right)^{\frac{1}{2}} \int_0^\pi e^{iqz \cos \theta} P_n(\cos \theta) \sin \theta \, d\theta. \end{aligned}$$

### Asymptotic expansions

$$\begin{aligned}\mathcal{I}_p(qz) &\cong \left(\frac{2}{\pi qz}\right)^{\frac{1}{2}} \left\{ \cos(qz - p\pi/2 - \pi/4) \sum_{m=0}^{\infty} \frac{(-1)^m \Gamma(p+2m+\frac{1}{2})}{(2qz)^{2m} m! \Gamma(p-2m+\frac{1}{2})} \right. \\ &\quad \left. - \sin(qz - p\pi/2 - \pi/4) \sum_{m=0}^{\infty} \frac{(-1)^m \Gamma(p+2m+\frac{3}{2})}{(2qz)^{2m+1} (m+1)! \Gamma(p-2m-\frac{1}{2})} \right\}, \\ \mathcal{Y}_p(qz) &\cong \left(\frac{2}{\pi qz}\right)^{\frac{1}{2}} \left\{ \sin(qz - p\pi/2 - \pi/4) \sum_{m=0}^{\infty} \frac{(-1)^m \Gamma(p+2m+\frac{1}{2})}{(2qz)^{2m} m! \Gamma(p-2m+\frac{1}{2})} \right. \\ &\quad \left. + \cos(qz - p\pi/2 - \pi/4) \sum_{m=0}^{\infty} \frac{(-1)^m \Gamma(p+2m+\frac{3}{2})}{(2qz)^{2m+1} (m+1)! \Gamma(p-2m-\frac{1}{2})} \right\}, \\ \mathcal{H}_p^{(1)}(qz) &\cong \left(\frac{2}{\pi qz}\right)^{\frac{1}{2}} e^{i(qz-p\pi/2-\pi/4)} \sum_{m=0}^{\infty} \frac{(-1)^m \Gamma(p+m+\frac{1}{2})}{(2izqz)^m m! \Gamma(p-m+\frac{1}{2})}, \\ \mathcal{H}_p^{(2)}(qz) &\cong \left(\frac{2}{\pi qz}\right)^{\frac{1}{2}} e^{-i(qz-p\pi/2-\pi/4)} \sum_{m=0}^{\infty} \frac{\Gamma(p+m+\frac{1}{2})}{(2izqz)^m m! \Gamma(p-m+\frac{1}{2})}.\end{aligned}$$

**Recursion formulas.** The same recursion formulas apply to all Bessel functions [38]. The same is true for formulas of differentiation and integration. Let  $\mathcal{Z}$  be a Bessel function of first, second, or third kind. Then

$$\begin{aligned}\mathcal{Z}_{p-1}(qz) + \mathcal{Z}_{p+1}(qz) &= \frac{2p}{qz} \mathcal{Z}_p(qz), \\ \mathcal{Z}_{p-1}(qz) - \mathcal{Z}_{p+1}(qz) &= \frac{2}{q} \frac{d}{dz} \mathcal{Z}_p(qz), \\ z \frac{d}{dz} \mathcal{Z}_p(qz) + p \mathcal{Z}_p(qz) &= qz \mathcal{Z}_{p-1}(qz), \\ z \frac{d}{dz} \mathcal{Z}_p(qz) - p \mathcal{Z}_p(qz) &= -qz \mathcal{Z}_{p+1}(qz), \\ \mathcal{Z}_p(qz) \mathcal{Z}_{1-p}(qz) + \mathcal{Z}_{-p}(qz) \mathcal{Z}_{p-1}(qz) &= \frac{2 \sin p\pi}{\pi qz}.\end{aligned}$$

**Differentiation.** Let  $\mathcal{Z}_p(qz)$  be a Bessel function of first, second, or third kind, or any linear combination of these functions. Then

$$\begin{aligned}\frac{d}{dz} [\mathcal{Z}_p(qz)] &= -\frac{p}{q} \mathcal{Z}_p(qz) + q \mathcal{Z}_{p-1}(qz) \\ &= \frac{p}{q} \mathcal{Z}_p(qz) - q \mathcal{Z}_{p+1}(qz) \\ &= \frac{q}{2} [\mathcal{Z}_{p-1}(qz) - \mathcal{Z}_{p+1}(qz)].\end{aligned}$$

$$\frac{d}{dz} [z^p \mathcal{Z}_p(qz)] = qz^p \mathcal{Z}_{p-1}(qz).$$

$$\frac{d}{dz} [z^{-p} \mathcal{Z}_p(qz)] = -qz^{-p} \mathcal{Z}_{p+1}(qz).$$

$$\frac{d}{dz} [z^{p/2} \mathcal{Z}_p[(qz)^{\frac{1}{2}}]] = \frac{q^{\frac{1}{2}}}{2} z^{(p-1)/2} \mathcal{Z}_{p-1}[(qz)^{\frac{1}{2}}].$$

$$\frac{d}{dz} [z^{-p/2} \mathcal{L}_p[(qz)^{\frac{1}{2}}]] = -\frac{q^{\frac{1}{2}}}{2} z^{-(p+1)/2} \mathcal{L}_{p+1}[(qz)^{\frac{1}{2}}].$$

$$\frac{d^2}{dz^2} [\mathcal{L}_p(qz)] = \left[ \frac{p(p-1)}{z^2} - q^2 \right] \mathcal{L}_p(qz) + \frac{q}{z} \mathcal{L}_{p+1}(qz).$$

$$\left( \frac{1}{qz} \frac{d}{dz} \right)^m [z^p \mathcal{L}_p(qz)] = (qz)^{p-m} \mathcal{L}_{p-m}(qz).$$

$$\left( \frac{1}{qz} \frac{d}{dz} \right)^m [z^{-p} \mathcal{L}_p(qz)] = (-1)^m (qz)^{-(p+m)} \mathcal{L}_{p+m}(qz).$$

$$\frac{d}{dz} [\mathcal{L}_0(qz)] = -q \mathcal{L}_1(qz).$$

$$\frac{d}{dz} [\mathcal{L}_1(qz)] = q \mathcal{L}_0(qz) - \frac{1}{z} \mathcal{L}_2(qz).$$

$$\frac{d^2}{dz^2} [\mathcal{L}_0(qz)] = -q^2 \mathcal{L}_0(qz) + \frac{q}{z} \mathcal{L}_1(qz).$$

$$\frac{d^2}{dz^2} [\mathcal{L}_1(qz)] = -q^2 \mathcal{L}_1(qz) + \frac{q}{z} \mathcal{L}_2(qz).$$

### Integration

$$\int z^{p+1} \mathcal{L}_p(qz) dz = \frac{z^{p+1}}{q} \mathcal{L}_{p+1}(qz).$$

$$\int z^{-p+1} \mathcal{L}_p(qz) dz = -\frac{z^{-p+1}}{q} \mathcal{L}_{p-1}(qz).$$

$$\int z [\mathcal{L}_p(qz)]^2 dz = \frac{z^2}{2} \{ [\mathcal{L}_p(qz)]^2 - \mathcal{L}_{p-1}(qz) \mathcal{L}_{p+1}(qz) \}.$$

$$\int z [\mathcal{L}_0(qz)]^2 dz = \frac{z^2}{2} \{ [\mathcal{L}_0(qz)]^2 + [\mathcal{L}_1(qz)]^2 \}.$$

$$\int \mathcal{L}_1(qz) dz = -\frac{1}{q} \mathcal{L}_0(qz).$$

$$\int z \mathcal{L}_0(qz) dz = \frac{z}{q} \mathcal{L}_1(qz).$$

$$\int z \mathcal{L}_p(\alpha z) \mathcal{L}_p(\beta z) dz = \frac{z}{\alpha^2 - \beta^2} [\beta \mathcal{L}_p(\alpha z) \mathcal{L}_{p-1}(\beta z) - \alpha \mathcal{L}_{p-1}(\alpha z) \mathcal{L}_p(\beta z)].$$

$$\int z \mathcal{L}_0(\alpha z) \mathcal{L}_0(\beta z) dz = \frac{z}{\alpha^2 - \beta^2} [\beta \mathcal{L}_0(\alpha z) \mathcal{L}_{-1}(\beta z) - \alpha \mathcal{L}_{-1}(\alpha z) \mathcal{L}_0(\beta z)].$$

$$\int z \mathcal{L}_1(\alpha z) \mathcal{L}_1(\beta z) dz = \frac{z}{\alpha^2 - \beta^2} [\beta \mathcal{L}_1(\alpha z) \mathcal{L}_0(\beta z) - \alpha \mathcal{L}_0(\alpha z) \mathcal{L}_1(\beta z)].$$

**Orthogonality.** Bessel wave functions are orthogonal on the interval  $(a, b)$  with respect to the weighting function  $z$ :

$$\int_a^b z \mathcal{L}_p(\kappa, q_m, z) \cdot \mathcal{L}_p(\kappa, q_n, z) dz = 0, \quad m \neq n, \quad (7.60)$$

where boundary conditions of the form

$$k_1 \mathcal{L}_p(\kappa, q_m, z) + k_2 \mathcal{L}'_p(\kappa, q_m, z) = 0$$

apply at  $z = a$  and at  $z = b$  (Sturm-Liouville conditions, § 7.07).

An arbitrary function  $f(z)$  can be expanded in a series of Bessel wave functions:

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{L}_p(\varkappa, q_m, z), \quad (7.61)$$

where

$$A_m = \frac{1}{N_m} \int_a^b z f(z) \mathcal{L}_p(\varkappa, q_m, z) dz \quad (7.62)$$

and

$$N_m = \int_a^b z [\mathcal{L}_p(\varkappa, q_m, z)]^2 dz.$$

If  $\varkappa = 0$ ,

$$\left. \begin{aligned} N_m &= \frac{b^2}{2} \{[\mathcal{L}_p(q_m b)]^2 - \mathcal{L}_{p-1}(q_m b) \cdot \mathcal{L}_{p+1}(q_m b)\} \\ &\quad - \frac{a^2}{2} \{[\mathcal{L}_p(q_m a)]^2 - \mathcal{L}_{p-1}(q_m a) \cdot \mathcal{L}_{p+1}(q_m a)\}. \end{aligned} \right\} \quad (7.63)$$

For  $p = 0$ , the coefficients of the expansion, Eq. (7.61), are

$$A_m = \frac{1}{N_m} \int_a^b z f(z) \cdot \mathcal{L}_0(\varkappa, q_m, z) dz$$

where

$$N_m = \int_a^b z [\mathcal{L}_0(\varkappa, q_m, z)]^2 dz.$$

If  $\varkappa = 0$ ,

$$\left. \begin{aligned} N_m &= \frac{b^2}{2} \{[\mathcal{L}_0(q_m b)]^2 + [\mathcal{L}_1(q_m b)]^2\} \\ &\quad - \frac{a^2}{2} \{[\mathcal{L}_0(q_m a)]^2 + [\mathcal{L}_1(q_m a)]^2\}. \end{aligned} \right\} \quad (7.63 \text{ a})$$

For boundary conditions  $\mathcal{L}_0(q_m a) = 0$  and  $\mathcal{L}_0(q_m b) = 0$ ,  $\mathcal{L}_0$  may be taken as a linear combination of  $\mathcal{I}_0$  and  $\mathcal{Y}_0$ , so adjusted that its zeros are at  $a$  and  $b$ . Then Eq. (7.63 a) becomes

$$N_m = \frac{b^2}{2} [\mathcal{L}_1(q_m b)]^2 - \frac{a^2}{2} [\mathcal{L}_1(q_m a)]^2. \quad (7.63 \text{ b})$$

In the special case of  $a = 0$ , Bessel wave functions of the second kind are not applicable, and the expansion is in terms of  $\mathcal{J}_p(\varkappa, q_m, z)$ . Then, for  $p = 0$ ,

$$\begin{aligned} f(z) &= \sum_{m=0}^{\infty} A_m \mathcal{J}_0(\varkappa, q_m, z), \\ A_m &= \frac{1}{N_m} \int_a^b z f(z) \mathcal{J}_0(\varkappa, q_m, z) dz, \end{aligned}$$

where

$$N_m = \int_0^b z [\mathcal{J}_0(\varkappa, q_m, z)]^2 dz.$$

If  $\varkappa = 0$ ,

$$N_m = \frac{b^2}{2} [\mathcal{J}_1(q_m b)]^2,$$

where  $q_m$  are determined by the relation,

$$\mathcal{J}_0(q_m b) = 0.$$

### 7.10 BAER FUNCTIONS

**The wave functions.** The Baer wave equation {114} may be written [39]

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-b} + \frac{1}{z-c} \right] \frac{dZ}{dz} + \left[ \frac{\kappa^2 z^2 - p(p+1)z - q(b+c)}{(z-b)(z-c)} \right] Z = 0. \quad (7.64)$$

For a solution about the origin, take

$$Z = z^\beta \sum_{j=0}^{\infty} C_j z^j.$$

Applying the method of § 7.03, we obtain

$$A_0 = 0,$$

$$A_1 = -\frac{b+c}{2bc},$$

$$A_2 = -\frac{(b^2+c^2)}{2(bc)^2},$$

$$A_j = \frac{1}{bc} [A_{j-1}(b+c) - A_{j-2}] \quad \text{for } j > 2;$$

$$B_0 = B_1 = 0,$$

$$B_2 = -\frac{q(b+c)}{bc},$$

$$B_3 = -\frac{1}{bc} \left[ p(p+1) + q \frac{(b+c)^2}{bc} \right],$$

$$B_4 = \frac{1}{bc} \left[ \kappa^2 - p(p+1) \left( \frac{b+c}{bc} \right) + q \left( \frac{b+c}{bc} \right) \left[ 1 - \frac{(b+c)^2}{bc} \right] \right],$$

$$B_j = \frac{1}{bc} [B_{j-1}(b+c) - B_{j-2}] \quad \text{for } j > 4.$$

The indicial equation is

$$\beta(\beta-1) = 0,$$

so  $b_1 = 1$ ,  $b_2 = 0$ . For the first root, the determinant is

$$A_j(1) = \begin{vmatrix} A_1 & 2 \cdot 1 & 0 & 0 & \dots & 0 \\ A_2 + B_2 & 2A_1 & 3 \cdot 2 & 0 & \dots & 0 \\ A_3 + B_3 & 2A_2 + B_2 & 3A_1 & 4 \cdot 3 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ A_j + B_j & \dots & \dots & \dots & \dots & j(j-1) \\ & & & & & jA_1 \end{vmatrix}. \quad (7.65)$$

Thus,

$$C_j(1) = \frac{(-1)^j C_0 A_j(1)}{[1 \cdot 2][2 \cdot 3][3 \cdot 4] \dots [j(j+1)]}$$

and the first solution is

$$Z_1 = C_0 z \sum_{j=0}^{\infty} \frac{(-1)^j A_j(1) z^j}{j!(j+1)!}. \quad (7.66)$$

For  $\beta = b_2 = 0$ ,

$$\Delta_j(0) = \begin{vmatrix} 0 & f_0(1) & 0 & 0 & \dots & 0 \\ B_2 & A_1 & 2 \cdot 1 & 0 & \dots & 0 \\ B_3 & A_2 + B_2 & 2A_1 & 3 \cdot 2 & \dots & 0 \\ B_4 & A_3 + B_3 & 2A_2 + B_2 & 3A_1 & 4 \cdot 3 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & (j-1)(j-2) \\ B_j & \dots & \dots & \dots & \dots & (j-1)A_1 \end{vmatrix},$$

and

$$C_j(0) = \frac{(-1)^j C_0 \Delta_j(0)}{f_0(1) [2 \cdot 1] [3 \cdot 2] \dots [j(j-1)]},$$

or

$$Z_2 = C_0 \left[ 1 + \sum_{m=1}^{\infty} \frac{(-1)^{m+1} \Delta'_m(0) z^{m+1}}{m! (m+1)!} \right], \quad (7.67)$$

where

$$\Delta'_m(0) = \begin{vmatrix} B_2 & 2 \cdot 1 & 0 & 0 & 0 & \dots & 0 \\ B_3 & 2A_1 & 3 \cdot 2 & 0 & 0 & \dots & 0 \\ B_4 & 2A_2 + B_2 & 3A_1 & 4 \cdot 3 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & m(m-1) & \dots \\ B_{m+1} & \dots & \dots & \dots & \dots & m A_1 & \dots \end{vmatrix}. \quad (7.68)$$

If  $C_0 = 1$ , the general solution of Eq. (7.64) may be written

$$Z = A \mathcal{B}_\beta^q(\kappa, z) + B \mathcal{C}_\beta^q(\kappa, z), \quad (7.69)$$

where

$$\left. \begin{aligned} \mathcal{B}_\beta^q(\kappa, z) &= z \sum_{m=0}^{\infty} \frac{(-1)^m \Delta'_m(1) z^m}{m! (m+1)!}, \\ \mathcal{C}_\beta^q(\kappa, z) &= 1 + \sum_{m=1}^{\infty} \frac{(-1)^{m+1} \Delta'_m(0) z^{m+1}}{m! (m+1)!}. \end{aligned} \right\} \quad (7.70)$$

These expressions converge within a circle about the origin with radius  $b$  ( $b < c$ ).

**Expansion about another singularity.** For an expansion about the point  $z_0 = b$ ,

$$Z = (z - b)^\beta \sum_{j=0}^{\infty} C_j (z - b)^j,$$

and

$$A_0 = \frac{1}{2},$$

$$A_1 = \frac{1}{2(b-c)},$$

$$A_j = -\frac{A_{j-1}}{b-c} \quad \text{for } j > 1;$$

$$B_0 = 0,$$

$$B_1 = -\frac{1}{b-c} [b^2 \kappa^2 - p(p+1)b - q(b+c)],$$

$$B_2 = \frac{1}{(b-c)^2} [b \kappa^2 - 2bc \kappa^2 + p(p+1)c + q(b+c)],$$

$$B_3 = \frac{1}{(b-c)^3} [c \kappa^2 - p(p+1)c - q(b+c)],$$

$$B_j = -\frac{B_{j-1}}{b-c} \quad \text{for } j > 3.$$

The indicial equation is

$$\beta(\beta - \frac{1}{2}) = 0,$$

so  $b_1 = \frac{1}{2}$ ,  $b_2 = 0$ . For  $\beta = b_1$ ,

$$A_j(\frac{1}{2}) = \begin{vmatrix} A_1/2 + B_1 & \frac{3}{2} & 0 & 0 & \dots & 0 \\ A_2/2 + B_2 & \frac{3}{2}A_1 + B_1 & 5 & 0 & \dots & 0 \\ A_3/2 + B_3 & \frac{3}{2}A_2 + B_2 & \frac{5}{2}A_1 + B_1 & \frac{21}{2} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & (j - \frac{1}{2})(j - 1) & \dots \\ A_j/2 + B_j & \dots & \dots & \dots & (j - \frac{1}{2})A_1 + B_1 & \dots \end{vmatrix}. \quad (7.71)$$

For  $\beta = b_2 = 0$ ,

$$A_j(0) = \begin{vmatrix} B_1 & \frac{1}{2} & 0 & 0 & \dots & 0 \\ B_2 & A_1 + B_1 & 3 & 0 & \dots & 0 \\ B_3 & A_2 + B_2 & 2A_1 + B_1 & \frac{15}{2} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & (j - 1)(j - \frac{3}{2}) & \dots \\ B_j & \dots & \dots & \dots & (j - 1)A_1 + B_1 & \dots \end{vmatrix}. \quad (7.72)$$

Thus the solutions are

$$\left. \begin{aligned} Z_1 &= C_0(z - b) \sum_{i=0}^{\infty} \frac{(-1)^i 2^{2i} A_j(\frac{1}{2}) \cdot (z - b)^i}{(2i + 1)!}, \\ Z_2 &= C_0 \sum_{i=0}^{\infty} \frac{(-1)^i 2^{2i} A_j(0) (z - b)^i}{(2i)!}. \end{aligned} \right\} \quad (7.73)$$

These series represent the same functions as Eq. (7.70) but over a different region in the  $z$ -plane. Accordingly, Eq. (7.73) with  $C_0 = 1$  represents the Baer functions  $\mathcal{B}_p^q(\kappa, z)$  and  $\mathcal{C}_p^q(\kappa, z)$  about the point  $z = b$  and extending out to the nearest singularity.

**Orthogonality.** According to § 7.07, a set of Baer functions with appropriate boundary conditions form an orthogonal set (with weighting function) on the arbitrary interval  $(a, b)$ . Thus, from Table 7.01,

$$\int_a^b \frac{z}{u} \mathcal{B}_{p_m}^q(\kappa, z) \mathcal{B}_{p_n}^q(\kappa, z) dz = 0, \quad m \neq n,$$

where  $u = [(z - b)(z - c)]^{\frac{1}{2}}$ . The same relation holds for  $\mathcal{C}_p^q(\kappa, z)$ ; and a similar relation holds if  $q$  is varied instead of  $p$  (Table 7.01).

An arbitrary function  $f(z)$  can be expanded in a series of Baer functions. For instance,

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{B}_{p_m}^q(\kappa, z),$$

where the coefficients are

$$\begin{aligned} A_m &= \frac{1}{N_m} \int_a^b \frac{z}{u} f(z) \mathcal{B}_{p_m}^q(\kappa, z) dz, \\ N_m &= \int_a^b \frac{z}{u} [\mathcal{B}_{p_m}^q(\kappa, z)]^2 dz. \end{aligned}$$

## 7.11 MATHIEU FUNCTIONS

The Baer equation {413} may be written

$$\frac{d^2Z}{d\xi^2} + \frac{1}{2} \left[ \frac{1}{\xi - b} + \frac{1}{\xi - c} \right] \frac{dZ}{d\xi} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 \xi}{(\xi - b)(\xi - c)} \right] Z = 0.$$

A special case of the Baer equation is the *Mathieu equation* [40]. Let  $b = 0$ ,  $c = 1$ ,  $\bar{A}_0 = -(2q + \lambda)$ ,  $\bar{A}_1 = 4q$ . Then, introducing the new variable defined by  $\zeta = \cos^2 z$ , we obtain the Mathieu equation,

$$\frac{d^2Z}{dz^2} + (\lambda - 2q \cos 2z) Z = 0, \quad (7.74)$$

where  $\lambda$  and  $q$  are arbitrary parameters [41].

**Periodic solutions.** For the special case of  $q = 0$  and  $\lambda = p^2$ , Eq. (7.74) reduces to the familiar {04}:

$$\frac{d^2Z}{dz^2} + p^2 Z = 0,$$

whose solutions are

$$Z = \begin{pmatrix} \cos \\ \sin \end{pmatrix} pz.$$

Periodic solutions are obtainable also for other values of  $q$ , but only when certain relations hold among  $\lambda$ ,  $p$ , and  $q$ . These periodic solutions are called *Mathieu functions of the first kind* and are denoted by

$$\text{ce}_p(q, z), \quad \text{se}_p(q, z),$$

corresponding to  $\cos pz$  and  $\sin pz$ .

Define the function  $\text{ce}_p(q, z)$  as a periodic function that reduces to  $\cos pz$  when  $q = 0$ . Let [42]

$$\begin{cases} Z = \text{ce}_p(q, z) = \cos pz + \sum_{j=1}^{\infty} q^j C_j(z), \\ \lambda = p^2 + \sum_{j=1}^{\infty} \alpha_j q^j. \end{cases}$$

Substitution into Eq. (7.74) yields the series

$$\left. \begin{aligned} \text{ce}_p(q, z) &= \cos pz - \frac{q}{4} \left[ \frac{\cos(p+2)z}{p+1} - \frac{\cos(p-2)z}{p-1} \right] \\ &\quad + \frac{q^2}{32} \left[ \frac{\cos(p+4)z}{(p+1)(p+2)} + \frac{\cos(p-4)z}{(p-1)(p-2)} \right] \\ &\quad - \frac{q^3}{128} \left[ \frac{(p^2+4p+7)\cos(p+2)z}{(p-1)(p+1)^3(p+2)} - \frac{(p^2-4p+7)\cos(p-2)z}{(p+1)(p-1)^3(p-2)} \right] \\ &\quad + \frac{\cos(p+6)z}{3(p+1)(p+2)(p+3)} - \frac{\cos(p-6)z}{3(p-1)(p-2)(p-3)} + \dots, \end{aligned} \right\} \quad (7.75)$$

and

$$\left. \begin{aligned} \lambda &= p^2 + \frac{q^2}{2(p^2-1)} \\ &\quad + \frac{(5p^2+7)q^4}{32(p^2-1)^3(p^2-4)} + \frac{(9p^4+58p^2+29)q^6}{64(p^2-1)^5(p^2-4)(p^2-9)} + \dots. \end{aligned} \right\} \quad (7.76)$$

Similarly,

$$\left. \begin{aligned} \text{se}_p(q, z) &= \sin pz - \frac{q}{4} \left[ \frac{\sin(p+2)z}{p+1} - \frac{\sin(p-2)z}{p-1} \right] \\ &\quad + \frac{q^2}{32} \left[ \frac{\sin(p+4)z}{(p+1)(p+2)} + \frac{\sin(p-4)z}{(p-1)(p-2)} \right] - \dots, \end{aligned} \right\} \quad (7.77)$$

where  $\lambda$  is again given by Eq. (7.76). The series of Eqs. (7.75), (7.76), and (7.77) fail if  $p$  is an integer other than zero.

For  $p \neq$  integer,  $\text{ce}_p(q, z)$  and  $\text{se}_p(q, z)$  are independent functions, and the general solution of Eq. (7.74) is

$$Z = A \text{ce}_p(q, z) + B \text{se}_p(q, z), \quad (7.78)$$

with  $p$  related to  $\lambda$  and  $q$  as in Eq. (7.76). For the more familiar case of  $p$  an integer, only one solution can be periodic and Eq. (7.78) cannot be used as the general solution [43].

**Integer  $p$ .** Now consider the special case where  $p = m$ , an integer [44]. If  $p = 0$ , Eqs. (7.75) and (7.76) reduce to

$$\begin{aligned} \text{ce}_0(q, z) &= 1 - \frac{q}{2} \cos 2z + \frac{q^2}{32} \cos 4z \\ &\quad - \frac{q^3}{128} \left( \frac{1}{9} \cos 6z - 7 \cos 2z \right) + \frac{q^4}{73728} (\cos 8z - 320 \cos 4z) + \dots \end{aligned}$$

with

$$\lambda = -\frac{q^2}{2} + \frac{7q^4}{128} - \frac{29q^6}{2304} + \dots$$

Improved methods of computation have been developed by INCE and by GOLDSTEIN [41]. For  $m = 0$ , a periodic solution requires the above relation between  $\lambda$  and  $q$ . However, the second solution obtained from Eq. (7.77) vanishes identically, so a non-periodic second solution must be found.

For  $p = 1$ , neither Eq. (7.75) nor Eq. (7.77) yields a convergent solution. However, somewhat similar periodic solutions, using an entirely different expression for  $\lambda$  can be found. These are Mathieu functions for integer values.

$$\begin{aligned} \text{ce}_1(q, z) &= \cos z - \frac{q}{8} \cos 3z + \frac{q^2}{64} \left( \frac{1}{3} \cos 5z - \cos 3z \right) \\ &\quad - \frac{q^3}{512} \left( \frac{1}{18} \cos 7z - \frac{4}{9} \cos 5z + \frac{1}{3} \cos 3z \right) \\ &\quad + \frac{q^4}{4096} \left( \frac{1}{180} \cos 9z - \frac{1}{12} \cos 7z + \frac{1}{6} \cos 5z + \frac{11}{9} \cos 3z \right) - \dots \end{aligned}$$

with

$$\lambda = 1 + q - \frac{q^2}{8} - \frac{q^3}{64} - \frac{q^4}{1536} + \frac{11q^5}{36864} + \dots$$

The interesting way in which a cosine changes into other Mathieu functions is indicated in Fig. 7.05, where  $\text{ce}_1$  is plotted as a function of  $q$  and  $z$ .

$$\begin{aligned} \text{ce}_2(q, z) &= \cos 2z - \frac{q}{8} \left( \frac{2}{3} \cos 4z - 2 \right) + \frac{q^2}{384} \cos 6z \\ &\quad - \frac{q^3}{512} \left( \frac{1}{45} \cos 8z + \frac{43}{27} \cos 4z + \frac{40}{3} \right) \\ &\quad + \frac{q^4}{4096} \left( \frac{1}{540} \cos 10z + \frac{293}{540} \cos 6z \right) - \dots \end{aligned}$$

with

$$\lambda = 4 + \frac{5}{12} q^2 - \frac{763}{13824} q^4 + \frac{1002401}{79626240} q^6 - \dots$$

The  $\lambda$ 's for the ce-functions are indicated by the curves marked  $a_0, a_1, \dots$  in Fig. 7.06. Corresponding curves marked  $b_1, b_2, b_3, \dots$  refer to the values of  $\lambda$




Fig. 7.05. The Mathieu function  $ce_1(qz)$ . For  $q=0$ , the function is an ordinary cosine, but as  $q$  increases, the cosine curve is progressively distorted

for the se-functions:

$$\begin{aligned} se_1(q, z) = & \sin z - \frac{q}{8} \sin 3z + \frac{q^2}{64} \left( \frac{1}{3} \sin 5z + \sin 3z \right) \\ & - \frac{q^3}{512} \left( \frac{1}{18} \sin 7z + \frac{4}{9} \sin 5z + \frac{1}{3} \sin 3z \right) \\ & + \frac{q^4}{4096} \left( \frac{1}{180} \sin 9z + \frac{1}{12} \sin 7z + \frac{1}{6} \sin 5z - \frac{11}{9} \sin 3z \right) - \dots \end{aligned}$$

with

$$\lambda = 1 - q - \frac{q^2}{8} + \frac{q^3}{64} - \frac{q^4}{1536} - \frac{11q^5}{36864} + \dots$$

$$\begin{aligned} se_2(q, z) = & \sin 2z - \frac{q}{12} \sin 4z + \frac{q^2}{384} \sin 6z - \frac{q^3}{512} \left( \frac{1}{45} \sin 8z - \frac{5}{27} \sin 4z \right) \\ & + \frac{q^4}{4096} \left( \frac{1}{540} \sin 10z - \frac{37}{540} \sin 6z \right) - \dots \end{aligned}$$

with

$$\lambda = 4 - \frac{q^2}{12} + \frac{5}{13824} q^4 - \dots$$

Since  $ce_m$  and  $se_m$  do not correspond to the same value of  $\lambda$ , their linear combination, Eq. (7.78), does not constitute the general solution of the Mathieu equation. Either  $ce_m$  or  $se_m$  may be used if a series of such terms fits the boundary conditions. If neither will satisfy the boundary conditions, one must introduce Mathieu functions of the second kind which are non-periodic.

We have seen that periodic solutions of the Mathieu equation occur only for certain relations between  $\lambda$ ,  $m$  and  $q$ , as shown by the heavy curves of Fig. 7.06. For all other conditions, the solutions are non-periodic. The shaded areas indicate non-periodic solutions which tend to finite values as  $z \rightarrow \infty$ ; the unshaded areas indicate non-periodic solutions which approach  $\infty$  as  $z \rightarrow \infty$ .




Fig. 7.06. Relation between  $\lambda$  and  $q$  for Mathieu functions. Periodic solutions are obtained only for points located on the curves. All other solutions are non-periodic, the shaded areas representing stable solutions and the unshaded areas unstable solutions

**Functions of the second kind.** Corresponding to Mathieu functions of the first kind, we have non-periodic functions of the second kind:

$$\left. \begin{aligned} \text{fe}_m(q, z) &= \text{ce}_m(q, z) \int_0^z \frac{dz}{[\text{ce}_m(q, z)]^2}, \\ \text{ge}_m(q, z) &= \text{se}_m(q, z) \int_0^z \frac{dz}{[\text{se}_m(q, z)]^2}. \end{aligned} \right\} \quad (7.79)$$

**Orthogonality.** If  $m$  and  $n$  are positive integers,

$$\int_0^{2\pi} \text{ce}_m(q, z) \text{ce}_n(q, z) dz = 0, \quad m \neq n;$$

$$\int_0^{2\pi} \text{se}_m(q, z) \text{se}_n(q, z) dz = 0, \quad m \neq n;$$

$$\int_0^{2\pi} \text{ce}_m(q, z) \text{se}_n(q, z) dz = 0,$$

$$\int_0^{2\pi} \text{ce}_m^2(q, z) dz = \int_0^{2\pi} \text{se}_m^2(q, z) dz = \pi, \quad m > 0.$$

Thus an arbitrary function  $f(z)$  can be expanded in a series of Mathieu functions:

$$f(z) = \sum_{m=0}^{\infty} A_m \text{ce}_m(q, z),$$

where

$$A_m = \frac{1}{\pi} \int_0^{2\pi} f(z) \text{ce}_m(q, z) dz.$$

Or

$$f(z) = \sum_{m=1}^{\infty} A_m \text{se}_m(q, z),$$

$$A_m = \frac{1}{\pi} \int_0^{2\pi} f(z) \text{se}_m(q, z) dz.$$

## 7.12 LEGENDRE FUNCTIONS

**The wave equation.** The Legendre wave equation {224} is

$$(z^2 - 1) \frac{d^2 Z}{dz^2} + 2z \frac{dZ}{dz} + \left[ \kappa^2 a^2 (z^2 - 1) - p(p+1) - \frac{q^2}{z^2 - 1} \right] Z = 0. \quad (7.80)$$

Expanding about the origin by the methods of § 7.03, we obtain

$$A_2 = A_4 = A_6 = \dots = -2,$$

$$B_2 = \kappa^2 a^2 + p(p+1) - q^2,$$

$$B_j = p(p+1) - jq^2/2$$

for  $j$  even and greater than 2. All other coefficients are zero. The indicial equation is

$$\beta(\beta - 1) = 0$$

with the roots  $b_1 = 1$ ,  $b_2 = 0$ . The determinants for these two cases are

$$\Delta'_m(1) = \begin{vmatrix} B_2 - 2 & 2 \cdot 3 & 0 & 0 & \dots & 0 \\ B_4 - 2 & B_2 - 6 & 4 \cdot 5 & 0 & \dots & 0 \\ B_6 - 2 & B_4 - 6 & B_2 - 10 & 6 \cdot 7 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ B_{2m-2} - 2 & B_{2m-4} - 6 & \dots & \dots & \dots & (2m-2)(2m-1) \\ B_{2m} - 2 & B_{2m-2} - 6 & \dots & \dots & \dots & B_2 - 4m+2 \end{vmatrix}, \quad (7.81)$$

$$\Delta'_m(0) = \begin{vmatrix} B_2 & 1 \cdot 2 & 0 & 0 & \dots & 0 \\ B_4 & B_2 - 4 & 3 \cdot 4 & 0 & \dots & 0 \\ B_6 & B_4 - 4 & B_2 - 8 & 5 \cdot 6 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ B_{2m-2} & B_{2m-4} - 4 & B_{2m-6} - 8 & \dots & \dots & (2m-3)(2m-2) \\ B_{2m} & B_{2m-2} - 4 & B_{2m-4} - 8 & \dots & \dots & B_2 - 4(m-1) \end{vmatrix}, \quad (7.82)$$

where

$$B_2 = \kappa^2 a^2 + p(p+1) - q^2,$$

$$B_{2l} = p(p+1) - lq^2, \quad l > 1.$$

The general solution of Eq. (7.80) is

$$Z = AZ_1 + BZ_2,$$

where

$$Z_1 = C_0 z \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(1) z^{2m}}{(2m+1)!}, \quad (7.83)$$

$$Z_2 = C_0 \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(0) z^{2m}}{(2m)!}. \quad (7.84)$$

This solution satisfies the differential equation for all values (integer or non-integer) of  $\kappa$ ,  $p$ , and  $q$ . It applies within a unit circle about the origin,  $|z| < 1$ . The same method may be employed to obtain a series solution for the region outside the unit circle.

The constant  $C_0$  is arbitrary. For simplicity, let  $C_0 = 1$  and define the *Legendre wave functions* by the above series:

$$\mathcal{P}_p^q(\kappa a, z) = z \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(1) z^{2m}}{(2m+1)!}, \quad (7.83 \text{ a})$$

$$\mathcal{Q}_p^q(\kappa a, z) = \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(0) z^{2m}}{(2m)!}. \quad (7.84 \text{ a})$$

The function  $\mathcal{P}_p^q$  is always an odd function, the function  $\mathcal{Q}_p^q$  is always even.

**Degenerate cases.** As indicated in § 7.01, the general Legendre wave functions of Eqs. (7.83 a) and (7.84 a) are solutions of {224}. These solutions may be written

$$\begin{aligned} \mathcal{P}_p^q(\kappa a, z) = z & \left\{ 1 - \frac{z^2}{3!} [\kappa^2 a^2 + p(p+1) - q^2 - 2] \right. \\ & + \frac{z^4}{5!} [\kappa^4 a^4 + 2\kappa^2 a^2 [p(p+1) - q^2 - 4] \\ & \left. + p(p+1) [p(p+1) - 2q^2 - 14] + q^4 + 20q^2 + 24] - \dots \right\}, \end{aligned}$$

$$\begin{aligned} \mathcal{Q}_p^q(\kappa a, z) = 1 & - \frac{z^2}{2!} [\kappa^2 a^2 + p(p+1) - q^2] \\ & + \frac{z^4}{4!} [\kappa^4 a^4 + 2\kappa^2 a^2 [p(p+1) - q^2 - 4] \\ & + p(p+1) [p(p+1) - 2q^2 - 6] + q^4 + 8q^2] - \dots. \end{aligned}$$

As degenerate cases, we have {222}, {220}, {114}, and {112}.

**I. Equation {222},  $\kappa = 0$ .** Then

$$\begin{aligned} \mathcal{P}_p^q(z) = z & \left\{ 1 - \frac{z^2}{3!} [p(p+1) - q^2 - 2] \right. \\ & + \frac{z^4}{5!} [p(p+1) [p(p+1) - 2q^2 - 14] + q^4 + 20q^2 + 24] - \dots \left. \right\}, \end{aligned}$$

$$\begin{aligned} \mathcal{Q}_p^q(z) = 1 & - \frac{z^2}{2!} [p(p+1) - q^2] \\ & + \frac{z^4}{4!} [p(p+1) [p(p+1) - 2q^2 - 6] + q^4 + 8q^2] - \dots \end{aligned}$$

II. Equation {220},  $p = 0$ .

$$\begin{aligned}\mathcal{P}_0^q(\kappa a, z) &= z \left\{ 1 - \frac{z^2}{3!} [\kappa^2 a^2 - q^2 - 2] \right. \\ &\quad \left. + \frac{z^4}{5!} [\kappa^4 a^4 - 2\kappa^2 a^2 (q^2 + 4) + q^4 + 20q^2 + 24] - \dots \right\}, \\ \mathcal{Q}_0^q(\kappa a, z) &= 1 - \frac{z^2}{2!} [\kappa^2 a^2 - q^2] \\ &\quad + \frac{z^4}{4!} [\kappa^4 a^4 - 2\kappa^2 a^2 [q^2 + 4] + q^4 + 8q^2] - \dots,\end{aligned}$$

III. Equation {114},  $q = 0$ .

$$\begin{aligned}\mathcal{P}_p(\kappa a, z) &= z \left\{ 1 - \frac{z^2}{3!} [\kappa^2 a^2 + p(p+1) - 2] + \frac{z^4}{5!} [\kappa^4 a^4 + 2\kappa^2 a^2 [p(p+1) - 4] \right. \\ &\quad \left. + p(p+1) [p(p+1) - 14] + 24] - \dots \right\}, \\ \mathcal{Q}_p(\kappa a, z) &= 1 - \frac{z^2}{2!} [\kappa^2 a^2 + p(p+1)] + \frac{z^4}{4!} [\kappa^4 a^4 + 2\kappa^2 a^2 [p(p+1) - 4] \\ &\quad + p(p+1) [p(p+1) - 6]] - \dots.\end{aligned}$$

IV. Equation {112},  $\kappa = 0$  and  $q = 0$ .

$$\begin{aligned}\mathcal{P}_p(z) &= z \left\{ 1 - \frac{z^2}{3!} [p(p+1) - 1 \cdot 2] + \frac{z^4}{5!} [p(p+1) - 1 \cdot 2] [p(p+1) - 3 \cdot 4] \right. \\ &\quad \left. - \frac{z^6}{7!} [p(p+1) - 1 \cdot 2] [p(p+1) - 3 \cdot 4] [p(p+1) - 5 \cdot 6] + \dots \right\}, \\ \mathcal{Q}_p(z) &= 1 - \frac{z^2}{2!} p(p+1) + \frac{z^4}{4!} p(p+1) [p(p+1) - 2 \cdot 3] \\ &\quad - \frac{z^6}{6!} p(p+1) [p(p+1) - 2 \cdot 3] [p(p+1) - 4 \cdot 5] + \dots. \\ \mathcal{P}_{p-\frac{1}{2}}(z) &= z \left\{ 1 - \frac{z^2}{3!} [(p^2 - \frac{1}{4}) - 1 \cdot 2] + \frac{z^4}{5!} [(p^2 - \frac{1}{4}) - 1 \cdot 2] [(p^2 - \frac{1}{4}) - 3 \cdot 4] \right. \\ &\quad \left. - \frac{z^6}{7!} [(p^2 - \frac{1}{4}) - 1 \cdot 2] [(p^2 - \frac{1}{4}) - 3 \cdot 4] [(p^2 - \frac{1}{4}) - 5 \cdot 6] + \dots \right\}, \\ \mathcal{Q}_{p-\frac{1}{2}}(z) &= 1 - \frac{z^2}{2!} (p^2 - \frac{1}{4}) + \frac{z^4}{4!} (p^2 - \frac{1}{4}) [(p^2 - \frac{1}{4}) - 2 \cdot 3] \\ &\quad - \frac{z^6}{6!} (p^2 - \frac{1}{4}) [(p^2 - \frac{1}{4}) - 2 \cdot 3] [(p^2 - \frac{1}{4}) - 4 \cdot 5] + \dots. \\ \mathcal{P}_{-\frac{1}{2}}(z) &= z \left\{ 1 + \frac{z^2}{3!} (1 \cdot 2 + \frac{1}{4}) + \frac{z^4}{5!} (1 \cdot 2 + \frac{1}{4}) (3 \cdot 4 + \frac{1}{4}) \right. \\ &\quad \left. + \frac{z^6}{7!} (1 \cdot 2 + \frac{1}{4}) (3 \cdot 4 + \frac{1}{4}) (5 \cdot 6 + \frac{1}{4}) + \dots \right\}, \\ \mathcal{Q}_{-\frac{1}{2}}(z) &= 1 + \frac{z^2}{2! 4} + \frac{z^4}{4! 4} [2 \cdot 3 + \frac{1}{4}] + \frac{z^6}{6! 4} [2 \cdot 3 + \frac{1}{4}] [4 \cdot 5 + \frac{1}{4}] + \dots.\end{aligned}$$

**Ordinary Legendre functions.** Note that the functions  $\mathcal{P}_p^q$  and  $\mathcal{Q}_p^q$  do not reduce directly to the familiar Legendre [45] functions  $P_n(z)$  and  $Q_n(z)$ . In obtaining the latter, one introduces other constants and makes other changes, as will be shown.

If  $\alpha = 0$ ,  $q = 0$ , and  $p = n$ , an integer, a simplification occurs because the infinite series become polynomials in certain cases. Evidently all  $B$ 's are equal,

$$B_{2l} = p(p+1),$$




Fig. 7.07. Ordinary Legendre polynomials  $P_n(\mu)$  for the range  $-1 \leq \mu \leq 1$ . The functions are alternately even and odd and all go through  $\pm 1$  at the ends of the interval




Fig. 7.08. Ordinary Legendre functions  $Q_n(\mu)$ . All functions become infinite at  $\mu = \pm 1$




Fig. 7.09. Ordinary Legendre functions  $P_n(x)$  over a wider range. Except for  $P_0(x)$ , all  $P_n(x) \rightarrow \infty$  as  $x \rightarrow \infty$

and the determinant  $A'_m(1)$  is zero if the last two rows are identical. Since the  $B$ 's are equal, the two rows are identical if

$$(2m-2)(2m-1) = p(p+1) - 4m + 2$$

or

$$p(p+1) = 2m(2m-1).$$

Therefore,  $p = 2m-1$ , an *odd integer*. Similarly,  $A'_m(0)$  becomes zero only if  $p = 2m-2$ , an *even integer*.

Thus, for  $\varkappa = 0$  and  $p = n$ , an integer, it is convenient to remove the polynomials from both  $Z_1$  and  $Z_2$  and to classify them as  $P_n$ . The remaining functions are called  $Q_n$ , and  $C_0$  is so chosen that

$$P_n(z) = \frac{(-1)^{n/2} n!}{2^n [(n/2)!]^2} \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(0) z^{2m}}{(2m)!}, \quad n \text{ even,} \quad (7.85)$$

$$P_n(z) = \frac{(-1)^{(n+1)/2} n! z}{2^{n-1} \left[ \left( \frac{n-1}{2} \right)! \right]^2} \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(1) z^{2m}}{(2m+1)!}, \quad n \text{ odd.} \quad (7.85)$$

$$Q_n(z) = \frac{(-1)^{n/2} 2^n [(n/2)!]^2 z}{n!} \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(1) z^{2m}}{(2m+1)!}, \quad n \text{ even,} \quad (7.86)$$

$$Q_n(z) = \frac{(-1)^{(n+1)/2} 2^{n-1} \left[ \left( \frac{n-1}{2} \right)! \right]^2}{n!} \sum_{m=0}^{\infty} \frac{(-1)^m A'_m(0) z^{2m}}{(2m)!}, \quad n \text{ odd.} \quad (7.86)$$




Fig. 7.10. Ordinary Legendre functions  $Q_n(x)$ . All are infinite at  $x = \pm 1$ ; all approach zero as  $x \rightarrow \infty$

$P_n(z)$  and  $Q_n(z)$  for the interval  $(-1, +1)$  are shown in Figs. 7.07 and 7.08. These are the *ordinary Legendre functions*, which will be employed only for  $\varkappa = 0$  and  $p = n$ , an integer. They must be clearly distinguished from the general functions  $\mathcal{P}_p^\varkappa(\varkappa a, z)$  and  $\mathcal{Q}_p^\varkappa(\varkappa a, z)$ , which do not reduce to Eqs. (7.85) and (7.86) but which are required except in the above special case.

The first six functions [46] obtained from Eq. (7.85) are

$$P_0(z) = 1,$$

$$P_1(z) = z,$$

$$P_2(z) = \frac{1}{2}(3z^2 - 1),$$

$$P_3(z) = \frac{1}{2}(5z^3 - 3z),$$

$$P_4(z) = \frac{1}{8}(35z^4 - 30z^2 + 3),$$

$$P_5(z) = \frac{1}{8}(63z^5 - 70z^3 + 15z).$$

Graphs are given in Figs. 7.07 and 7.09 for real values of  $z$ .

Legendre functions of the second kind may be obtained from Eq. (7.86):

$$Q_0(z) = \tanh^{-1} z = \frac{1}{2} \ln \left( \frac{1+z}{1-z} \right), \quad |z| < 1;$$

$$Q_0(z) = \coth^{-1} z = \frac{1}{2} \ln \left( \frac{z+1}{z-1} \right), \quad |z| > 1;$$




Fig. 7.11. Ordinary Legendre functions  $P_{1/2}(x)$ ,  $P_{3/2}(x)$ ,  $P_{5/2}(x)$ ,  $P_0(x)$ ,  $P_{1/2}(x)$




Fig. 7.12. Ordinary Legendre functions  $Q_{1/2}(x)$ ,  $Q_{3/2}(x)$ ,  $Q_{5/2}(x)$ ,  $Q_0(x)$ ,  $Q_{1/2}(x)$

$$Q_1(z) = z Q_0(z) - 1,$$

$$Q_2(z) = P_2(z) Q_0(z) - \frac{3}{2} z,$$

$$Q_3(z) = P_3(z) Q_0(z) - \frac{5}{2} z^2 + \frac{2}{3},$$

$$Q_4(z) = P_4(z) Q_0(z) - \frac{35}{8} z^3 + \frac{55}{24} z,$$

$$Q_5(z) = P_5(z) Q_0(z) - \frac{63}{8} z^4 + \frac{49}{8} z^2 - \frac{8}{15}, \dots$$

These functions are indicated in Figs. 7.08 and 7.10. Other graphs are given in Figs. 7.11 to 7.14. The general solution of Legendre's equation {112D}, with  $p = n$ , is

$$Z = A P_n(z) + B Q_n(z).$$




Fig. 7.13. Ordinary Legendre functions of the first kind for imaginary arguments




Fig. 7.14. Ordinary Legendre functions of the second kind for imaginary arguments

**Legendre associated functions [45].** The general form of the Legendre equation {222} is

$$(1 - z^2) \frac{d^2 Z}{dz^2} - 2z \frac{dZ}{dz} + \left[ p(p+1) - \frac{q^2}{1-z^2} \right] Z = 0. \quad (7.87)$$

If the constants are integers, the general solution is

$$Z = A P_n^m(z) + B Q_n^m(z),$$

where  $P_n^m$  and  $Q_n^m$  are the Legendre associated functions. For  $|z| < 1$ ,

$$\left. \begin{aligned} P_n^m(z) &= (1-z^2)^{m/2} \frac{d^m P_n(z)}{dz^m}, \\ Q_n^m(z) &= (1-z^2)^{m/2} \frac{d^m Q_n(z)}{dz^m}. \end{aligned} \right\} \quad (7.88)$$

For  $|z| > 1$ ,

$$\left. \begin{aligned} P_n^m(z) &= (z^2-1)^{m/2} \frac{d^m P_n(z)}{dz^m}, \\ Q_n^m(z) &= (z^2-1)^{m/2} \frac{d^m Q_n(z)}{dz^m}. \end{aligned} \right\} \quad (7.88a)$$

Definite integrals

$$P_n(z) = \frac{1}{\pi} \int_0^\pi [z \pm (z^2-1)^{\frac{1}{2}} \cos u]^n du.$$

$$P_n(z) = \frac{1}{\pi} \int_0^{\cos^{-1} z} \frac{du}{[z \mp (z^2-1)^{\frac{1}{2}} \cos u]^{n+1}}.$$

$$P_n(z) = \frac{2}{\pi} \int_0^{\cos^{-1} z} \frac{\cos(n+\frac{1}{2}) u du}{[2(\cos u - z)]^{\frac{1}{2}}}.$$

$$P_n(z) = \frac{2}{\pi} \int_{\cos^{-1} z}^{\infty} \frac{\sin(n+\frac{1}{2}) u du}{[2(z - \cos u)]^{\frac{1}{2}}}.$$

$$Q_n(z) = \frac{1}{2} \int_{-1}^1 \frac{P_n(u)}{z-u} du.$$

$$Q_n(z) = \int_0^\infty \frac{du}{[z + (z^2-1)^{\frac{1}{2}} \cosh u]^{n+1}}.$$

$$P_n^m(z) = \frac{2^m m! (n+m)!}{(2m)! (n-m)!} (z^2-1)^{m/2} \int_0^\pi \frac{\sin^{2m} u du}{[z + (z^2-1)^{\frac{1}{2}} \cos u]^{n+m+1}}.$$

$$Q_n^m(z) = (-1)^m \frac{2^m (n+m)! m!}{(n-m)! (2m)!} (z^2-1)^{m/2} \int_0^\infty \frac{\sinh^{2m} u du}{[z + (z^2-1)^{\frac{1}{2}} \cosh u]^{n+m+1}},$$

$(n-m) \geq 0$  and  $z$  not on real axis between  $\pm 1$ .

$$Q_n^m(z) = (-1)^m \frac{n!}{(n-m)!} \int_0^\infty \frac{\cosh mu du}{[z + (z^2-1)^{\frac{1}{2}} \cosh u]^{n+1}}$$

$n \geq m$  and  $z$  not on real axis between  $\pm 1$ .

Relations

$$nP_n - (2n-1)zP_{n-1} + (n-1)P_{n-2} = 0.$$

$$(z^2-1) \frac{dP_n}{dz} = -(n+1)(zP_n - P_{n+1}).$$

$$(n+1)(P_{n+1} - zP_n) - n(zP_n - P_{n-1}) = 0.$$

$$nP_n = z \frac{dP_n}{dz} - \frac{dP_{n-1}}{dz}.$$

$$(n+1)P_n = -z \frac{dP_n}{dz} + \frac{dP_{n+1}}{dz}.$$

$$\begin{aligned}
& (2n+1)(z^2-1) \frac{dP_n}{dz} = n(n+1)(P_{n+1} - P_{n-1}). \\
& nQ_n - (2n-1)zQ_{n-1} + (n-1)Q_{n-2} = 0. \\
& (2n+1)Q_n = \frac{dQ_{n+1}}{dz} - \frac{dQ_{n-1}}{dz}. \\
& P_n Q_{n-1} - P_{n-1} Q_n = 1/n. \\
& P_n Q_{n-2} - P_{n-2} Q_n = \frac{(2n-1)z}{(n-1)n}. \\
& P_n \frac{dQ_n}{dz} - Q_n \frac{dP_n}{dz} = (1-z^2)^{-1}. \\
& P_n^{m+2} + 2(m+1) \frac{z}{(z^2-1)^{\frac{1}{2}}} P_n^{m+1} - (n-m)(n+m+1) P_n^m = 0. \\
& (2n+1)zP_n^m - (n-m+1)P_{n+1}^m - (n+m)P_{n-1}^m = 0. \\
& (n-m+2)P_{n+2}^m - (2n+3)zP_{n+1}^m + (n+m+1)P_n^m = 0.
\end{aligned}$$

**Orthogonality.** Solutions of the Legendre wave equation {224} are

$$\mathcal{P}_p^q(\varkappa a, z) \quad \text{and} \quad \mathcal{Q}_p^q(\varkappa a, z).$$

Either of these functions, with proper boundary conditions, forms an orthogonal set on an arbitrary interval  $(a, b)$ . According to Table 7.01,

$$\int_a^b \mathcal{P}_{p_m}^q(\varkappa a, z) \mathcal{P}_{p_n}^q(\varkappa a, z) dz = 0,$$

and

$$\int_a^b \mathcal{Q}_{p_m}^q(\varkappa a, z) \mathcal{Q}_{p_n}^q(\varkappa a, z) dz = 0, \quad m \neq n.$$

Therefore, an arbitrary function  $f(z)$  may be expressed as a series of either  $\mathcal{P}$  or  $\mathcal{Q}$  functions,

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{P}_{p_m}^q(\varkappa a, z),$$

where

$$A_m = \frac{1}{N_m} \int_a^b f(z) \mathcal{P}_{p_m}^q(\varkappa a, z) dz,$$

$$N_m = \int_a^b [\mathcal{P}_{p_m}^q(\varkappa a, z)]^2 dz.$$

For the ordinary Legendre polynomials,

$$\int_{-1}^1 z^k P_n(z) dz = 0, \quad k = 0, 1, 2, \dots (n-1).$$

$$\int_0^1 z^k P_n(z) dz = \frac{k(k-1)(k-2)\dots(k-n+2)}{(k+n+1)(k+n-1)\dots(k-n+3)}.$$

$$\int_{-1}^1 P_m(z) P_n(z) dz = 0, \quad m \neq n.$$

$$\int_{-1}^1 [P_m(z)]^2 dz = \frac{2}{2m+1}.$$

An arbitrary function  $f(z)$  may be expressed as

$$f(z) = \sum_{m=0}^{\infty} A_m P_m(z),$$

where

$$A_m = \frac{2m+1}{2} \int_{-1}^1 f(z) P_m(z) dz.$$

### 7.13 LAMÉ FUNCTIONS

**The wave functions.** In the Lamé wave equation {1113},  $a_1$  may be set equal to zero without loss of generality. Then

$$\left. \begin{aligned} \frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{1}{z-b} + \frac{1}{z-c} \right] \frac{dZ}{dz} \\ + \frac{1}{4} \left[ \frac{(b^2+c^2)q - p(p+1)z + \kappa^2 z^2}{z(z-b)(z-c)} \right] Z = 0. \end{aligned} \right\} \quad (7.89)$$

To obtain a series solution about the origin, let

$$Z = z^\beta \sum_{j=0}^{\infty} C_j z^j.$$

Then, according to § 7.03,

$$A_0 = \frac{1}{2},$$

$$A_1 = -\frac{b+c}{2bc},$$

$$A_2 = -\frac{1}{2(bc)^2} [b^2 + c^2],$$

$$A_j = \frac{1}{bc} [A_{j-1}(b+c) - A_{j-2}] \quad \text{for } j > 2.$$

Also,

$$B_0 = 0,$$

$$B_1 = \frac{q(b^2+c^2)}{4bc},$$

$$B_2 = \frac{1}{4(bc)^2} [q(b+c)(b^2+c^2) - p(p+1)bc],$$

$$B_3 = \frac{1}{4(bc)^3} [q(b^2+c^2)(b^2+bc+c^2) - p(p+1)bc(b+c) + \kappa^2(bc)^2],$$

$$B_j = \frac{1}{bc} [B_{j-1}(b+c) - B_{j-2}] \quad \text{for } j > 3.$$

The indicial equation is

$$\beta(\beta - \frac{1}{2}) = 0,$$

$b_1 = \frac{1}{2}$ ,  $b_2 = 0$ . Therefore

$$A_j(\frac{1}{2}) = \left| \begin{array}{cccccc} \frac{1}{2}A_1 + B_1 & \frac{3}{2} & 0 & 0 & \dots & 0 \\ \frac{1}{2}A_2 + B_2 & \frac{3}{2}A_1 + B_1 & 5 & 0 & \dots & 0 \\ \frac{1}{2}A_3 + B_3 & \frac{3}{2}A_2 + B_2 & \frac{5}{2}A_1 + B_1 & \frac{21}{2} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & (j-\frac{1}{2})(j-1) & \\ \frac{1}{2}A_j + B_j & \dots & \dots & \dots & (j-\frac{1}{2})A_1 + B_1 & \end{array} \right|, \quad (7.90)$$

and

$$\mathcal{A}_j(0) = \begin{vmatrix} B_1 & \frac{1}{2} & 0 & 0 & \dots & 0 \\ B_2 & A_1 + B_1 & 3 & 0 & \dots & 0 \\ B_3 & A_2 + B_2 & 2A_1 + B_1 & \frac{15}{2} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & (j-1)(j-\frac{3}{2}) & \dots \\ B_j & A_{j-1} + B_{j-1} & \dots & \dots & (j-1)A_1 + B_1 & \dots \end{vmatrix}. \quad (7.91)$$

The solutions of the Lamé wave equation are

$$\left. \begin{aligned} Z_1 &= C_0 z^{\frac{1}{2}} \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} \mathcal{A}_j(\frac{1}{2}) z^j}{(2j+1)!}, \\ Z_2 &= C_0 \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} \mathcal{A}_j(0) z^j}{(2j)!}. \end{aligned} \right\} \quad (7.92)$$

It is customary to express Lamé functions with a different variable. If  $z = \xi^2$  and  $C_0 = 1$ , we have the *Lamé wave functions*

$$\left. \begin{aligned} \mathcal{E}_p^q(\varkappa, \xi) &= \xi \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} \mathcal{A}_j(\frac{1}{2}) \xi^{2j}}{(2j+1)!}, \\ \mathcal{F}_p^q(\varkappa, \xi) &= \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} \mathcal{A}_j(0) \xi^{2j}}{(2j)!}. \end{aligned} \right\} \quad (7.92a)$$

The general solution of Eq. (7.89) is then

$$Z = A \mathcal{E}_p^q(\varkappa, \xi) + B \mathcal{F}_p^q(\varkappa, \xi). \quad (7.93)$$

The series are convergent within a circle about the origin and extending out to the nearest singularity. The same procedure, however, may be employed in expanding about any other point.

**Polynomial solutions.** The solutions (7.92a) and (7.93) apply also to the ordinary Lamé equation {1112} and to the degenerate case {1111} where  $p = 0$ . There are a few very special cases in which the infinite series reduce to polynomials [47], much as with Legendre polynomials. These are the ordinary Lamé functions  $E_p^q(z)$  treated in books on the subject. Not only does  $\varkappa$  have to be zero and  $p$  an integer, but  $q$  must be related to  $p$  in a definite way to allow a polynomial.

**Case I.**  $C_0 = 1, \beta = 0, \varkappa = p = q = 0, \mathcal{F}_0^0(0, z) = 1$ .

**Case II.**  $C_0 = 1, \beta = 1, \varkappa = 0, p = 2, q = 1, \mathcal{E}_1^2(0, z) = z$ .

**Case III.**  $C_0 = 1, \beta = 0, \varkappa = 0, p = 2,$

$$q_1 = 2b^2c^2 \left[ 1 + \left( 1 - \frac{3b^2c^2}{(b^2+c^2)^2} \right)^{\frac{1}{2}} \right],$$

$$q_2 = 2b^2c^2 \left[ 1 - \left( 1 - \frac{3b^2c^2}{(b^2+c^2)^2} \right)^{\frac{1}{2}} \right],$$

$$\mathcal{F}_2^{q_1}(0, z) = 1 - \frac{3z^2}{(b^2+c^2) + [(b^2+c^2)^2 - 3b^2c^2]^{\frac{1}{2}}},$$

$$\mathcal{F}_2^{q_1}(0, z) = 1 - \frac{3z^2}{(b^2+c^2) - [(b^2+c^2)^2 - 3b^2c^2]^{\frac{1}{2}}}.$$

**Case IV.**  $C_0 = 1$ ,  $\beta = 1$ ,  $\varkappa = 0$ ,  $\phi = 3$ ,

$$q_1 = 1 - \frac{6b^2c^2}{5} \left[ 2 + \left( 4 - \frac{15b^2c^2}{(b^2+c^2)^2} \right)^{\frac{1}{2}} \right],$$

$$q_2 = 1 - \frac{6b^2c^2}{5} \left[ 2 - \left( 4 - \frac{15b^2c^2}{(b^2+c^2)^2} \right)^{\frac{1}{2}} \right],$$

$$\mathcal{E}_3^{q_1}(0, z) = 1 - \frac{5z^3}{2(b^2+c^2) + [4(b^2+c^2)^2 - 15b^2c^2]^{\frac{1}{2}}},$$

$$\mathcal{E}_3^{q_2}(0, z) = 1 - \frac{5z^3}{2(b^2+c^2) - [4(b^2+c^2)^2 - 15b^2c^2]^{\frac{1}{2}}}.$$

Polynomial solutions of higher degree also occur for discrete values of  $\phi$  and  $q$ .

Other interesting special cases can be expressed in closed form:

$$\mathcal{F}_1^q(0, z) = \frac{1}{ib} (z^2 - b^2)^{\frac{1}{2}} \quad \text{if } q = \frac{c^2}{b^2 + c^2},$$

$$\mathcal{F}_1^q(0, z) = \frac{1}{ic} (z^2 - c^2)^{\frac{1}{2}} \quad \text{if } q = \frac{b^2}{b^2 + c^2},$$

$$\mathcal{F}_2^1(0, z) = \frac{1}{bc} [(z^2 - b^2)(z^2 - c^2)]^{\frac{1}{2}}.$$

Related to these solutions are further solutions which reduce to polynomials times square roots, for suitable values of  $\phi$  and  $q$ .

**Orthogonality.** Solutions of the Lamé wave equation  $\{1113\}$  are  $\mathcal{E}_p^q(\varkappa, \zeta)$  and  $\mathcal{F}_p^q(\varkappa, \zeta)$ . These functions may be employed in expressing an arbitrary function  $f(\zeta)$  over an arbitrary interval  $(a, b)$ . According to Table 7.01, orthogonality is obtained with a weighting function, assuming Sturm-Liouville boundary conditions. Thus, for example,

$$\int_a^b \frac{z}{u} \mathcal{E}_{p_m}^q(\varkappa, \zeta) \mathcal{E}_{p_n}^q(\varkappa, \zeta) d\zeta = 0, \quad m \neq n.$$

Thus,

$$f(\zeta) = \sum_{m=0}^{\infty} A_m \mathcal{E}_{p_m}^q(\varkappa, \zeta),$$

where

$$A_m = \frac{1}{N_m} \int_a^b \frac{\zeta^2}{u} f(\zeta) \mathcal{E}_{p_m}^q(\varkappa, \zeta) d\zeta,$$

$$N_m = \int_a^b \frac{\zeta^2}{u} [\mathcal{E}_{p_m}^q(\varkappa, \zeta)]^2 d\zeta.$$

A similar expansion can be made on  $q_m$ , or the  $\mathcal{F}$ -functions can be employed instead of the  $\mathcal{E}$ -functions.

### 7.14 WANGERIN FUNCTIONS

Equation  $\{1122\}$  is written [48]

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z-b} + \frac{1}{z-c} + \frac{2}{z} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2}{z^2(z-b)(z-c)} \right] Z = 0, \quad (7.94)$$

if  $a_3 = 0$ . For a solution about the origin,

$$Z = z^\beta \sum_{j=0}^{\infty} C_j z^j.$$

The method of § 7.02 then gives

$$\begin{aligned} A_0 &= 1, \\ A_1 &= -\frac{b+c}{2bc}, \\ A_2 &= -\frac{1}{2(bc)^2} [b^2 + c^2], \\ A_j &= \frac{1}{bc} [A_{j-1}(b+c) - A_{j-2}], \quad j > 2. \end{aligned}$$

Also,

$$\begin{aligned} B_0 &= \frac{\bar{A}_0}{4bc}, \\ B_1 &= \frac{1}{4(bc)^2} [\bar{A}_0(b+c) + \bar{A}_1 bc], \\ B_2 &= \frac{1}{4(bc)^3} [\bar{A}_0(b^2 + bc + c^2) + \bar{A}_1 bc(b+c) + \bar{A}_2(bc)^2], \\ B_j &= \frac{1}{bc} [B_{j-1}(b+c) - B_{j-2}], \quad j > 2. \end{aligned}$$

Therefore, the determinant is

$$\Delta_j(\beta) = \left| \begin{array}{cccccc} \beta A_1 + B_1 & (\beta+1)^2 + B_0 & 0 & 0 & \dots & 0 \\ \beta A_2 + B_2 & (\beta+1) A_1 + B_1 & (\beta+2)^2 + B_0 & 0 & \dots & 0 \\ \beta A_3 + B_3 & (\beta+1) A_2 + B_2 & (\beta+2) A_1 + B_1 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & (\beta+j-1)^2 + B_0 \\ \beta A_j + B_j & (\beta+1) A_{j-1} + B_{j-1} & \dots & \dots & \dots & (\beta+j-1) A_1 + B_1 \end{array} \right|. \quad (7.95)$$

The indicial equation gives

$$b_1 = +\left(-\frac{\bar{A}_0}{4bc}\right)^{\frac{1}{2}}, \quad b_2 = -\left(-\frac{\bar{A}_0}{4bc}\right)^{\frac{1}{2}},$$

and the solutions are

$$\left. \begin{aligned} Z_1 &= C_0 z^{b_1} \sum_{j=0}^{\infty} \frac{(-1)^j \Delta_j(b_1) z^j}{\prod_{l=1}^j [(b_1+l)^2 + B_0]}, \\ Z_2 &= C_0 \sum_{j=0}^{\infty} \frac{(-1)^j \Delta_j(b_2) z^j}{\prod_{l=1}^j [(b_2+l)^2 + B_0]}, \end{aligned} \right\} \quad (7.96)$$

if  $b_1$  and  $b_2$  are distinct and do not differ by an integer.

The *Wangerin functions* may be defined for  $C_0 = 1$ ,  $b = 1$ ,  $c = 1/k^2$ ,  $\bar{A}_0 = -q^2 c$ ,  $\bar{A}_1 = -p^2 c$ ,  $\bar{A}_2 = 1 - q^2$ . Then Eq. (7.96) becomes

$$\left. \begin{aligned} \mathcal{S}_p^q(k, z) &= z^{q/2} \sum_{m=0}^{\infty} \frac{(-1)^m \Delta_m(q/2) z^m}{\prod_{l=1}^m [(l+q/2)^2 - q^2/4]}, \\ \mathcal{T}_p^q(k, z) &= \sum_{m=0}^{\infty} \frac{(-1)^m \Delta_m(-q/2) z^m}{\prod_{l=1}^m [(l-q/2)^2 - q^2/4]} \end{aligned} \right\} \quad (7.96a)$$

where

$$\begin{aligned} A_0 &= 1, \\ A_1 &= -\left(\frac{1+c}{2c}\right), \\ A_2 &= -\left(\frac{(1+c^2)}{2c^2}\right), \\ A_j &= \frac{1}{c} [A_{j-1}(1+c) - A_{j-2}] \quad \text{for } j > 2: \\ B_0 &= -q^2/4, \\ B_1 &= -\frac{1}{4c} [q^2(1+c) + p^2], \\ B_2 &= -\frac{1}{4c^2} [q^2(1+c)^2 + p^2(1+c) - c], \\ B_j &= \frac{1}{c} [B_{j-1}(1+c) - B_{j-2}]. \end{aligned}$$

Equation (7.96a) converges for a circular region with center at the origin and extending out to the next singularity. The same procedure allows the  $\mathcal{S}$  and  $\mathcal{T}$  functions to be expanded about any other point.

**Orthogonality.** The Wangerin functions  $\mathcal{S}_p^q(k, z)$  and  $\mathcal{T}_p^q(k, z)$  form orthogonal sets, with weighting function, on an arbitrary interval  $(a, b)$ . With Sturm-Liouville boundary conditions and with weighting function  $w = z/u$ ,

$$\int_a^b \frac{z}{u} \mathcal{S}_{p_m}^q(k, z) \mathcal{S}_{p_n}^q(k, z) dz = 0, \quad m \neq n.$$

An arbitrary function  $f(z)$  may be expressed therefore as

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{S}_{p_m}^q(k, z) dz,$$

where

$$\begin{aligned} A_m &= \frac{1}{N_m} \int_a^b \frac{z}{u} f(z) \mathcal{S}_{p_m}^q(k, z) dz, \\ N_m &= \int_a^b \frac{z}{u} [\mathcal{S}_{p_m}^q(k, z)]^2 dz. \end{aligned}$$

This result applies equally well to the  $\mathcal{T}$ -function.

### 7.15 HEINE FUNCTIONS

The Heine equation {1222} is

$$\frac{d^2Z}{dz^2} + \frac{1}{2} \left[ \frac{1}{z} + \frac{2}{z-b} + \frac{2}{z-c} \right] \frac{dZ}{dz} + \frac{1}{4} \left[ \frac{\bar{A}_0 + \bar{A}_1 z + \bar{A}_2 z^2 + \bar{A}_3 z^3}{z(z-b)^2(z-c)^2} \right] Z = 0, \quad (7.97)$$

if  $a_1 = 0$ . By the method of § 7.03, expanding about  $z = 0$ ,

$$\begin{aligned} A_0 &= \frac{1}{2}, \\ A_1 &= -\frac{(b+c)}{bc}, \\ A_2 &= -\frac{(b^2+c^2)}{(bc)^2}, \\ A_j &= \frac{1}{bc} [A_{j-1}(b+c) - A_{j-2}] \quad \text{for } j > 2. \end{aligned}$$

Also,

$$B_0 = 0,$$

$$B_1 = \frac{\bar{A}_0}{4(bc)^2},$$

$$B_2 = \frac{1}{4(bc)^3} [2(b+c)\bar{A}_0 + bc\bar{A}_1],$$

$$B_3 = \frac{1}{4(bc)^4} [(3b^2 + 4bc + 3c^2)\bar{A}_0 + 2bc(b+c)\bar{A}_1 + (bc)^2\bar{A}_2],$$

$$B_4 = \frac{1}{4(bc)^5} [2(b+c)(2b^2 + bc + 2c^2)\bar{A}_0 + bc(3b^2 + bc + 3c^2)\bar{A}_1 + 2(bc)^2(b+c)\bar{A}_2 + (bc)^3\bar{A}_3],$$

$$B_j = \frac{1}{(bc)^j} [2B_{j-1}bc(b+c) - B_{j-2}(b^2 + 4bc + c^2) + 2B_{j-3}(b+c) - B_{j-4}] \quad \text{for } j > 4.$$

The indicial equation is

$$\beta(\beta - \frac{1}{2}) = 0,$$

so  $b_1 = \frac{1}{2}$ ,  $b_2 = 0$ . For  $\beta = b_1 = \frac{1}{2}$ , the determinant is

$$A_j(\frac{1}{2}) = \begin{vmatrix} \frac{1}{2}\bar{A}_1 + B_1 & \frac{3}{2} & 0 & 0 & \dots & 0 \\ \frac{1}{2}\bar{A}_2 + B_2 & \frac{3}{2}\bar{A}_1 + B_1 & \frac{1}{2}0 & 0 & \dots & 0 \\ \frac{1}{2}\bar{A}_3 + B_3 & \frac{3}{2}\bar{A}_2 + B_2 & \frac{5}{2}\bar{A}_1 + B_1 & \frac{2}{2}1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & (j-\frac{1}{2})(j-1) & \dots \\ \frac{1}{2}\bar{A}_j + B_j & \dots & \dots & \dots & (j-\frac{1}{2})A_1 + B_1 & \dots \end{vmatrix}.$$

For  $\beta = b_2 = 0$ ,

$$A_j(0) = \begin{vmatrix} B_1 & \frac{1}{2} & 0 & 0 & \dots & 0 \\ B_2 & A_1 + B_1 & 3 & 0 & \dots & 0 \\ B_3 & A_2 + B_2 & 2A_1 + B_1 & \frac{1}{2}5 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & (j-1)(j-1-\frac{1}{2}) & \dots \\ B_j & A_{j-1} + B_{j-1} & \dots & \dots & (j-1)A_1 + B_1 & \dots \end{vmatrix}.$$

Thus the solutions of Eq. (7.97) are

$$\left. \begin{aligned} Z_1 &= C_0 z^{\frac{1}{2}} \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} A_j(\frac{1}{2}) z^j}{(2j+1)!}, \\ Z_2 &= C_0 \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} A_j(0) z^j}{(2j)!}. \end{aligned} \right\} \quad (7.98)$$

These series are convergent within a circular region whose center is at the origin and whose perimeter passes through the nearest singularity ( $z = b$ ).

The form of Eq. (7.97) needed in field theory has  $b = 1$ ,  $\bar{A}_0 = -\dot{p}^2 c^2$ ,  $\bar{A}_1 = \dot{p}^2 c(1+c) - q^2(1-c)^2 + 2c$ ,  $\bar{A}_2 = -[\dot{p}^2 c + 2(1+c)]$ ,  $\bar{A}_3 = 2$ . Setting  $C_0 = 1$  and  $c = 1/k^2$ , we obtain the *Heine functions*:

$$\left. \begin{aligned} \mathcal{U}_p^q(k, z) &= z^{\frac{1}{2}} \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} A_j(\frac{1}{2}) z^j}{(2j+1)!}, \\ \mathcal{V}_p^q(k, z) &= \sum_{j=0}^{\infty} \frac{(-1)^j 2^{2j} A_j(0) z^j}{(2j)!}, \end{aligned} \right\} \quad (7.99)$$

where

$$A_0 = \frac{1}{2},$$

$$A_1 = -(1+c)/c,$$

$$A_2 = -(1+c^2)/c^2,$$

$$A_j = \frac{1}{c} [A_{j-1}(1+c) - A_{j-2}] \quad \text{for } j > 2;$$

$$B_0 = 0,$$

$$B_1 = -\frac{\dot{p}^2}{4c},$$

$$B_2 = -\frac{1}{4c^2} [\dot{p}^2(1+c) + q^2 - 2c],$$

$$B_3 = -\frac{1}{4c^3} [\dot{p}^2(1+c+c^2) + 2q^2(1+c) - 2c(1+c)], \dots$$

**Orthogonality.** Like the other functions of this book, the Heine functions form orthogonal sets on the arbitrary interval  $(a, b)$  with respect to a weighting function  $w(z)$ . With Sturm-Liouville boundary conditions,

$$\int_a^b w(z) \mathcal{U}_{p_m}^q(k, z) \mathcal{U}_{p_n}^q(k, z) dz = 0, \quad m \neq n.$$

Therefore, an arbitrary function of  $z$  may be expressed as

$$f(z) = \sum_{m=0}^{\infty} A_m \mathcal{U}_{p_m}^q(k, z),$$

where

$$A_m = \frac{1}{N_m} \int_a^b w(z) f(z) \mathcal{U}_{p_m}^q(k, z) dz,$$

$$N_m = \int_a^b w(z) [\mathcal{U}_{p_m}^q(k, z)]^2 dz.$$

These relations hold also for the functions  $\mathcal{V}_p^q(k, z)$ , the weighting functions being taken from Table 7.01.

## BIBLIOGRAPHY

*For Section I*

- [1] LAMÉ, G.: Leçons sur les coordonnées curvilignes et leurs diverses applications. Paris: Mallet-Bachelier 1859.
- [2] DARBOUX, G.: Sur une classe remarquable de courbes et de surfaces algébriques. Paris 1896. — Leçons sur les systèmes orthogonaux et les coordonnées curvilignes. Paris: Gauthier-Villars 1910.
- MÜLLER, E.: Die verschiedenen Koordinatensysteme. Encyk. Math. Wissen., Bd. III.1.1., S. 596. Leipzig: B. G. Teubner 1907—1910.
- [3] EISENHART, L. P.: Separable systems of STÄCKEL. Ann. Math. **35**, 284 (1934). — Stäckel systems in conformal euclidean space. Ann. Math. **36**, 57 (1935). ROBERTSON, H. P.: Bemerkung über separierbare Systeme in der Wellenmechanik. Math. Ann. **98**, 749 (1927).
- [4] BÖCHER, M.: Über die Reihenentwickelungen der Potentialtheorie. Leipzig: B. G. Teubner 1894.
- [5] MOON, P., and D. E. SPENCER: Foundations of electrodynamics. Princeton, N. J.: D. Van Nostrand Co. 1960.
- [6] MOON, P., and D. E. SPENCER: The meaning of the vector Laplacian. J. Franklin Inst. **256**, 551 (1953).
- [7] MOON, P., and D. E. SPENCER: Field theory for engineers, Chap. 11. Princeton, N. J.: D. Van Nostrand Co. 1961.
- [8] STÄCKEL, P.: Über die Integration der Hamilton-Jacobischen Differentialgleichung mittels Separation der Variablen. Habil.-Schr. Halle 1891. — Sur une classe de problèmes de dynamique. C.R., Acad. Sci., Paris **116**, 485 (1893). — Über die Integration der Hamiltonschen Differentialgleichung mittels Separation der Variablen. Math. Ann. **49**, 145 (1897).
- [9] MOON, P., and D. E. SPENCER: Separability conditions for the Laplace and Helmholtz equations. J. Franklin Inst. **253**, 585 (1952). — Theorems on separability in Riemannian  $n$ -space. Amer. Math. Soc. Proc. **3**, 635 (1952). — Recent investigations of the separation of LAPLACE's equation. Amer. Math. Soc. Proc. **4**, 302 (1953). — Separability in a class of coordinate systems. J. Franklin Inst. **254**, 227 (1952).
- [10] BLASCHKE, W.: Eine Verallgemeinerung der Theorie der konfokalen  $E_2$ . Math. Z. **27**, 653 (1928). LEVI-CIVITA, T.: Sulla integrazione della equazione di Hamilton-Jacobi per separazione di variabili. Math. Ann. **59**, 383 (1904). TURRIERE, E.: Démonstration du théorème de STÄCKEL par l'élimination du temps entre les équations de LAGRANGE. L'enseignement Math. **22**, 337 (1923). FOGELSANG, W.: Eine Verallgemeinerung der konfokalen Flächen zweiter Ordnung. Math. Z. **35**, 25 (1932).
- [11] *Textbooks dealing with separation of variables*
- BATEMAN, H.: Partial differential equations of mathematical physics. Cambridge: Cambridge Univ. Press 1932; New York: Dover Publications 1944. BYERLY, W. E.: Fourier series and spherical, cylindrical, and ellipsoidal harmonics. Boston: Ginn & Co. 1893; New York: Dover Publications 1959.

- CHURCHILL, R. V.: Fourier series and boundary value problems. New York: McGraw-Hill Book Co. 1941.
- COURANT, R., u. D. HILBERT: Methoden der mathematischen Physik. Berlin: Springer 1931.
- FRANK, P., u. R. v. MISES: Die Differential- und Integralgleichungen der Mechanik und Physik, 8th edit. of RIEMANN-WEBER. Braunschweig: Vieweg & Sohn 1930.
- HILDEBRAND, F. B.: Advanced calculus for engineers. Englewood Cliffs, N. J.: Prentice-Hall 1949.
- JEFFREYS, H., and B. S. JEFFREYS: Methods of mathematical physics. Cambridge: Cambridge Univ. Press 1950.
- KELLOGG, O. D.: Foundations of potential theory. Berlin: Springer 1929; New York: Dover Publications 1953.
- LENSE, J.: Reihenentwicklungen in der mathematischen Physik. Berlin: Walter de Gruyter 1933.
- MAXWELL, J. C.: Electricity and magnetism. London: Oxford Univ. Press 1904; first edit. 1873.
- MOON, P., and D. E. SPENCER: Field theory for engineers. Princeton, N. J.: D. Van Nostrand Co. 1961.
- MORSE, P. M., and H. FESHBACH: Methods of theoretical physics. New York: McGraw-Hill Book Co. 1953.
- MURNAGHAN, F. D.: Introduction to applied mathematics. New York: John Wiley & Sons 1948.
- SMYTHE, W. R.: Static and dynamic electricity. New York: McGraw-Hill Book Co. 1950.
- SOMMERFELD, A.: Partial differential equations in physics. New York: Academic Press 1949.
- STRATTON, J. A.: Electromagnetic theory. New York: McGraw-Hill Book Co. 1941.
- WEBER, E.: Electromagnetic fields. New York: John Wiley & Sons 1950.
- WEBSTER, A. G.: Partial differential equations of mathematical physics. Leipzig: B. G. Teubner 1933.

*For Sections II, III, and IV*

- [12] KLEIN, F.: Vorlesungen über lineare Differentialgleichungen der zweiten Ordnung. Bearbeitet von E. RITTER. Göttingen 1894.
- [13] *Cyclides*
- CASEY, M.: On cyclides and spherico-quartics. Phil. Trans. Roy. Soc. Lond. **161**, 585 (1871).
- DARBOUX, G.: Remarques sur la théorie des surfaces orthogonales. C. R. Acad. Sci., Paris **59**, 240 (1864). — Sur l'application des méthodes de la physique mathématique à l'étude des corps terminés par des cyclides. C. R. Acad. Sci., Paris **83**, 1037, 1099 (1876).
- MAXWELL, J. C.: On the cyclide. Quart. J. Math. 9, 111 (1868).
- [14] *Inversion*
- COOLIDGE, J. L.: A treatise on the circle and the sphere. London: Oxford Univ. Press 1916.
- DARBOUX, G.: Leçons sur les systèmes orthogonaux et les coordonnées curvilignes, p. 277. Paris: Gauthier-Villars 1910.
- JEANS, J. H.: Electricity and magnetism. Cambridge: Cambridge Univ. Press 1925.
- MAXWELL, J. C.: Electricity and magnetism. London: Oxford Univ. Press 1904.
- SCHMIDT, H.: Die Inversion und ihre Anwendung. München: R. Oldenbourg 1950.
- THOMSON, W.: Extrait d'un lettre à M. LIOUVILLE. J. Math. pures appl. **10**, 364 (1845). (The first use of inversion in electrostatics.) — Extraits de deux lettres adressées à M. LIOUVILLE. J. Math. pures appl. **12**, 256 (1847).

- LIOUVILLE, J.: Note au sujet de l'article précédent. *J. Math. pures appl.* **12**, 265 (1847).
- THOMPSON, W., and P. G. TAIT: Treatise on natural philosophy, Part II, p. 62. Cambridge: Cambridge Univ. Press 1890.
- WANGERIN, A.: Theorie des Potentials und der Kugelfunktionen, Bd. II, S. 147. Berlin: Walter de Gruyter 1921.
- WEBER, E.: Electromagnetic fields, p. 244. New York: John Wiley & Sons 1950.
- WRINCH, D. M.: Inverted prolate spheroids. *Phil. Mag.* **14**, 1061 (1932).
- [15] MOON, P., and D. E. SPENCER: Cylindrical and rotational coordinate systems. *J. Franklin Inst.* **252**, 327 (1951). — Some coordinate systems associated with elliptic functions. *J. Franklin Inst.* **255**, 531 (1953).
- [16] *Theory of a complex variable*
- CHURCHILL, R. V.: Complex variables and applications. New York: McGraw-Hill Book Co. 1960.
- KNOPP, K.: Theory of functions. New York: Dover Publications 1945.
- NEHARI, Z.: Conformal mapping. New York: McGraw-Hill Book Co. 1952.
- OSGOOD, W. F.: Lehrbuch der Funktionentheorie. Leipzig: B. G. Teubner 1928.
- ROTHE, R., F. OLENDORFF and K. POHLHAUSEN: Theory of functions. Cambridge, Mass.: Technology Press 1933.
- TITCHMARSH, E. C.: Theory of functions. London: Oxford Univ. Press 1932.
- TOWNSEND, E. J.: Functions of a complex variable. New York: Henry Holt & Co. 1915.
- WALKER, M.: Conjugate functions for engineers. London: Oxford Univ. Press 1933.
- [17] *Complex transformations*
- KOBER, H.: Dictionary of conformal representations. New York: Dover Publications 1952.
- LOVE, A. E. H.: Some electrostatic distributions in two dimensions. *Proc. London Math. Soc.* **22**, 337 (1924).
- MICHELL, J. H.: A map of the complex  $Z$ -function: a condenser problem. *Messenger of Math.* **23**, 72 (1894).
- GREENHILL, G.: Theory of a stream line past a plane barrier. Advisory Comm. for Aeronautics, Report No. 19, London 1910.
- [18] *Applications of complex transformations*
- ANDRONESCU, P.: Das parallel- und meridiane Ebene Feld nebst Beispielen. *Arch. Elektrotechn.* **14**, 379 (1924).
- BAIRSTOW, L.: Resistance of a cylinder moving in a viscous fluid. *Phil. Trans. Roy. Soc. Lond.* **223**, 383 (1923).
- GLAUERT, H.: Elements of aerofoil and airscrew theory. Cambridge: Cambridge Univ. Press 1926.
- GRÖSSER, W.: Einige elektrostatische Probleme der Hochspannungstechnik. *Arch. Elektrotechn.* **25**, 193 (1931).
- HOLTZMÜLLER, O.: Über die logarithmische Abbildung und die aus ihr entstehenden orthogonalen Kurvensysteme. *Z. Math. Phys.* **16**, 269 (1871). Beiträge zur Theorie der isogonalen Verwandtschaften. *Z. Math. Phys.* **18**, 227 (1873). — Weitere Beiträge zur Theorie der isogonalen Verwandtschaften. *Z. Math. Phys.* **20**, 1 (1875). — Einführung in die Theorie der isogonalen Verwandtschaften. Leipzig: B. G. Teubner 1882.
- KEHREN, E.: Anwendung der konformen Abbildung in der Elektrostatik. *Ann. d. Phys.* **14**, 367 (1932).
- KNIGHT, R. C.: The potential of a circular cylinder between two infinite planes. *Proc. London Math. Soc.* **39**, 272 (1933).
- LABUS, J.: Berechnung des elektrischen Feldes von Hochspannungstransformatoren mit Hilfe der konformen Abbildung. *Arch. Elektrotechn.* **19**, 82 (1927). — Der Potential- und Feldverlauf längs einer Transformatorkwicklung. *Arch. Elektrotechn.* **21**, 250 (1928).

- LEVI-CIVITA, T.: Sopra un problema di elettrostatica che si è presentato nella costruzione dei cavi. *Rend. Circ. Math. Palermo* **20** (1905).
- LEVY, H.: Discontinuous fluid motion past a curved boundary. *Proc. Roy. Soc. Lond.* **92**, 285 (1915).
- MCLACHLAN, N. W.: Heat conduction in elliptical cylinder and an analogous electromagnetic problem. *Phil. Mag.* **36**, 600 (1945).
- MEYER, E.: Zwei Beispiele zweidimensionaler elektrostatischer Kraftlinienbilder. *Math. Ann.* **93**, 157 (1925).
- MORTON, W. B.: The electrification of two intersecting planes. *Phil. Mag.* **1**, 337 (1926). — On the parallel-plate condenser and other two-dimensional fields specified by elliptic functions. *Phil. Mag.* **2**, 827 (1926).
- NICHOLSON, J. W.: The electrification of two parallel circular disks. *Phil. Trans. Roy. Soc. Lond.* **224**, 303 (1923/24).
- PAGE, W. M.: Some two-dimensional problems in electrostatics and hydrodynamics. *Proc. London Math. Soc.* **11**, 313 (1913).
- PETERSOHN, H.: Zweidimensionale elektrostatische Probleme. *Z. Physik* **38**, 727 (1926).
- POOLE, E. G. C.: On the discontinuous motion produced in an infinite stream by two plane obstacles. *Proc. London Math. Soc.* **22**, 425 (1924).
- PORITSKY, H.: Field due to two equally charged parallel conducting cylinders. *J. Math. Phys.* **11**, 213 (1932).
- RICHMOND, H. W.: On the electrostatic field of a plane or circular grating formed of thick rounded bars. *Proc. London Math. Soc.* **22**, 389 (1924). — Notes on the use of the Schwartz-Christoffel transformation in electrostatics (and hydrodynamics). *Proc. London Math. Soc.* **22**, 483 (1924).
- ROGOWSKI, W.: Die elektrische Festigkeit am Rande des Plattenkondensators. *Arch. Elektrotechn.* **12**, 1 (1923).
- SIEBECK, F. H.: Über eine Gattung von Curven vierten Grades, welche mit den elliptischen Funktionen zusammenhängen. *J. reine angew. Math.* **57**, 359 (1860).
- THOMSON, J. J.: Recent researches in electricity and magnetism, Chap. 3. London: Oxford Univ. Press 1893.
- WEBER, E.: Die konforme Abbildung in der elektrischen Festigkeitslehre. *Arch. Elektrotechn.* **17**, 174 (1926).
- WRIGHT, C. E.: Note on the potential and attraction of rectangular bodies. *Phil. Mag.* **10**, 110 (1930).

*For Section V*

- [19] PANOFSKY, W. K. H., and M. PHILLIPS: Classical electricity and magnetism, p. 166. Reading, Mass.: Addison-Wesley Publ. Co. 1955.
- MORSE, P. M. and H. FESHBACH: Methods of theoretical physics. New York: McGraw-Hill Book Co. 1953.
- MOON, P., and D. E. SPENCER: Foundations of electrodynamics. Princeton, N. J.: D. Van Nostrand Co. 1960. — Field theory for engineers. Princeton, N. J.: D. Van Nostrand Co. 1961. — The meaning of the vector Laplacian. *J. Franklin Inst.* **256**, 551 (1953). — TEM waves in cylindrical systems. *J. Franklin Inst.*, **256**, 325 (1953).
- [20] SPENCER, D. E.: Separation of variables in electromagnetic theory. *J. Appl. Phys.* **22**, 386 (1951).

*For Section VI*

- [21] KLEIN, F.: Vorlesungen über lineare Differentialgleichungen der zweiten Ordnung. Bearbeitet von E. RITTER. Göttingen 1894.
- [22] BÖCHER, M.: Über die Reihenentwicklungen der Potentialtheorie. Leipzig: B. G. Teubner 1894.
- [23] INCE, E. L.: Ordinary differential equations, Chap. 20. London: Longmans, Green & Co. 1927.

- [24] MOON, P., and D. E. SPENCER: On the specification of Bôcher equations. *J. Franklin Inst.* **260**, 41 (1955).
- [25] MOON, P., and D. E. SPENCER: On the classification of the ordinary differential equations of field theory. *Quart. Appl. Math.* **14**, 1 (1956).
- [26] MOON, P., and D. E. SPENCER: Field theory for engineers, Chap. 5. Princeton, N. J.: D. Van Nostrand Co. 1961.

*For Section VII*

- [27] FROBENIUS, G.: Über die Integration der linearen Differentialgleichungen durch Reihen. *J. reine angew. Math.* **76**, 214 (1873).
- [28] FORSYTH, A. R.: Theory of differential equations. Cambridge: Cambridge Univ. Press 1902; New York: Dover Publications 1959, Vol. IV.
- INCE, E. L.: Ordinary differential equations. London: Longmans, Green & Co. 1927; New York: Dover Publications 1959, p. 396.
- [29] MOON, P., and D. E. SPENCER: Series solutions of Bôcher equations. (To be published.)
- [30] NIELSEN, N.: Handbuch der Theorie der Cylinderfunktionen. Leipzig: B. G. Teubner 1904.
- WATSON, G. N.: Theory of Bessel functions, p. 64. Cambridge: Cambridge Univ. Press 1944.
- [31] STURM, C., and J. LIOUVILLE: Extrait d'un mémoire sur le développement des fonctions en séries. *J. de Math.* **2**, 220 (1837).
- LIOUVILLE, J.: Premier mémoire sur la théorie des équations différentielles linéaires et sur le développement des fonctions en séries. *J. de Math.* **3**, 561 (1838).
- CHURCHILL, R. V.: Fourier series and boundary value problems, p. 46. New York: McGraw-Hill Book Co. 1941.
- [32] STRATTON, J. A.: Electromagnetic theory. New York: McGraw-Hill Book Co. 1941.
- RAMO, S., and J. R. WHINNERY: Fields and waves in modern radio. New York: John Wiley & Sons 1953.
- SMYTHE, W. R.: Static and dynamic electricity. New York: McGraw-Hill Book Co. 1950.
- SOUTHWORTH, C.: Principles and applications of waveguide transmission. Princeton, N. J.: D. Van Nostrand Co. 1950.
- MARCUVITZ, N.: Waveguide handbook. New York: McGraw-Hill Book Co. 1951.
- McLACHLAN, N. W.: Bessel functions for engineers. London: Oxford Univ. Press 1934. — Theory and applications of Mathieu functions. London: Oxford Univ. Press 1947.
- AHARONI, J.: Antennae. London: Oxford Univ. Press 1946.
- KRAUS, J. D.: Antennas. New York: McGraw-Hill Book Co. 1950.
- SILVER, S.: Microwave antenna theory and design. New York: McGraw-Hill Book Co. 1949.
- OLLENDORFF, F.: Potentialfelder der Elektrotechnik. Berlin: Springer 1932.
- SLATER, J. C.: Microwave electronics. Princeton, N. J.: D. Van Nostrand Co. 1950.
- FANO, R. M., L. J. CHU and R. B. ADLER: Electromagnetic fields, energy, and forces. New York: John Wiley & Sons 1960.
- ADLER, R. B., L. J. CHU and R. M. FANO: Electromagnetic energy transmission and radiation. New York: John Wiley & Sons 1960.
- WAIT, J. R.: Electromagnetic radiation from cylindrical structures. New York: Pergamon Press 1959.
- MORSE, P. M., and H. FESHBACH: Methods of theoretical physics. New York: McGraw-Hill Book Co. 1953.

CARSLAW, H. S., and J. C. JAEGER: Conduction of heat in solids. London: Oxford Univ. Press 1947.

INGERSOLL, L. R., O. J. ZOBEL and A. C. INGERSOLL: Heat conduction, with engineering and geological applications. New York: McGraw-Hill Book Co. 1948.

MUSKAT, M.: The flow of homogeneous fluids through porous media. New York: McGraw-Hill Book Co. 1937.

BORN, M., and E. WOLF: Principles of optics. New York: Pergamon Press 1959.

#### Weber functions

[33] WEBER, H.: Über die Integration der partiellen Differentialgleichung:  $\epsilon^2 u/\partial x^2 + \partial^2 u/\partial y^2 + k^2 u = 0$ . Math. Ann. **1**, 1 (1869).

[34] HERMITE, C.: Sur un nouveau développement en série des fonctions. C. R. Acad. Sci., Paris **58**, 93 (1864).

[35] WHITTAKER, E. T., and G. N. WATSON: Modern analysis, p. 347. Cambridge: Cambridge Univ. Press 1946.

INCE, E. L.: Ordinary differential equations, p. 159. London: Longmans, Green & Co. 1927.

WHITTAKER, E. T.: On the functions associated with the parabolic cylinder in harmonic analysis. Proc. London Math. Soc. **35**, 417 (1903).

WATSON, G. N.: The harmonic functions associated with the parabolic cylinder. Proc. London Math. Soc. **8**, 393 (1910).

MILNE, A.: On the equation of the parabolic cylinder functions. Proc. Edinburgh Math. Soc. **32**, 2 (1914).

HILLE, E.: On the zeros of the functions of the parabolic cylinder. Ark. Math., Astron. Fys. **18**, 26 (1924).

WELLS, C. P., and R. D. SPENCE: The parabolic cylinder functions. J. Math. Phys. **24**, 51 (1945).

DARWIN, C. G.: On Weber's function. Quart. J. Mech. Appl. Math. **2**, 311 (1949).

MILLER, J. C. P.: On the choice of standard solutions for a homogeneous linear differential equation of the second order. Quar. J. Mech. Appl. Math. **3**, 225 (1950). — A method for the determination of converging factors, applied to the asymptotic expansions for the parabolic cylinder functions. Proc. Cambridge Phil. Soc. **48**, 243 (1952).

MAGNUS, W., and F. OBERHETTINGER: Formeln und Sätze für die speziellen Funktionen der mathematischen Physik. Berlin: Springer 1943; New York: Chelsea Publ. Co. 1949.

ERDÉLYI, A., W. MAGNUS, F. OBERHETTINGER, and F. G. TRICOMI: Higher transcendental functions, Vol. II, p. 115. New York: McGraw-Hill Book Co. 1953.

#### Tables of Weber functions

[36] JANKE, E., u. F. EMDE: Tafeln höherer Funktionen, 4. Aufl. p. 27. Leipzig: B. G. Teubner 1948. Tables of the confluent hypergeometric function  $F(\frac{1}{2}n, \frac{1}{2}; x)$  and related functions. Washington, D. C.: Nat. Bu. Stds. 1949. Tables of Weber parabolic cylinder functions. London: H. M. Stationery Office 1955.

#### Bessel functions

[37] BESEL, F. W.: Untersuchung des Teils der planetarischen Störung, welcher aus Bewegung der Sonne entsteht. Berl. Abh. **1** (1824).

WATSON, G. N.: Theory of Bessel functions. Cambridge: Cambridge Univ. Press 1944.

WHITTAKER, E. T., and G. N. WATSON: Modern analysis, Chap. 17. Cambridge: Cambridge Univ. Press 1946.

GRAY, A., G. B. MATHEWS and T. M. MACROBERT: A treatise on Bessel functions. London: Macmillan Co. 1931.

MC LACHLAN, N. W.: Bessel functions for engineers. London: Oxford Univ. Press 1941.

NEUMANN, F.: Beiträge zur Theorie der Kugelfunktionen. Leipzig: B. G. Teubner 1878.

HEINE, H. E.: Handbuch der Kugelfunktionen. Berlin: G. Reimer 1878.

NIELSEN, N.: Handbuch der Theorie der Cylinderfunktionen. Leipzig: B. G. Teubner 1904.

PRASAD, G.: A treatise on spherical harmonics and the functions of Bessel and Lamé. 2 vols. Benares: S. C. Chatterji 1932.

LENSE, J.: Kugelfunktionen. Leipzig: Akad. Verlagsges. 1950.

BYERLY, W. E.: Fourier's series and spherical, cylindrical, and ellipsoidal harmonics. Boston: Ginn & Co. 1893.

POCKELS, F.: Über die partielle Differentialgleichung  $Au + K^2u = 0$  und deren Auftreten in der mathematischen Physik. Leipzig: B. G. Teubner 1891.

#### *Tables of Bessel functions*

[38] BATEMAN, H., and R. C. ARCHIBALD: A guide to tables of Bessel functions. Math. Tables and Other Aids to Comp. **1**, 205 (1944).

FLETCHER, A., J. C. P. MILLER and L. ROSENHEAD: Index of mathematical tables. London: Sci. Computing Service 1946.

JAHNKE, E., and F. EMDE: Tables of functions, p. 126. New York: Dover Publications 1943.

WATSON, G. N.: Theory of Bessel functions, p. 666. Cambridge: Cambridge Univ. Press 1944.

GRAY, A., G. B. MATHEWS and T. M. MACROBERT: A treatise on Bessel functions. London: Macmillan Co. 1931.

MCLACHLAN, N. W.: Bessel functions for engineers. London: Oxford Univ.

Press 1941. British Assn. for Adv. Sci.: Bessel functions, Vol. 6 and 10.

Cambridge: Cambridge Univ. Press 1937 and 1952. Harvard Computation Lab.: Tables of the Bessel functions of the first kind, Vols. 3 to 14. Cambridge, Mass.: Harvard Univ. Press 1947 to 1951. — Tables of the modified Hankel functions of order one-third and their derivatives. Cambridge, Mass.: Harvard Univ. Press 1945.

CAMBI, E.: Tables of Bessel functions of the first kind, to all significant orders.

New York: Dover Publications 1948. Nat. Bu. Stds.: Table of  $J_0(z)$  and  $J_1(z)$  for complex arguments. New York: Columbia Univ. Press 1947. —

Tables of spherical Bessel functions. New York: Columbia Univ. Press 1947. — Tables of Bessel functions  $Y_0(z)$  and  $Y_1(z)$  for complex arguments.

New York: Columbia Univ. Press 1950. — Tables of Bessel functions of fractional order, 2 vols. New York: Columbia Univ. Press 1948.

ONOUE, M.: Tables of modified quotients of Bessel functions of the first kind and real and imaginary arguments. New York: Columbia Univ. Press 1958. — Tables of the Bessel functions  $Y_0(x)$ ,  $Y_1(x)$ ,  $K_0(x)$ ,  $K_1(x)$ ,  $0 \leq x \leq 1$ .

Washington, D. C.: Nat. Bu. Stds. 1948.

[39] BAER, C.: Die Funktion des parabolischen Cylinders. Gymnasialprogramm, Cüstrin, 1883. — Parabolische Koordinaten. Frankfurt 1888.

#### *Mathieu functions*

[40] MATHIEU, E.: Mémoire sur le mouvement vibratoire d'une membrane de forme elliptique. J. de Math. **13**, 137 (1868).

[41] CURTIS, M. F.: The existence of the functions of the elliptic cylinder. Ann. of Math. **20**, 23 (1918).

HUMBERT, P.: Fonctions de Lamé et fonctions de Mathieu. (Mém. des sci. math.) Paris: Gauthier-Villars 1926.

DHAR, S. C.: Elliptic cylinder functions of second kind. Bull. Calcutta Math. Soc. **18**, 11 (1927).

INCE, E. L.: The elliptic cylinder functions of the second kind. Proc. Edinburgh Math. Soc. **33**, 2 (1914). — Researches into the characteristic numbers of the Mathieu equation. Proc. Roy. Soc. Edinburgh **46**, 20, 316 (1925); **47**, 294 (1927). — Ordinary differential equations, p. 175. London: Longmans, Green & Co. 1927. — Tables of elliptic-cylinder functions. Proc. Roy. Soc. Edinburgh **52**, 355 (1932).

- GOLDSTEIN, S.: Mathieu functions. *Trans. Cambridge Phil. Soc.* **23**, 303 (1927). — The second solution of Mathieu's differential equation. *Proc. Cambridge Phil. Soc.* **24**, 223 (1928).
- VARMA, R. S.: On Mathieu functions. *J. Indian Math. Soc.* **19**, 49 (1931).
- BARROW, W. L.: Untersuchen über den Heulsummer. *Ann. d. Phys.* **11**, 147 (1931).
- STRUTT, M. J. O.: Lamésche, Mathieusche und verwandte Funktionen in Physik und Technik. Berlin: Springer 1932.
- WHITTAKER, E. T., and G. N. WATSON: Modern analysis, Chap. 19. Cambridge: Cambridge Univ. Press. 1946.
- BLANCH, G.: Computation of Mathieu functions. *J. Math. Phys.* **25**, 1 (1946).
- MCLACHLAN, N. W.: Mathieu functions and their classification. *J. Math. Phys.* **25**, 209 (1946).
- [42] MCLACHLAN, N. W.: Theory and application of Mathieu functions, p. 19. London: Oxford Univ. Press 1947.
- [43] MORSE, P. M., and H. FESHBACH: Methods of mathematical physics. New York: McGraw-Hill Book Co. 1953.
- MEIXNER, J., u. F. W. SCHÄRKE: Mathieusche Funktionen und Sphäroidfunktionen. Berlin: Springer 1954.
- CAMPBELL, R.: Théorie générale de l'équation de Mathieu. Paris: Masson & Cie. 1955.

*Tables of Mathieu functions*

- [44] FLETCHER, A. F., J. C. P. MILLER and L. ROSENHEAD: An index of mathematical tables, p. 328. London: Sci. Comp. Service 1946.
- STRATTON, J. A., P. M. MORSE, L. J. CHU and R. A. HUTNER: Elliptic cylinder and spheroidal wave functions. New York: John Wiley & Sons 1941. Tables relating to Mathieu functions. New York: Columbia Univ. Press 1951.
- JANKE, E., u. F. EMDE: Tafeln höherer Funktionen. Leipzig: B. G. Teubner 1948.
- INCE, E. L.: Tables of elliptic-cylinder functions. *Proc. Roy. Soc. Edinburgh* **52**, 355 (1932).
- GOLDSTEIN, S.: Mathieu functions. *Trans. Cambridge Phil. Soc.* **23**, 303 (1927).

*Legendre functions*

- [45] LEGENDRE, A. M.: Sur l'attraction des sphéroïdes. *Mém. Math. Phys.* **10** (1785).
- HOBSON, E. W.: The theory of spherical and ellipsoidal harmonics. Cambridge: Cambridge Univ. Press 1931.
- HEINE, E.: Kugelfunktionen. Berlin: G. Reimer 1878.
- BYERLY, W. E.: Fourier's series and spherical, cylindrical, and ellipsoidal harmonics. Boston: Ginn & Co. 1893.
- PRASAD, G.: Spherical harmonics and the functions of Bessel and Lamé, 2 vols. Benares: S. C. Chatterji 1930.
- WHITTAKER, E. T., and G. N. WATSON: Modern analysis, Chap. 15. Cambridge: Cambridge Univ. Press 1946.
- MACROBERT, T. M.: Spherical harmonics. New York: Dover Publications 1948.
- ROBIN, L.: Fonctions sphériques de Legendre et fonctions sphéroidales, 3 vols. Paris: Gauthier-Villars 1959.

*Tables of Legendre functions*

- [46] DAVIS, H. T.: A bibliography and index of mathematical tables. Evanston, Ill. 1949.
- JAHNKE, E., and F. EMDE: Tables of functions, 3rd Edition p. 119. Leipzig: B. G. Teubner 1938.
- FOWLE, F. E.: Smithsonian physical tables, p. 66. Washington, D.C. 1933. Nat. Bu. Stds.: Table of associated Legendre functions. New York: Columbia Univ. Press 1945.
- MURSI, Z.: Table of Legendre associated functions. Cairo: Fouad I Univ. 1941.

TALLQUIST, H.: Tafeln der Kugelfunktionen. *Acta Soc. Sci. Fenn.* **32**, No. 6 (1906); **33**, Nos. 4 and 9 (1908); **6**, Nos. 3 and 10 (1932); **2**, No. 4 (1937); **2**, No. 11 (1938).

STRATTON, J. A., P. M. MORSE, L. J. CHU, J. D. C. LITTLE, and F. J. CORBÁTO: Spheroidal wave functions. New York: John Wiley & Sons 1956.

FLAMMER, C.: Spheroidal wave functions. Stanford, Cal.: Stanford Univ. Press 1957.

### *Lamé functions*

[47] LAMÉ, G.: Leçons sur les coordonnées curvilignes et leurs diverses applications. Paris: Mallet-Bachelier 1859.

TODHUNTER, I.: An elementary treatise on Laplace's functions, Lamé's functions, and Bessel's functions. London: Macmillan Co. 1875.

HEINE, H. E.: Handbuch der Kugelfunktionen. Berlin: G. Reimer 1878.

KLEIN, F.: Über Lamésche Funktionen. *Math. Ann.* **18**, 237 (1881).

BYERLY, W. E.: Fourier's series and spherical, cylindrical, and ellipsoidal harmonics. Boston: Ginn & Co. 1893.

NIVEN, W. D.: On ellipsoidal harmonics. *Phil. Trans. Roy. Soc. Lond.* **182**, 231 (1892).

MACLAUREN, R.: On the solutions of the equation  $(\nabla^2 + K^2)\psi = 0$  in elliptic coordinates and their physical applications. *Trans. Cambridge Phil. Soc.* **17**, 41 (1898).

DARWIN, G. H.: Ellipsoidal harmonic analysis. *Phil. Trans. Roy. Soc. Lond.* **197**, 461 (1901).

HUMBERT, P.: Fonctions de Lamé et fonctions de Mathieu. Paris: Gauthier-Villars 1926.

HOBSON, E. W.: The theory of spherical and ellipsoidal harmonics, Chap. 11. Cambridge: Cambridge Univ. Press 1931.

PRASAD, G.: A treatise on spherical harmonics and the functions of Bessel and Lamé, 2 vols. Benares: S. C. Chatterji 1932.

STRUTT, M. J. O.: Lamésche, Mathieusche und verwandte Funktionen in Physik und Technik. Berlin: Springer 1932.

[48] WANGERIN, A.: Über die Reduktion der Gleichung  $\partial^2 V/\partial x^2 + \partial^2 V/\partial y^2 + \partial^2 V/\partial z^2 = 0$  auf gewöhnliche Differentialgleichungen, *Mber. Akad. Wiss. Berlin* 152 (1878).

[49] HEINE, H. E.: Handbuch der Kugelfunktionen. Berlin: G. Reimer 1878.

## APPENDIX

### SYMBOLS USED IN THE TEXT

$A, B, C, D$  = constants.

$A_i$  = coefficients in series expansion of  $[(z - z_0) P(z)]$ , Section VII.

$\bar{A}_0, \bar{A}_1, \dots$  = constants in  $Q$ -term of a Bôcher equation.

$\mathcal{A}$  = area.

$a$  = distance to focus.

$a, b, c, \{$  = constants.  
 $a_1, a_2, a_3 \}$

$\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3 \}$  = unit vectors.  
 $\mathbf{a}_x, \mathbf{a}_y, \mathbf{a}_z \}$

$B_i$  = coefficients in series expansion of  $[(z - z_0)^2 Q(z)]$ , Section VII.

$\mathcal{B}_p^q$  = Baer function.

$b_1, b_2$  = roots of indicial equation, § 7.02.

$C_n$  = coefficients in series expansion of  $Z$ .

$\mathcal{C}_p^q$  = Baer function.

$c$  = a constant.

$\text{ce}_m$  = a Mathieu function.

$D_j(\beta) = (\beta - b_2) \cdot C_j(\beta)$ , § 7.01.

$\mathbf{E}$  = a vector.

$E_1, E_2, E_3$  = components of the vector  $\mathbf{E}$ .

$E_n^q$  = ordinary Lamé polynomial.

$\mathcal{E}_p^q$  = generalized Lamé function.

$e = 2.71828 \dots$

$F_n^q$  = ordinary Lamé function of second kind.

$\mathcal{F}_p^q$  = generalized Lamé function of the second kind.

$f_i$  = a function of  $u^i$ .

$\text{fe}_m$  = a Mathieu function.

$g_{11}, g_{22}, g_{33}$  = metric coefficients, § 1.04.

$g = g_{11} g_{22} g_{33}$ .

$\text{ge}_m$  = a Mathieu function.

$h$  = a constant.

$i = \sqrt{-1}$ .

$\mathcal{J}_p$  = Bessel function of first kind.

$K = \int_0^{\pi/2} \frac{d\varphi}{(1 - k^2 \sin^2 \varphi)^{1/2}}$ , complete elliptic integral.

$K' = \int_0^{\pi/2} \frac{d\varphi}{(1 - k'^2 \sin^2 \varphi)^{1/2}}$ , complete elliptic integral.

$k = b_1 - b_2$ , § 7.02.

$k$  = modulus of elliptic function.

$k' = (1 - k^2)^{\frac{1}{2}}$ .

$L$  = an operator, Section VII.

$M_{ij}$  = cofactor of element  $\Phi_{ij}$  in the Stäckel matrix.

$m, n$  = integers.

$N_n$  =  $n$ th norm.

$P_n(z)$  = ordinary Legendre polynomial.

$P_n^m(z)$  = Legendre associated function.

$P(z)$  = coefficient in Bôcher equation, Section VI.

$\mathcal{P}_p^q$  = generalized Legendre functions of first kind.

$p, q$  = constants, not necessarily integers.

$Q_n(z)$  = ordinary Legendre function of second kind.

$Q_n^m(z)$  = Legendre associated function.

$Q$  = a function of  $u^1, u^2, u^3$  used in  $R$ -separation (Section IV).

$Q(z)$  = coefficient in Bêcher equation.

$\mathcal{Q}_p^q$  = generalized Legendre function of second kind.

$R(r)$  = a function of  $r$ .

$R$  = a function of  $u^1, u^2, u^3$  used in  $R$ -separation (Section IV).

$r, \theta, \lambda$  = conical coordinates (Table 1.09).

$r, \theta, \psi$  = spherical coordinates (Table 1.05).

$r, \psi, z$  = circular-cylinder coordinates (Table 1.02).

$S$  = Stäckel determinant.

$[S]$  = Stäckel matrix.

$\mathcal{S}_p^q$  = Wangerin function.

$s$  = distance.

$\text{se}_m$  = a Mathieu function.

$T$  = a function of time.

$\mathcal{T}_p^q$  = Wangerin function.

$t$  = time.

$U^i$  = a function of  $u^i$ .

$\mathcal{U}_p^q$  = Heine function.

$u^1, u^2, u^3$  = general coordinates.

$u(z), v(z), w(z)$  = parameters in the Sturm-Liouville equation (§ 7.07).

$V$  = volume.

$\mathcal{V}_p^q$  = Heine function.

$\mathcal{W}_e, \mathcal{W}_o$  = Weber functions.

$w = u + iv$ , a complex number (Section II).

$X, Y, Z$  = functions of  $x, y, z$ , respectively.

$x, y, z$  = rectangular coordinates.

$\mathcal{Y}_n$  = Bessel function of the second kind.

$Z$  = a function of  $z$ .

$Z_1, Z_2$  = independent solutions of a Bôcher equation.

$\mathcal{Z}_p$  = generic designation of Bessel functions, including Bessel functions of first, second, and third kinds and their linear combinations.

$z_0$  = singular point.

$z = x + iy$ , a complex number.

$\bar{z}$  = complex conjugate of  $z$  (Section II).

- $\alpha_j$  = separation constant.  
 $\beta$  = a variable (Section VII).  
 $\Gamma$  = gamma function (Section VII).  
 $\gamma$  = a constant.  
 $\gamma = 0.5772157\dots$ , Euler's constant (Section VII).  
 $\Delta_j$  = a determinant.  
 $\zeta$  = independent variable.  
 $H(\eta)$  = a function of  $\eta$ .  
 $\eta$  = a coordinate in elliptic-cylinder coordinates (and in other systems).  
 $\eta, \theta, \lambda$  = ellipsoidal coordinates (Table 1.10).  
 $\eta, \psi, z$  = elliptic-cylinder coordinates (Table 1.03), and other cylindrical systems (Section III).  
 $\Theta$  = a function of  $\theta$ .  
 $\theta$  = angle from  $z$ -axis.  
 $\varkappa$  = constant in Helmholtz equation.  
 $\Lambda$  = a function of  $\lambda$ .  
 $M(\mu)$  = a function of  $\mu$ .  
 $\mu, \nu$  = coordinates.  
 $\mu, \nu, z$  = parabolic-cylinder coordinates (Table 1.04), and other cylindrical systems (Section III).  
 $\mu, \nu, \psi$  = parabolic coordinates (Table 1.08), and other rotational systems (Section IV).  
 $\mu, \nu, \lambda$  = paraboloidal coordinates (Table 1.11).  
 $N(\nu)$  = a function of  $\nu$ .  
 $\xi$  = a coordinate in alternative circular-cylinder coordinates and in alternative spherical coordinates.  
 $\pi = 3.14159\dots$   
 $\Phi_{ij}$  = element in Stäckel matrix.  
 $\varphi$  = scalar potential.  
 $\Psi$  = a function of  $\psi$ .  
 $\psi$  = angle about the  $z$ -axis.  
 $\Omega$  = a solution of Laplace's equation (Section I).  
 $\nabla^2$  = scalar Laplacian.  
 $\hat{\nabla}^2$  = vector Laplacian.

## AUTHOR INDEX

- |   |   |
|---|---|
| <p>Adler, R. B. 221<br/>         Aharoni, J. 221<br/>         Andronescu, P. 219<br/>         Archibald, R. C. 223</p> <p>Baer, C. 223<br/>         Bairstow, L. 219<br/>         Barrow, W. L. 224<br/>         Bateman, H. 217, 223<br/>         Bessel, F. W. 222<br/>         Blanch, G. 224<br/>         Blaschke, W. 217<br/>         Bôcher, M. 49, 144, 217,<br/>             220<br/>         Born, M. 222<br/>         Byerly, W. E. 217, 223,<br/>             224, 225</p> <p>Cambi, E. 223<br/>         Campbell, R. 224<br/>         Carslaw, H. S. 222<br/>         Casey, M. 218<br/>         Chu, L. J. 221, 224, 225<br/>         Churchill, R. V. 218, 219,<br/>             221<br/>         Coolidge, J. L. 218<br/>         Corbáto, F. J. 225<br/>         Courant, R. 218<br/>         Curtis, M. F. 223</p> <p>Darboux, G. 1, 217, 218<br/>         Darwin, C. G. 222<br/>         Darwin, G. H. 225<br/>         Davis, H. T. 224<br/>         Dhar, S. C. 223</p> <p>Eisenhart, L. P. 1, 217<br/>         Emde, F. 187, 222, 223,<br/>             224<br/>         Erdélyi, A. 222</p> <p>Fano, R. M. 221<br/>         Feshbach, H. 218, 220, 221,<br/>             224<br/>         Flammer, C. 225<br/>         Fletcher, A. F. 223, 224<br/>         Fogelsang, W. 217<br/>         Forsyth, A. R. 221<br/>         Fowle, F. E. 224<br/>         Frank, P. 218<br/>         Frobenius, G. 221</p> | <p>Glauert, H. 219<br/>         Goldstein, S. 224<br/>         Gray, A. 222, 223<br/>         Greenhill, G. 219<br/>         Grösser, W. 219</p> <p>Heine, H. E. 223, 224,<br/>             225<br/>         Hermite, C. 222<br/>         Hilbert, D. 218<br/>         Hildebrand, F. B. 218<br/>         Hille, E. 222<br/>         Hobson, E. W. 224, 225<br/>         Holtzmüller, O. 219<br/>         Humbert, P. 223, 225<br/>         Hutner, R. A. 224</p> <p>Ince, E. L. 144, 220, 221,<br/>             222, 223, 224<br/>         Ingwersoll, A. C. 222<br/>         Ingwersoll, L. R. 222</p> <p>Jaeger, J. C. 222<br/>         Jahnke, E. 187, 223, 224<br/>         Jeans, J. H. 218<br/>         Jeffreys, B. S. 218<br/>         Jeffreys, H. 218</p> <p>Kehren, E. 219<br/>         Kellogg, O. D. 218<br/>         Klein, F. 49, 144, 218, 220,<br/>             225<br/>         Knight, R. C. 219<br/>         Knopp, K. 219<br/>         Kober, H. 219<br/>         Kraus, J. D. 221</p> <p>Labus, J. 219<br/>         Lamé, G. 1, 217, 225<br/>         Legendre, A. M. 224<br/>         Lense, J. 218, 223<br/>         Levi-Civita, T. 217, 220<br/>         Levy, H. 220<br/>         Liouville, J. 176, 219, 221<br/>         Little, J. D. C. 225<br/>         Love, A. E. H. 219</p> <p>McLachlan, N. W. 220, 221,<br/>             222, 223, 224<br/>         Maclaurin, R. 225</p> |
|---|---|
- MacRobert, T. M. 222, 223,  
     224  
 Magnus, W. 222  
 Marcuvitz, N. 221  
 Mathews, G. B. 222, 223  
 Mathieu, E. 223  
 Maxwell, J. C. 218  
 Meixner, J. 224  
 Meyer, E. 220  
 Michell, J. H. 219  
 Miller, J. C. P. 222, 223,  
     224  
 Milne, A. 222  
 Mises, R. v. 218  
 Moon, P. 217, 218, 219, 220,  
     221  
 Morse, P. M. 218, 220, 221,  
     224, 225  
 Morton, W. B. 220  
 Müller, E. 217  
 Murnaghan, F. D. 218  
 Mursi, Z. 224  
 Muskat, M. 222
- Nehari, Z. 219  
 Neumann, F. 222  
 Nicholson, J. W. 220  
 Nielson, N. 173, 175, 221,  
     223  
 Niven, W. D. 225
- Oberhettinger, F. 222  
 Ollendorff, F. 219, 221  
 Onoe, M. 223  
 Osgood, W. F. 219
- Page, W. M. 220  
 Panofsky, W. K. H. 220  
 Petersohn, H. 220  
 Phillips, M. 220  
 Pockels, F. 223  
 Pohlhausen, K. 219  
 Poole, E. G. C. 220  
 Poritsky, H. 220  
 Prasad, G. 223, 224, 225
- Ramo, S. 221  
 Richmond, H. W. 220  
 Robertson, H. P. 217  
 Robin, L. 224

- |   |                                       |  |
|---|---------------------------------------|--|
| Rogowski, W. 220                          | Stratton, J. A. 218, 221,<br>224, 225 | Wait, J. R. 221                        |
| Rosenhead, L. 223, 224                    | Strutt, M. J. O. 224,<br>225          | Walker, M. 219                         |
| Rothe, R. 219                             | Sturm, C. 176, 221                    | Wangerin, A. 219, 225                  |
| Schärke, F. W. 224                        | Tait, P. G. 219                       | Watson, G. N. 221, 222,<br>223, 224    |
| Schmidt, H. 218                           | Tallquist, H. 225                     | Weber, E. 218, 219, 220                |
| Siebeck, F. H. 220                        | Thomson, J. J. 220                    | Weber, H. 222                          |
| Silver, S. 221                            | Thomson, W. 218, 219                  | Webster, A. G. 218                     |
| Slater, J. C. 221                         | Titchmarsh, E. C. 219                 | Wells, C. P. 222                       |
| Smythe, W. R. 218,<br>221                 | Todhunter, I. 225                     | Whinnery, J. R. 221                    |
| Sommerfeld, A. 218                        | Townsend, E. J. 219                   | Whittaker, E. T. 183, 222,<br>223, 224 |
| Southworth, C. 221                        | Tricomi, F. G. 222                    | Wolf, E. 222                           |
| Spence, R. D. 222                         | Turriere, E. 217                      | Wright, C. E. 220                      |
| Spencer, D. E. 217, 218,<br>219, 220, 221 | Varma, R. S. 224                      | Wrinch, D. M. 219                      |
| Stäckel, P. 217                           |                                       | Zobel, O. J. 222                       |

## SUBJECT INDEX

The numbers in brackets {} refer to the classification of differential equations, Section VI

|  |  |  |
|--|--|--|
| Alternative cylindrical system 13<br>tabulated data 13   | Canonical equations 148<br>Cap-cyclide coordinates, tabulated data 103, 132  | oblate spheroidal 31<br>parabolic 34<br>conical 37<br>ellipsoidal 40<br>paraboloidal 44  |
| Alternative spherical system 25<br>tabulated data 25   | Cardioids 51, 52<br>map 58<br>Cardioid coordinates, tabulated data 99, 107<br>Cardioid-cylinder coordinates 79, 86<br>Cassinian-ovals 51, 52<br>map 62<br>Cassinian-oval cylinder coordinates 80, 88<br>Cauchy-Riemann equations 49<br>Circles 51, 52<br>map 61<br>Circular-cylinder coordinates 7, 12, 80<br>alternative system 13<br>vector Helmholtz equation 138 | Coordinate systems in which Laplace equation is $R$ -separable 97<br>tangent-sphere 99, 104<br>cardioid 99, 107<br>bisphere 100, 110<br>toroidal 101, 112<br>inverse prolate spheroidal 101, 115<br>inverse oblate spheroidal 101, 119   |
| Area 2   |  | 6-sphere 122   |
| Asymptotic expansion, Weber functions 183<br>Bessel functions 191  |  | bi-cyclide 102, 124<br>flat-ring cyclide 102, 126<br>disk-cyclide 103, 129<br>cap-cyclide 103, 132   |
| Baer equation 148, 150<br>wave 148, 150  |  | Coordinate systems with rotational symmetry 97   |
| Baer functions 194<br>{113} 46, 157<br>orthogonality 179   |  | spherical 24, 100<br>prolate spheroidal 28, 101<br>oblate spheroidal 31, 101<br>parabolic 34, 99<br>tangent-sphere 99, 104<br>cardioid 99, 107<br>hyperbolic 100<br>bispherical 100, 110<br>toroidal 101, 112<br>inverse prolate spheroidal 101, 115<br>inverse oblate spheroidal 101, 119 |
| Baer wave functions 194<br>{114} 48, 157<br>orthogonality 179, 196   |  | 6-sphere 122   |
| Bessel equation 146, 148, 149, 178, 185<br>wave 148, 149, 178  |  | bi-cyclide 102, 124  |
| Bessel functions 185, 194<br>first kind 172, 187<br>second kind 173, 187<br>Hankel 190<br>wave, 1st kind 172, 185<br>wave, 2nd kind 175, 185<br>orthogonality 178, 192 |  | flat-ring cyclide 102, 126   |
| Bessel functions, {24} 14, 15, 16, 27, 35, 40, 105, 108, 139, 140, 141, 142, 155<br>{14} 14, 15, 16, 17, 36, 106, 109, 154<br>{06} 23, 24                              |  | disk-cyclide 103, 129  |
| Bessel wave functions, {26} 36, 143, 155, 171<br>{16} 36, 154  |  | cap-cyclide 103, 132   |
| Bibliography 217   |  | Critically damped 5  |
| Bi-cyclide coordinates, tabulated data 102, 124  |  | Curl 2   |
| Bi-cylinder coordinates 81, 89   |  | in rectangular coordinates 9   |
| Bipolar circles 51, 53<br>map 64   |  | circular cylinder 12   |
| Bi-spherical coordinates 100, 110  |  | elliptic cylinder 17   |
| Bôcher equations 144<br>classification 148   |  | parabolic cylinder 21  |
|  |  | spherical 24   |
|  |  | prolate spheroidal 28  |

|   |  |  |
|---|--|--|
| parabolic-cylinder 21                           | hyperbolic 79, 87                                    | alternative circular-cylinder 13                               |
| spherical 25                                    | rose 80, 87  | elliptic-cylinder 17   |
| alternative spherical 25                        | Cassinian oval 80, 88                                | parabolic-cylinder 21  |
| prolate spheroidal 29                           | inverse Cassinian oval 81, 88                        | spherical 25   |
| oblate spheroidal 32                            | bi-cylinder 81, 89                                   | alternative spherical 25                                       |
| parabolic 35                                    | Maxwell 81, 89                                       | prolate spheroidal 29  |
| conical 38                                      | logarithmic 81, 90                                   | oblate spheroidal 31   |
| ellipsoidal 42                                  | In tan 82, 90  | parabolic 35   |
| paraboloidal 45                                 | In cosh 82, 91                                       | conical 38   |
| tangent-cylinder 86                             | inverse elliptic 82, 91                              | ellipsoidal 42   |
| cardioid-cylinder 86                            | sn 83, 92  | paraboloidal 45  |
| hyperbolic-cylinder 87                          | cn 83, 92  | tangent-cylinder 86  |
| rose-cylinder 87                                | inverse sn 83, 93                                    | cardioid-cylinder 86   |
| Cassinian-oval 88                               | In sn 84, 94   | hyperbolic-cylinder 87   |
| inverse Cassinian-oval 88                       | In cn 84, 94   | rose-cylinder 87   |
| bi-cylinder 89                                  | zeta 85, 95  | Cassinian-oval 88  |
| Maxwell-cylinder 89                             | Cylindrical systems 7, 49, 77                        | inverse Cassinian-oval 88                                      |
| logarithmic-cylinder 90                         | vector Helmholtz equation 136, 138                   | bi-cylinder 89   |
| In tan-cylinder 90                              | Damped wave equation 4                               | Maxwell-cylinder 89  |
| In cosh-cylinder 91                             | Definite integrals, for Bessel functions 190         | logarithmic-cylinder 90  |
| inverse elliptic cylinder 92                    | for Legendre functions 208                           | In tan-cylinder 90   |
| sn-cylinder 92                                  | Differential equations, partial 3                    | In cosh-cylinder 91  |
| cn-cylinder 93                                  | ordinary 144   | inverse elliptic cylinder 91                                   |
| inverse sn-cylinder 93                          | Differential equations 144                           | sn-cylinder 92   |
| In sn-cylinder 94                               | specification 145                                    | cn-cylinder 93   |
| In cn-cylinder 95                               | classification 148                                   | inverse sn-cylinder 93   |
| zeta-cylinder 95                                | Weber 148, 154, 178, 182                             | In sn-cylinder 94  |
| tangent sphere 105                              | Bessel 146, 149, 155, 171, 178, 185                  | In cn-cylinder 95  |
| cardioid 108                                    | Baer 150, 157, 179, 194                              | zeta-cylinder 95   |
| bi-spherical 111                                | Mathieu 150, 156                                     | tangent-sphere 104   |
| toroidal 114                                    | Legendre 151, 155, 156, 157, 158, 159, 179, 180, 201 | cardioid 107   |
| inverse prolate spheroidal 117                  | Lamé 151, 152, 159, 160, 180, 210                    | bi-spherical 111   |
| inverse oblate spheroidal 120                   | Wangerin 152, 161, 181, 212                          | toroidal 113   |
| 6-sphere 123                                    | Heine 152, 162, 181, 214                             | inverse prolate spheroidal 117                                 |
| bi-cyclide 125                                  | Differentiation of Bessel functions 191              | inverse oblate spheroidal 120                                  |
| flat-ring cyclide 128                           | Diffusion equation 4                                 | 6-sphere 122   |
| disk-cyclide 131                                | Dirichlet conditions 176                             | bi-cyclide 125   |
| cap-cyclide 134                                 | Disk-cyclide coordinates, tabulated data 103, 129    | flat-ring cyclide 128  |
| Curvilinear squares 49                          | Distance 1   | disk-cyclide 131   |
| Cyclide coordinate systems, bi-cyclide 102, 124 | Divergence 2   | cap-cyclide 134  |
| flat-ring cyclide 102, 126                      | in rectangular coordinates 9                         | Eigenvalue 176   |
| disk-cyclide 103, 129                           | circular-cylinder 12                                 | Electric field distorted by introduction of spheroid 8, 47, 48 |
| cap-cyclide 103, 132                            | Ellipses 51, 53                                      | Elementary functions 148                                       |
| Cyclides 49                                     | map 69   | orthogonality 178  |
| Cylindrical coordinates 50, 78                  |  | Ellipses 51, 53  |
| circular 12, 80                                 |  | map 69   |
| elliptic 17, 82                                 |  |  |
| parabolic 21, 79                                |  |  |
| tangent 79, 86                                  |  |  |
| cardioid 79, 86                                 |  |  |

Ellipsoidal coordinates 7  
     tabulated data 40  
 Elliptic-cylinder coordinates 7, 82  
     tabulated data 17  
     vector Helmholtz equation 140  
 Example of electrostatic field 8  
 Exponential functions, {04} 10, 11, 14, 15, 19, 22, 105, 106, 111, 112, 123  
 Flat-ring cyclide coordinates, tabulated data 102, 126  
 4-leaf roses 51, 52  
     map 60  
 Frobenius method 168  
 Functions 163  
     Weber 182  
     Bessel 171, 173, 185  
     Baer 194  
     Mathieu 197  
     Legendre 201  
     Lamé 210  
     Wangerin 212  
     Heine 214  
 Gradient 2  
     in rectangular coordinates 9  
     circular-cylinder 12  
     alternative circular cylinder 13  
     elliptic cylinder 17  
     parabolic cylinder 21  
     spherical 25  
     alternative spherical 25  
     prolate spheroidal 28  
     oblate spheroidal 31  
     parabolic 35  
     conical 38  
     ellipsoidal 42  
     paraboloidal 45  
     tangent-cylinder 86  
     cardioid-cylinder 86  
     hyperbolic-cylinder 87  
     rose-cylinder 87  
     Cassinian-oval 88  
     inverse Cassinian-oval 88  
     bi-cylinder 89  
     Maxwell-cylinder 89  
     logarithmic-cylinder 90  
     ln tan-cylinder 90

ln cosh-cylinder 91  
     inverse elliptic cylinder 91  
     sn-cylinder 92  
     cn-cylinder 93  
     inverse sn-cylinder 93  
     ln sn-cylinder 94  
     ln cn-cylinder 95  
     zeta-cylinder 95  
     tangent-sphere 104  
     cardioid 107  
     bi-spherical 110  
     toroidal 113  
     inverse prolate spheroidal 117  
     inverse oblate spheroidal 120  
     6-sphere 122  
     bi-cyclide 125  
     flat-ring cyclide 128  
     disk-cyclide 131  
     cap-cyclide 134  
 Heine equation 148, 152, 162, 181  
 Heine functions 214  
     {1222} 126, 162  
     orthogonality 181, 216  
 Helmholtz equation 1  
     scalar 4  
     separation in rectangular coordinates 10, 11  
         circular-cylinder 15, 16, 17  
         elliptic-cylinder 19, 20  
         parabolic-cylinder 23, 24  
         spherical 27  
         prolate spheroidal 30  
         oblate spheroidal 33, 34  
         parabolic 36  
         conical 39, 40  
         ellipsoidal 43, 44  
         paraboloidal 47, 48  
     vector 136  
     separation in cylindrical systems 136  
         rotational 137  
         rectangular coordinates 138  
         circular-cylinder 138  
         elliptic-cylinder 140  
         parabolic-cylinder 140  
         spherical 141  
         prolate spheroidal 142  
         oblate spheroidal 142  
         parabolic 143  
 Hermite polynomials 182  
 Hyperbolas 51, 52  
     map 59  
 Hyperbolic-cylinder coordinates 79, 87  
 Hyperbolic coordinates 100  
 Hypergeometric function 183  
 Indical equation 166  
 Integral representation, Weber functions 184  
     Bessel functions 190  
     Legendre functions 208  
 Integration of Bessel functions 192  
 Inverse Cassinian-ovals 51, 53  
     map 63  
 Inverse Cassinian-oval coordinates 81, 88  
 Inverse elliptic-cylinder coordinates 82, 91  
 Inverse ellipses 51, 53  
     map 70  
 Inverse oblate spheroidal coordinates 101, 119  
 Inverse prolate spheroidal coordinates 101, 115  
 Inverse sn curves 51, 54  
     map 73  
 Inverse sn-cylinder coordinates 83, 93  
 Inversion 49  
 Lamé equation 148, 152, 160, 180  
     wave 148, 151  
 Lamé functions 210  
     wave functions 210  
     polynomials 211  
     orthogonality 180, 212  
     {1112} 38, 39, 40, 43, 160  
     {1111} 39, 159  
 Lamé wave functions  
     {1113} 44, 160  
     orthogonality 180  
 Laplace equation 1, 3  
     cylindrical coordinates 78  
     separation in rectangular coordinates 9, 10  
         circular-cylinder 13, 14  
         elliptic-cylinder 18, 19  
         parabolic-cylinder 22, 23  
         spherical 26, 27

|   |  |  |
|---|--|--|
| prolate spheroidal 29,<br>30<br>oblate spheroidal 32,<br>33<br>parabolic 35, 36<br>conical 38, 39<br>ellipsoidal 42, 43<br>paraboloidal 46<br>tangent-sphere 105<br>cardioid 108<br>bispherical 111<br>toroidal 114<br>inverse prolate spher-<br>oidal 117<br>inverse oblate spher-<br>oidal 120<br>6-sphere 123<br>bi-cyclide 125<br>flat-ring cyclide 128<br>disk-cyclide 131<br>cap-cyclide 134  | ln cn-cylinder 95<br>zeta-cylinder 95<br>tangent-sphere 105<br>cardioid 108<br>bi-spherical 111<br>toroidal 114<br>inverse prolate spher-<br>oidal 117<br>inverse oblate spher-<br>oidal 120<br>6-sphere 123<br>bi-cyclide 125<br>flat-ring cyclide 128<br>disk-cyclide 131<br>cap-cyclide 134   | ln sn curves 51, 54<br>map 74<br>ln sn-cylinder coordinates<br>84, 94<br>ln tan curves 51, 53<br>map 67<br>ln tan cylinder coordinates<br>82, 90<br>logarithmic curves 51, 53<br>map 66<br>logarithmic-cylinder co-<br>ordinates 81, 90  |
| Laplacian, scalar 3<br>in rectangular coordina-<br>tes 9<br>circular-cylinder 12<br>alternative circular-<br>cylinder 13<br>elliptic-cylinder 17<br>parabolic-cylinder 21<br>spherical 25<br>alternative spherical<br>25<br>prolate spheroidal 29<br>oblate spheroidal 32<br>parabolic 35<br>conical 38<br>ellipsoidal 42<br>paraboloidal 45<br>tangent-cylinder 86<br>cardioid-cylinder 86<br>hyperbolic-cylinder 87<br>rose-cylinder 87<br>Cassinian-oval 88<br>inverse Cassinian-oval<br>88<br>bi-cylinder 89<br>Maxwell-cylinder 89<br>logarithmic-cylinder<br>90 | Laplacian, vector 3<br>Legendre equation 146, 148,<br>151, 158, 179<br>wave 148, 151<br>Legendre functions 201<br>wave 201<br>degenerate cases 202<br>ordinary functions 203<br>polynomials, first kind<br>205<br>polynomials, second kind<br>206<br>associated functions 207<br>orthogonality, 179, 209<br>$\{222\}$ 26, 27, 29, 32, 111,<br>114, 117, 118, 120, 121,<br>158<br>$\{220\}$ 26<br>$\{112\}$ 26, 27, 29, 33,<br>112, 114, 115, 118, 121,<br>142, 155 | Mathieu equation 150<br>Mathieu functions 197<br>$\{113\}$ 18, 19, 20, 156<br>orthogonality 200<br>Maxwell curves 51, 53<br>map 65<br>Maxwell-cylinder coordi-<br>nates 81, 89<br>Metric coefficients 1<br>in rectangular coordina-<br>tes 9<br>circular-cylinder 12,<br>52, 80<br>alternative circular-<br>cylinder 13<br>elliptic-cylinder 18, 53,<br>82<br>parabolic-cylinder 21,<br>52<br>spherical 25<br>alternative spherical 25<br>prolate spheroidal 28,<br>101<br>oblate spheroidal 31,<br>101<br>parabolic 34, 99<br>conical 37<br>ellipsoidal 41<br>paraboloidal 45<br>tangent-cylinder 52,<br>79, 86<br>cardioid-cylinder 52,<br>79, 86<br>hyperbolic-cylinder 52,<br>79, 87<br>rose-cylinder 52, 80, 87<br>Cassinian-oval cylin-<br>der 52, 80, 88<br>inverse Cassinian oval<br>cylinder 53, 81, 88<br>bipolar-cylinder 53,<br>81, 89<br>Maxwell-cylinder 53,<br>81, 89 |
| ln tan-cylinder 90<br>ln cosh-cylinder 91<br>inverse elliptic-cylin-<br>der 92<br>sn-cylinder 92<br>cn-cylinder 93<br>inverse sn-cylinder 93<br>ln sn-cylinder 94   | ln cn curves 51, 55<br>map 75<br>ln cn cylinder coordinates<br>84, 94<br>ln cosh curves 51, 53<br>map 68<br>ln cosh cylinder coordina-<br>tes 82, 91   |  |

|   |  |  |
|---|--|--|
| logarithmic-cylinder<br>53, 81, 90        | Parabolas 51, 52<br>map 57   | inverse prolate spheroidal 101, 115  |
| ln tan-cylinder 53, 82,<br>90             | Parabolic coordinates 2, 3,<br>6, 7, 99<br>tabulated data 34<br>vector Helmholtz equa-<br>tion 143                 | inverse oblate spheroidal<br>101, 119  |
| ln cosh-cylinder 53, 82,<br>91            | Parabolic-cylinder coordi-<br>nates 7, 79, 52<br>map 57<br>tabulated data 21<br>vector Helmholtz equa-<br>tion 140 | 6-sphere 122<br>bi-cyclide 102, 124<br>flat-ring cyclide 102, 126<br>disk-cyclide 103, 129<br>cap-cyclide 103, 132 |
| inverse elliptic cylin-<br>der 53, 82, 91 | Paraboloidal coordinates 7<br>tabulated data 44  | Rotational systems 7, 49,<br>96<br>vector Helmholtz equa-<br>tion 137, 141   |
| sn-cylinder 54, 83, 92                    | Partial differential equa-<br>tions 3  | separability conditions<br>98  |
| cn-cylinder 54, 83, 92                    | Periodic solutions,<br>of Mathieu equation 197   | R-separation 96  |
| inverse sn-cylinder 54,<br>83, 93         | Poisson equation 3   | Scalar Helmholtz equa-<br>tion 4   |
| ln sn-cylinder 54, 84,<br>94              | Poles 145  | Scalar Laplacian 3   |
| ln cn-cylinder 55, 84,<br>94              | Polynomials,<br>Hermite 182<br>Legendre 205<br>Lamé 211  | Scalar potential 4   |
| zeta-cylinder 55, 85,<br>95               | Potential 4  | Schwarz-Christoffel me-<br>thod 49   |
| tangent-sphere 99, 104                    | Power functions, {04} 14,<br>109<br>{22} 26, 38, 154<br>{01} 27  | Separation 5, 96<br>table 98   |
| cardioid 99, 107                          | Prolate spheroidal coordi-<br>nates 7, 101<br>tabulated data 28<br>vector Helmholtz equa-<br>tion 142              | Separation constants 6   |
| bispherical 100, 110                      | Rectangular coordinates 7<br>tabulated data 9<br>vector Helmholtz equa-<br>tion 138                                | Separation equations 6, 152  |
| toroidal 101, 113                         | Recursion formulas,<br>Weber functions 184<br>Bessel functions 191   | Separation of Helmholtz<br>equation,<br>in rectangular coordina-<br>tes 10   |
| inverse prolate spher-<br>oidal 101, 116  | Regular singularity 165,<br>168  | circular-cylinder 15   |
| inverse oblate spher-<br>oidal 101, 119   | Rose coordinates 80, 87  | elliptic-cylinder 19   |
| 6-sphere 122                              | Rotational coordinates 50,<br>96, 97   | parabolic-cylinder 23  |
| bi-cyclide 102, 125                       | spherical 24, 100  | spherical 27   |
| flat-ring cyclide 102,<br>128             | prolate spheroidal 28, 101   | prolate spheroidal 30  |
| disk-cyclide 103, 130                     | oblate spheroidal 31, 101  | oblate spheroidal 33   |
| cap-cyclide 103, 133                      | parabolic 34, 99   | parabolic 36   |
| Neumann conditions 176                    | tangent-sphere 99, 104   | conical 39   |
| Oblate spheroidal coordina-<br>tes 7, 101 | cardioid 99, 107   | ellipsoidal 43   |
| tabulated data 31                         | hyperbolic 100   | paraboloidal 47  |
| vector Helmholtz equa-<br>tion 142        | bispherical 100, 110   | Separation of Laplace's<br>equation,<br>in rectangular coordina-<br>tes 9  |
| Order of a pole 145                       | toroidal 101, 112  | circular-cylinder 13   |
| Ordinary point 165                        |  | elliptic-cylinder 18   |
| Orthogonal coordinate sys-<br>tem 1       |  | parabolic-cylinder 22  |
| Orthogonal map 50                         |  | spherical 26   |
| Orthogonality 175                         |  | prolate spheroidal 29  |
| table 178                                 |  | oblate spheroidal 32   |
| Weber functions 184                       |  | parabolic 35   |
| Bessel 192                                |  | conical 38   |
| Baer 196                                  |  | ellipsoidal 42   |
| Mathieu 200                               |  | paraboloidal 46  |
| Legendre 209                              |  | tangent-sphere 105   |
| Lamé 212                                  |  | cardioid 108   |
| Wangerin 214                              |  |  |
| Heine 216                                 |  |  |
| Oscillatory 5                             |  |  |
| Overdamped 5                              |  |  |

|  |  |   |
|--|--|---|
| bi-sphere 111<br>toroid 114<br>inverse prolate spheroidal 117<br>inverse oblate spheroidal 120<br>6-sphere 123<br>bi-cyclide 124<br>flat-ring cyclide 128<br>disk-cyclide 131<br>cap-cyclide 134   | cardioid 108<br>bi-sphere 111<br>toroid 114<br>inverse prolate 117<br>inverse oblate 120<br>6-sphere 123<br>bi-cyclide 126<br>flat-ring cyclide 129<br>disk-cyclide 132<br>cap-cyclide 135   | Tangent-circles 51, 52<br>map 56<br>Tangent-cylinder coordinates, 52, 77, 79, 86<br>map 56<br>Tangent-sphere coordinates 99, 104<br>Toroidal coordinates 97, 101<br>tabulated data 112<br>Transformations of Bôcher equations 146<br>Transformations in complex plane 49, 51<br>power functions 52<br>exponential 52<br>logarithmic 53<br>hyperbolic 53<br>elliptic 54, 55                  |
| Series solutions 165<br>Simple separation 5<br>necessary and sufficient conditions 7<br>definition 96<br>Singular point 165<br>Singularity 165<br>6-sphere coordinates 122<br>sn curves 51, 54<br>map 71<br>sn-cylinder coordinates 83, 92<br>Solutions of Helmholtz equation,<br>in rectangular coordinates 11<br>circular-cylinder 15<br>elliptic-cylinder 20<br>parabolic-cylinder 23<br>spherical 27<br>prolate spheroidal 30<br>oblate spheroidal 33<br>parabolic 36<br>conical 40<br>ellipsoidal 44<br>paraboloidal 48 | Spherical coordinates 7, 100<br>tabulated data 24<br>vector Helmholtz equation 141<br>Spheroid in uniform field 8<br>Stäckel matrix 5<br>determinant 5<br>for cylindrical systems 7<br>for rotational systems 7<br>in rectangular coordinates 9<br>circular-cylinder 10<br>alternative circular-cylinder 13<br>elliptic-cylinder 17<br>parabolic-cylinder 21<br>spherical 24<br>alternative spherical 25<br>prolate spheroidal 25<br>oblate spheroidal 31<br>parabolic 34<br>conical 37<br>ellipsoidal 41<br>paraboloidal 45<br>tangent-sphere 104<br>cardioid 107<br>bispherical 110<br>toroidal 112<br>inverse prolate spheroidal 116<br>inverse oblate spheroidal 119<br>6-sphere 122<br>bi-cyclide 125<br>flat-ring cyclide 127<br>disk-cyclide 130<br>cap-cyclide 132 | Transmission-line equation 4<br>Trigonometric functions, {04} 10, 11, 14, 15, 16, 17, 18, 19, 20, 22, 23, 26, 27, 29, 30, 32, 33, 35, 36, 40, 78, 105, 106, 108, 109, 111, 114, 115, 118, 121, 123, 126, 129, 132, 135, 139, 140, 141<br>other forms 153<br>Two-dimensional field 50<br>Vector Helmholtz equation 136<br>Vector Laplacian 3, 136, 137<br>Vector wave equation 4<br>Volume 2 |
| Solutions of Laplace equation,<br>in rectangular coordinates 10<br>circular-cylinder 14<br>elliptic-cylinder 18<br>parabolic-cylinder 22<br>spherical 26<br>prolate spheroidal 29<br>oblate spheroidal 32<br>parabolic 35<br>conical 38<br>ellipsoidal 43<br>paraboloidal 46<br>cylindrical coordinates 78<br>tangent-sphere coordinates 105   | Sturm-Liouville systems 176<br>Surfaces 1<br>Symbols, table 226  | Wangerin equation 148, 152, 161, 181, 212<br>Wangerin functions 212<br>{1122} 129, 132, 135, 161<br>orthogonality 181, 214<br>Wave equation 4<br>Weber equation 148, 149, 154, 178<br>Weber functions 154, 182<br>{06} 22, 23<br>orthogonality 178, 184<br>Weighting function 176<br>Zeta function curves 51, 55<br>map 76<br>Zeta-cylinder coordinates 85, 95                              |