



UPPSALA
UNIVERSITET

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Bayesian Statistics and Data Analysis

Lecture 7

Måns Magnusson

Department of Statistics, Uppsala University
Thanks to Aki Vehtari, Aalto University



UPPSALA
UNIVERSITET

Assignment 6

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Difficulties with stan: R-studio can check stan syntax + take small steps
- Repetition - there is a purpose behind this.
- \hat{R} as a posterior statistic of interest



UPPSALA
UNIVERSITET

- **Hierarchical models**
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Section 1

Hierarchical models



- Hierarchical models


- Rats example
- Factory example
- 8 schools example

- Exchangeability

Hierarchical model

- Example: Treatment effectiveness

- in hospital j the survival probability is θ_j
- observations y_{ij} tell whether patient i survived in hospital





- Hierarchical models


- Rats example
- Factory example
- 8 schools example

- Exchangeability


Hierarchical model

- Example: Treatment effectiveness

- in hospital j the survival probability is θ_j
- observations y_{ij} tell whether patient i survived in hospital



- sensible to assume that θ_j are similar



- natural to think that θ_j have common population distribution
- θ_j is not directly observed and the population distribution is unknown




● Hierarchical models

- Rats example
- Factory example
- 8 schools example

● Exchangeability

Hierarchical model: terms

Lvl 1: observations given parameters $p(y_{ij}|\theta_j)$



Joint posterior

$$\begin{aligned} p(\theta, \tau|y) &\propto p(y|\theta, \tau)p(\theta, \tau) \\ &\propto p(y|\theta)p(\theta|\tau)p(\tau) \end{aligned}$$



● Hierarchical models


- Rats example
- Factory example
- 8 schools example

● Exchangeability

Hierarchical model: terms

Lvl 1: observations given parameters $p(y_{ij}|\theta_j)$

Lvl 2: parameters given hyperparameters $p(\theta_j|\tau)$



Joint posterior

$$\begin{aligned} p(\theta, \tau|y) &\propto p(y|\theta, \tau)p(\theta, \tau) \\ &\propto p(y|\theta)p(\theta|\tau)p(\tau) \end{aligned}$$



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Comparisons

- "Separate model" (model with separate/independent effects)

θ_1	θ_2	\dots	θ_n
\downarrow	\downarrow		\downarrow
y_1	y_2		y_n




- Hierarchical models

- Rats example
- Factory example
- 8 schools example


- Exchangeability

Comparisons

- "Separate model" (model with separate/independent effects)



- "Joint/pooled model" (model with a common effect / pooled model)







- **Hierarchical models**
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Comparisons


- "Separate model" (model with separate/independent effects)



- "Joint/pooled model" (model with a common effect / pooled model)



- Hierarchical model






Predictive distributions for hierarchical models

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Two types of predictive distributions
 1. A new observation in **an existing group**
 2. A new observation in **a new group**





UPPSALA
UNIVERSITET

Hierarchical binomial model: rats

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability
- Medicine testing
- Type F344 female rats in control group given placebo
 - count how many get endometrial stromal polyps
 - familiar binomial model example




- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Medicine testing
- Type F344 female rats in control group given placebo
 - count how many get endometrial stromal polyps
 - familiar binomial model example
- Experiment has been repeated 71 times

0/20	0/20	0/20	0/20	0/20	0/20	0/20	0/19	0/19	0/19
0/19	0/18	0/18	0/17	1/20	1/20	1/20	1/20	1/19	1/19
1/18	1/18	2/25	2/24	2/23	2/20	2/20	2/20	2/20	2/20
2/20	1/10	5/49	2/19	5/46	3/27	2/17	7/49	7/47	3/20
3/20	2/13	9/48	10/50	4/20	4/20	4/20	4/20	4/20	4/20
4/20	10/48	4/19	4/19	4/19	5/22	11/46	12/49	5/20	5/20
6/23	5/19	6/22	6/20	6/20	6/20	16/52	15/46	15/47	9/24
4/14									




Pooled model



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability



Separate model



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability




Hierarchical binomial model: rats

- Hierarchical binomial model for rats
prior parameters α and β are unknown

$$\theta_j | \alpha, \beta \sim \text{Beta}(\theta_j | \alpha, \beta)$$

$$y_j | n_j, \theta_j \sim \text{Bin}(y_j | n_j, \theta_j)$$



- Joint posterior $p(\theta_1, \dots, \theta_J, \alpha, \beta | y)$
 - multiple parameters




Hierarchical binomial model: rats

- Hierarchical binomial model for rats
prior parameters α and β are unknown

$$\theta_j | \alpha, \beta \sim \text{Beta}(\theta_j | \alpha, \beta)$$

$$y_j | n_j, \theta_j \sim \text{Bin}(y_j | n_j, \theta_j)$$



- Joint posterior $p(\theta_1, \dots, \theta_J, \alpha, \beta | y)$
 - multiple parameters
 - factorize $\prod_{j=1}^J p(\theta_j | \alpha, \beta, y) p(\alpha, \beta | y)$



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Population prior $\text{Beta}(\theta_j | \alpha, \beta)$
- Hyperprior $p(\alpha, \beta)$?
 - α, β both affect the location and scale
 - BDA3 (p. 110) has (vague) $p(\alpha, \beta) \propto (\alpha + \beta)^{-5/2}$



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability


- Population prior $\text{Beta}(\theta_j | \alpha, \beta)$
- Hyperprior $p(\alpha, \beta)$?
 - α, β both affect the location and scale
 - BDA3 (p. 110) has (vague) $p(\alpha, \beta) \propto (\alpha + \beta)^{-5/2}$
- What type of predictive distributions can we have?



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability


Hierarchical binomial model: rats

The marginal of α and β






Beta(α, β) given posterior draws of α and β



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability



Population distribution (prior) for θ_j



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability




UPPSALA
UNIVERSITET

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical binomial model: rats

Separate model






UPPSALA
UNIVERSITET


- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical binomial model: rats

Separate model



Hierarchical model






UPPSALA
UNIVERSITET


- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical binomial model: rats

Hierarchical model



Population distribution (prior) for θ_j






- Hierarchical models
 - Rats example
 - **Factory example**
 - 8 schools example
- Exchangeability

Hierarchical normal model: factory

- Factory has 6 machines which quality is evaluated
- Assume hierarchical model
 - each machine has its own (average) quality θ_j and **common variance** σ^2

$$\theta_j | \mu_P, \sigma_P^2 \sim \mathcal{N}(\mu_P, \sigma_P^2)$$

$$y_{ij} | \theta_j \sim \mathcal{N}(\theta_j, \sigma^2)$$




- Can be used to predict the future quality produced by each machine and quality produced by a new similar machine



- Hierarchical models
 - Rats example
 - **Factory example**
 - 8 schools example
- Exchangeability

Hierarchical normal model: factory

- Factory has 6 machines which quality is evaluated
- Assume hierarchical model
 - each machine has its own (average) quality θ_j and **own variance** σ_j^2






- Hierarchical models
 - Rats example
 - **Factory example**
 - 8 schools example
- Exchangeability

Hierarchical normal model: factory

- Factory has 6 machines which quality is evaluated
- Assume hierarchical model
 - each machine has its own (average) quality θ_j and **own variance** σ_j^2




- What type of predictive distributions can we have?



- Hierarchical models
 - Rats example
 - **Factory example**
 - 8 schools example
- Exchangeability

Hierarchical normal model: factory

- Factory has 6 machines which quality is evaluated
- Assume hierarchical model
 - each machine has its own (average) quality θ_j and **own variance** σ_j^2



- **What type of predictive distributions can we have?**
- Can be used to predict the future quality produced by each machine and quality produced by a new similar machine



UPPSALA
UNIVERSITET

Hierarchical normal model: 8 schools

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: SAT coaching effectiveness
 - in USA commonly used Scholastic Aptitude Test (SAT) is designed so that short term practice should not improve the results significantly
 - schools have anyway coaching courses
 - test the effectiveness of the coaching courses



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: SAT coaching effectiveness
 - in USA commonly used Scholastic Aptitude Test (SAT) is designed so that short term practice should not improve the results significantly
 - schools have anyway coaching courses
 - test the effectiveness of the coaching courses
- SAT
 - standardized multiple choice test
 - mean about 500 and standard deviation about 100
 - most scores between 200 and 800
 - different topics, e.g., V=Verbal, M=Mathematics
 - pre-test PSAT



UPPSALA
UNIVERSITET

Hierarchical normal model: 8 schools

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Effectiveness of the SAT coaching
 - students had made pre-tests PSAT-M and PSAT-V
 - part of students were coached
 - linear regression was used to estimate the coaching effect y_j for the school j (could be denoted with $\bar{y}_{.j}$, too) and variances σ_j^2
 - y_j approximately normally distributed, with variances assumed to be known based on about 30 students per school
 - data is group means and variances (not personal results)



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Effectiveness of the SAT coaching
 - students had made pre-tests PSAT-M and PSAT-V
 - part of students were coached
 - linear regression was used to estimate the coaching effect y_j for the school j (could be denoted with $\bar{y}_{.j}$, too) and variances σ_j^2
 - y_j approximately normally distributed, with variances assumed to be known based on about 30 students per school
 - data is group means and variances (not personal results)
- Data:

School	A	B	C	D	E	F	G	H
y_j	28	8	-3	7	-1	1	18	12
σ_j	15	10	16	11	9	22	20	28



Hierarchical normal model for group means

- J experiments, unknown θ_j and known σ^2

$$y_{ij}|\theta_j \sim \mathcal{N}(\theta_j, \sigma^2), \quad i = 1, \dots, n_j; \quad j = 1, \dots, J$$

- Group j sample mean and sample variance

$$\bar{y}_{.j} = \frac{1}{n_j} \sum_{i=1}^{n_j} y_{ij}$$

$$\sigma_j^2 = \frac{\sigma^2}{n_j}$$

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical normal model for group means

- J experiments, unknown θ_j and known σ^2

$$y_{ij}|\theta_j \sim \mathcal{N}(\theta_j, \sigma^2), \quad i = 1, \dots, n_j; \quad j = 1, \dots, J$$

- Group j sample mean and sample variance

$$\bar{y}_{.j} = \frac{1}{n_j} \sum_{i=1}^{n_j} y_{ij}$$

$$\sigma_j^2 = \frac{\sigma^2}{n_j}$$

- Use model

$$\bar{y}_{.j}|\theta_j \sim \mathcal{N}(\theta_j, \sigma_j^2)$$

this model can be generalized so that, σ_j^2 can be different from each other for other reasons than n_j




- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical normal model for group means

$$\theta_j | \mu, \tau \sim \mathcal{N}(\mu, \tau)$$


$$\bar{y}_j | \theta_j \sim \mathcal{N}(\theta_j, \sigma_j^2)$$





- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Separate model







Hierarchical normal model: 8 schools

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Separate model



Pooled model






- Hierarchical models

- Rats example
- Factory example
- 8 schools example


- Exchangeability

Hierarchical normal model: 8 schools


Separate model



Pooled model



Hierarchical model






UPPSALA
UNIVERSITET

Hierarchical normal model: 8 schools

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability







- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical normal model: 8 schools

Marginal posterior $p(\tau|y)$



Conditional means $E[\theta_j|\tau, y]$







- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Hierarchical normal model: 8 schools


Marginal posterior $p(\tau|y)$



Conditional means $E[\theta_j|\tau, y]$



Conditional standard deviations $sd[\theta_j|\tau, y]$





UPPSALA
UNIVERSITET

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

Section 2

Exchangeability



UPPSALA
UNIVERSITET

Exchangeability

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Justifies why we can use
 - a joint model for data
 - a joint prior for a set of parameters
- Less strict than independence (IID)
- IID \rightarrow exchangeability



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- *Exchangeability*

Parameters $\theta_1, \dots, \theta_J$ (or observations y_1, \dots, y_J) are exchangeable if the joint distribution p is invariant to the permutation of indices $(1, \dots, J)$

e.g.

$$p(\theta_1, \theta_2, \theta_3) = p(\theta_2, \theta_3, \theta_1)$$



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- *Exchangeability*

Parameters $\theta_1, \dots, \theta_J$ (or observations y_1, \dots, y_J) are exchangeable if the joint distribution p is invariant to the permutation of indices $(1, \dots, J)$

e.g.

$$p(\theta_1, \theta_2, \theta_3) = p(\theta_2, \theta_3, \theta_1)$$

Can we come up with a situation where this doesn't hold?



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- *Exchangeability*

Parameters $\theta_1, \dots, \theta_J$ (or observations y_1, \dots, y_J) are exchangeable if the joint distribution p is invariant to the permutation of indices $(1, \dots, J)$

e.g.

$$p(\theta_1, \theta_2, \theta_3) = p(\theta_2, \theta_3, \theta_1)$$

Can we come up with a situation where this doesn't hold?

- Exchangeability implies symmetry:
If there is no information which can be used *a priori* to separate θ_j from each other, we can assume exchangeability. ("Ignorance implies exchangeability")



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Exchangeability does not mean that the results of the experiments could not be different
 - e.g. if we know that the experiments have been in two different laboratories, and we know that the other laboratory has better conditions for the rats, but we do not know which experiments have been made in which laboratory
 - a priori experiments are exchangeable
 - model could have unknown parameter for the laboratory with a conditional prior for rats assumed to come from the same place (clustering model)



UPPSALA
UNIVERSITET

Exchangeability and additional information

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: bioassay
 - y_i number of dead animals are not exchangeable alone



UPPSALA
UNIVERSITET

Exchangeability and additional information

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: bioassay
 - y_i number of dead animals are not exchangeable alone
 - x_i dose is additional information



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: bioassay
 - y_i number of dead animals are not exchangeable alone
 - x_i dose is additional information
 - (x_i, y_i) exchangeable and logistic regression was used

$$p(\alpha, \beta | y, n, x) \propto \prod_{i=1}^n p(y_i | \alpha, \beta, n_i, x_i) p(\alpha, \beta)$$



UPPSALA
UNIVERSITET

Hierarchical exchangeability

- Hierarchical models

- Rats example
- Factory example
- 8 schools example

- Exchangeability

- Example: hierarchical rats example
 - all rats not exchangeable



UPPSALA
UNIVERSITET

Hierarchical exchangeability

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: hierarchical rats example
 - all rats not exchangeable
 - in a single laboratory rats exchangeable



UPPSALA
UNIVERSITET

Hierarchical exchangeability

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: hierarchical rats example
 - all rats not exchangeable
 - in a single laboratory rats exchangeable
 - laboratories exchangeable



UPPSALA
UNIVERSITET

Hierarchical exchangeability

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: hierarchical rats example
 - all rats not exchangeable
 - in a single laboratory rats exchangeable
 - laboratories exchangeable



UPPSALA
UNIVERSITET

Hierarchical exchangeability

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Example: hierarchical rats example
 - all rats not exchangeable
 - in a single laboratory rats exchangeable
 - laboratories exchangeable
 - hierarchical model can be used



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Conditional exchangeability
 - if y_i is connected to an additional information x_i , so that y_i are not exchangeable, but (y_i, x_i) exchangeable use joint model or conditional model $(y_i | x_i)$.



- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability

- Conditional exchangeability
 - if y_i is connected to an additional information x_i , so that y_i are not exchangeable, but (y_i, x_i) exchangeable use joint model or conditional model $(y_i | x_i)$.
- Partial exchangeability
 - if the observations can be grouped (a priori), then we can use a hierarchical model



Exchangeability

- The simplest form of the exchangeability (but not the only one) for the parameters θ is **conditional independence**

$$p(x_1, \dots, x_J | \theta) = \prod_{j=1}^J p(x_j | \theta)$$

- Hierarchical models
 - Rats example
 - Factory example
 - 8 schools example
- Exchangeability



- The simplest form of the exchangeability (but not the only one) for the parameters θ is **conditional independence**

$$p(x_1, \dots, x_J | \theta) = \prod_{j=1}^J p(x_j | \theta)$$

- Let $(x_n)_{n=1}^{\infty}$ to be an infinite sequence of exchangeable random variables. De Finetti's theorem then says that there is some random variable θ so that x_j are conditionally independent given θ , and joint density for x_1, \dots, x_J can be written in the *iid mixture* form

$$p(x_1, \dots, x_J) = \int \left[\prod_{j=1}^J p(x_j | \theta) \right] p(\theta) d\theta$$