



UPPSALA
UNIVERSITET

Bayesian Statistics and Data Analysis

Lecture 3

Måns Magnusson

Department of Statistics, Uppsala University
Thanks to Aki Vehtari, Aalto University

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



UPPSALA
UNIVERSITET

Assignment 2

- **Introduction**
- **Multiple parameter models**
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- **Bioassay example**
- **Additional examples**
 - Multinomial model
 - Multivariate Gaussian

- "Task 2b and 2d was unclear" (on our list now)



Assignment 2

- **Introduction**
- **Multiple parameter models**
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- **Bioassay example**
- **Additional examples**
 - Multinomial model
 - Multivariate Gaussian

- "Task 2b and 2d was unclear" (on our list now)
- "Simulate from the predictive distribution in two steps?"



Assignment 2

- **Introduction**
- **Multiple parameter models**
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- **Bioassay example**
- **Additional examples**
 - Multinomial model
 - Multivariate Gaussian

- "Task 2b and 2d was unclear" (on our list now)
- "Simulate from the predictive distribution in two steps?"



UPPSALA
UNIVERSITET

Practicalities

- **Introduction**
- **Multiple parameter models**
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- **Bioassay example**
- **Additional examples**
 - Multinomial model
 - Multivariate Gaussian
- **Master thesis proposals**
 - Deadline for interest today at 23.59.
 - I'll stay if you have questions.



UPPSALA
UNIVERSITET

Practicalities

- Introduction

- Multiple parameter models

- Marginalization

- Gaussian Model

- Example
- Conjugate prior

- Bioassay example

- Additional examples

- Multinomial model
- Multivariate Gaussian

- Master thesis proposals

- Deadline for interest today at 23.59.
- I'll stay if you have questions.

- Next week (Wednesday): Deadline mini-project proposal



UPPSALA
UNIVERSITET

Practicalities

- **Introduction**

- **Multiple parameter models**

- Marginalization

- **Gaussian Model**

- Example
- Conjugate prior

- **Bioassay example**

- **Additional examples**

- Multinomial model
- Multivariate Gaussian

- **Master thesis proposals**

- Deadline for interest today at 23.59.
- I'll stay if you have questions.


- **Next week (Wednesday): Deadline mini-project proposal**

- **Submit to the correct assignment - its anonymous**



Example of uncertainty in modeling

Data




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Example of uncertainty in modeling

Posterior mean




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Example of uncertainty in modeling


Predictive distribution given posterior mean





Example of uncertainty in modeling

Posterior draws




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Example of uncertainty in modeling

Posterior draws and predictive distribution




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Example of uncertainty in modeling

Posterior draws and predictive distribution



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



UPPSALA
UNIVERSITET

Monte Carlo and Posterior Draws

- **Introduction**
 - **Multiple parameter models**
 - Marginalization
 - **Gaussian Model**
 - Example
 - Conjugate prior
 - **Bioassay example**
 - **Additional examples**
 - Multinomial model
 - Multivariate Gaussian
- Assume we can get draws from $p(\theta | y)$



- **Introduction**
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Assume we can get draws from $p(\theta | y)$
- $\theta^{(s)}$ draws from $p(\theta | y)$ can be used
 - for visualization
 - to approximate expectations (integrals)

$$E_{p(\theta|y)}[\theta] = \int \theta p(\theta | y) \approx \frac{1}{S} \sum_{s=1}^S \theta^{(s)}$$



- **Introduction**
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Assume we can get draws from $p(\theta | y)$
- $\theta^{(s)}$ draws from $p(\theta | y)$ can be used
 - for visualization
 - to approximate expectations (integrals)

$$E_{p(\theta|y)}[\theta] = \int \theta p(\theta | y) \approx \frac{1}{S} \sum_{s=1}^S \theta^{(s)}$$

- to approximate uncertainty intervals for θ



UPPSALA
UNIVERSITET

- Introduction
- **Multiple parameter models**
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Section 2

Multiple parameter models



- Joint posterior distribution of multiple parameters

$$p(\theta_1, \theta_2 \mid y) \propto p(y \mid \theta_1, \theta_2)p(\theta_1, \theta_2)$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Joint posterior distribution of multiple parameters

$$p(\theta_1, \theta_2 | y) \propto p(y | \theta_1, \theta_2)p(\theta_1, \theta_2)$$

- Marginalization

$$p(\theta_1 | y) = \int p(\theta_1, \theta_2 | y) d\theta_2$$

$p(\theta_1 | y)$ is a marginal distribution

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Joint posterior distribution of multiple parameters

$$p(\theta_1, \theta_2 \mid y) \propto p(y \mid \theta_1, \theta_2)p(\theta_1, \theta_2)$$

- Marginalization

$$p(\theta_1 \mid y) = \int p(\theta_1, \theta_2 \mid y) d\theta_2$$

$p(\theta_1 \mid y)$ is a marginal distribution

- Goal is often to find marginal posterior of an interesting quantity
 - a **parameter** $p(\theta|y)$
 - a **potential observation** $p(\tilde{y}|y)$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Joint distribution of unknown future observation and parameters

$$\begin{aligned} p(\tilde{y}, \theta | y) &= p(\tilde{y} | \theta, y) p(\theta | y) \\ &= p(\tilde{y} | \theta) p(\theta | y) \quad (\text{often}) \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Joint distribution of unknown future observation and parameters

$$\begin{aligned}p(\tilde{y}, \theta | y) &= p(\tilde{y} | \theta, y)p(\theta | y) \\ &= p(\tilde{y} | \theta)p(\theta | y) \quad (\text{often})\end{aligned}$$

- Marginalization over posterior distribution

$$\begin{aligned}p(\tilde{y} | y) &= \int p(\tilde{y} | \theta)p(\theta | y)d\theta \\ &= \int p(\tilde{y}, \theta | y)d\theta\end{aligned}$$

$p(\tilde{y} | y)$ is a predictive distribution



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Observation model


$$\frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(y - \mu)^2\right)$$

- Uninformative prior

$$p(\mu, \sigma^2) \propto \sigma^{-2}$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian






- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Gaussian fit with posterior mean




$$p(\textcolor{red}{y} \mid \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(\textcolor{red}{y} - \mu)^2\right)$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Gaussian fit with posterior mean




$$p(y \mid \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(y - \mu)^2\right)$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Gaussian fit with posterior mean




$$p(y \mid \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(y - \mu)^2\right)$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Gaussians with posterior draw parameters




$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Gaussians with posterior draw parameters




$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Draws from the joint posterior distribution




$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian example

Draws from the joint posterior distribution




$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$



Joint posterior

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$




- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Joint posterior

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$

with $p(\mu, \sigma^2) \propto \sigma^{-2}$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian




- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma | y)$$

with $p(\mu, \sigma^2) \propto \sigma^{-2}$

$$p(\mu, \sigma^2 | y) \propto \sigma^{-2} \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(y_i - \mu)^2\right)$$






- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$

with $p(\mu, \sigma^2) \propto \sigma^{-2}$

$$p(\mu, \sigma^2 \mid y) \propto \sigma^{-n-2} \exp \left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2 \right)$$





- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian


Joint posterior

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma | y)$$

with $p(\mu, \sigma^2) \propto \sigma^{-2}$

$$\begin{aligned} p(\mu, \sigma^2 | y) &\propto \sigma^{-n-2} \exp \left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2 \right) \\ &= \sigma^{-n-2} \exp \left(-\frac{1}{2\sigma^2} \left[\sum_{i=1}^n (y_i - \bar{y})^2 + n(\bar{y} - \mu)^2 \right] \right) \end{aligned}$$

$$\text{where } \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$$






Joint posterior

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma | y)$$

with $p(\mu, \sigma^2) \propto \sigma^{-2}$



$$\begin{aligned} p(\mu, \sigma^2 | y) &\propto \sigma^{-n-2} \exp \left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2 \right) \\ &= \sigma^{-n-2} \exp \left(-\frac{1}{2\sigma^2} \left[\sum_{i=1}^n (y_i - \bar{y})^2 + n(\bar{y} - \mu)^2 \right] \right) \end{aligned}$$

$$\text{where } \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$$

$$= \sigma^{-n-2} \exp \left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2] \right)$$

$$\text{where } s^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$$



UPPSALA
UNIVERSITET

Gaussian: Completing the square

$$\sum_{i=1}^n (y_i - \mu)^2$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



$$\sum_{i=1}^n (y_i - \mu)^2$$
$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2)$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian: Completing the square

$$\sum_{i=1}^n (y_i - \mu)^2$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2)$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2 - \bar{y}^2 + \bar{y}^2 - 2y_i\bar{y} + 2y_i\bar{y})$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian: Completing the square

$$\sum_{i=1}^n (y_i - \mu)^2$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2)$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2 - \bar{y}^2 + \bar{y}^2 - 2y_i\bar{y} + 2y_i\bar{y})$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\bar{y} + \bar{y}^2) + \sum_{i=1}^n (\mu^2 - 2y_i\mu - \bar{y}^2 + 2y_i\bar{y})$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian: Completing the square

$$\sum_{i=1}^n (y_i - \mu)^2$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2)$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2 - \bar{y}^2 + \bar{y}^2 - 2y_i\bar{y} + 2y_i\bar{y})$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\bar{y} + \bar{y}^2) + \sum_{i=1}^n (\mu^2 - 2y_i\mu - \bar{y}^2 + 2y_i\bar{y})$$

$$\sum_{i=1}^n (y_i - \bar{y})^2 + n(\mu^2 - 2\bar{y}\mu - \bar{y}^2 + 2\bar{y}\bar{y})$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Gaussian: Completing the square

$$\sum_{i=1}^n (y_i - \mu)^2$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2)$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\mu + \mu^2 - \bar{y}^2 + \bar{y}^2 - 2y_i\bar{y} + 2y_i\bar{y})$$

$$\sum_{i=1}^n (y_i^2 - 2y_i\bar{y} + \bar{y}^2) + \sum_{i=1}^n (\mu^2 - 2y_i\mu - \bar{y}^2 + 2y_i\bar{y})$$

$$\sum_{i=1}^n (y_i - \bar{y})^2 + n(\mu^2 - 2\bar{y}\mu - \bar{y}^2 + 2\bar{y}\bar{y})$$


$$\sum_{i=1}^n (y_i - \bar{y})^2 + n(\bar{y} - \mu)^2$$




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu)$ and $p(\sigma^2)$

Joint posterior



Marginal of mu




$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma \mid y)$$




- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu)$ and $p(\sigma^2)$


Joint posterior



Marginal of mu



Marginal of sigma



$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma | y)$$

marginals

$$p(\mu | y) = \int p(\mu, \sigma | y) d\sigma$$

$$p(\sigma | y) = \int p(\mu, \sigma | y) d\mu$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Suppose we care about μ in the Gaussian model.
- Which parameter is a nuisance parameter?
- Why integrate it out instead of plug in one estimate?



Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$p(\sigma^2 | y) \propto \int p(\mu, \sigma^2 | y) d\mu$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$\begin{aligned} p(\sigma^2 | y) &\propto \int p(\mu, \sigma^2 | y) d\mu \\ &\propto \int \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\mu \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$\begin{aligned} p(\sigma^2 | y) &\propto \int p(\mu, \sigma^2 | y) d\mu \\ &\propto \int \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\mu \\ &\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \cdot \\ &\quad \int \exp\left(-\frac{n}{2\sigma^2} (\bar{y} - \mu)^2\right) d\mu \end{aligned}$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$\begin{aligned} p(\sigma^2 | y) &\propto \int p(\mu, \sigma^2 | y) d\mu \\ &\propto \int \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\mu \\ &\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \cdot \\ &\quad \int \exp\left(-\frac{n}{2\sigma^2} (\bar{y} - \mu)^2\right) d\mu \end{aligned}$$

Note! $\int \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y - \mu)^2\right) d\mu = 1$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$\begin{aligned}p(\sigma^2 | y) &\propto \int p(\mu, \sigma^2 | y) d\mu \\&\propto \int \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\mu \\&\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \cdot \\&\quad \int \exp\left(-\frac{n}{2\sigma^2} (\bar{y} - \mu)^2\right) d\mu\end{aligned}$$

Note!

$$\begin{aligned}&\int \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y - \mu)^2\right) d\mu = 1 \\&\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \sqrt{2\pi\sigma^2/n}\end{aligned}$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$\begin{aligned}p(\sigma^2 | y) &\propto \int p(\mu, \sigma^2 | y) d\mu \\&\propto \int \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\mu \\&\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \cdot\end{aligned}$$

$$\begin{aligned}&\int \exp\left(-\frac{n}{2\sigma^2} (\bar{y} - \mu)^2\right) d\mu \\ \text{Note!} \quad &\int \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y - \mu)^2\right) d\mu = 1 \\&\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \sqrt{2\pi\sigma^2/n} \\&\propto (\sigma^2)^{-(n+1)/2} \exp\left(-\frac{(n-1)s^2}{2\sigma^2}\right)\end{aligned}$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal $p(\mu|y)$ and $p(\sigma^2|y)$

Marginal posterior $p(\sigma^2 | y)$ (easier for σ^2 than σ)

$$\begin{aligned}
 p(\sigma^2 | y) &\propto \int p(\mu, \sigma^2 | y) d\mu \\
 &\propto \int \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\mu \\
 &\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \cdot
 \end{aligned}$$

$$\begin{aligned}
 &\int \exp\left(-\frac{n}{2\sigma^2} (\bar{y} - \mu)^2\right) d\mu \\
 \text{Note!} \quad &\int \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y - \mu)^2\right) d\mu = 1 \\
 &\propto \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} (n-1)s^2\right) \sqrt{2\pi\sigma^2/n} \\
 &\propto (\sigma^2)^{-(n+1)/2} \exp\left(-\frac{(n-1)s^2}{2\sigma^2}\right)
 \end{aligned}$$

$$p(\sigma^2 | y) = \text{Inv-}\chi^2(\sigma^2 | n-1, s^2)$$



Known mean

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

$$\sigma^2 \mid y \sim \text{Inv-}\chi^2(n, \nu)$$

$$\text{where } \nu = \frac{1}{n} \sum_{i=1}^n (y_i - \mu)^2$$



Known mean

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior

$$\sigma^2 \mid y \sim \text{Inv-}\chi^2(n, \nu)$$

$$\text{where } \nu = \frac{1}{n} \sum_{i=1}^n (y_i - \mu)^2$$

Unknown mean

- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

$$\sigma^2 \mid y \sim \text{Inv-}\chi^2(n - 1, s^2)$$

$$\text{where } s^2 = \frac{1}{n - 1} \sum_{i=1}^n (y_i - \bar{y})^2$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Marginal posterior $p(\mu | y)$


$$p(\mu | y) = \int_0^\infty p(\mu | \sigma^2, y) p(\sigma^2 | y) d\sigma^2$$

- Marginal posterior of μ , a mixture of normal distributions where mixing density is the marginal posterior of σ^2




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior



Marginal of sigma




Factorization

$$p(\mu, \sigma^2 | y) = p(\mu | \sigma^2, y) p(\sigma^2 | y)$$




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior



Marginal of sigma




Factorization

$$p(\mu, \sigma^2 | y) = p(\mu | \sigma^2, y) p(\sigma^2 | y)$$
$$(\sigma^2)^{(s)} \sim p(\sigma^2 | y)$$




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior



Marginal of sigma



Factorization

$$p(\mu, \sigma^2 | y) = p(\mu | \sigma^2, y) p(\sigma^2 | y)$$

$$(\sigma^2)^{(s)} \sim p(\sigma^2 | y)$$


$$p(\mu | (\sigma^2)^{(s)}, y) = \mathcal{N}(\mu | \bar{y}, (\sigma^2)^{(s)} / n)$$




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior


60
 - Exact contour plot — Cond. distribution of μ
 Sample from joint post. — Sample from the marg.



Cond distr of μ for 25 draws



Marginal of sigma



Factorization

$$p(\mu, \sigma^2 | y) = p(\mu | \sigma^2, y) p(\sigma^2 | y)$$

$$(\sigma^2)^{(s)} \sim p(\sigma^2 | y)$$


$$p(\mu | (\sigma^2)^{(s)}, y) = \mathcal{N}(\mu | \bar{y}, (\sigma^2)^{(s)} / n)$$




- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior


60
- Exact contour plot — Cond. distribution of μ
Sample from joint post. — Sample from the marg.



Cond distr of μ for 25 draws



Marginal of sigma



Factorization

$$p(\mu, \sigma^2 | y) = p(\mu | \sigma^2, y) p(\sigma^2 | y)$$

$$(\sigma^2)^{(s)} \sim p(\sigma^2 | y)$$


$$p(\mu | (\sigma^2)^{(s)}, y) = \mathcal{N}(\mu | \bar{y}, (\sigma^2)^{(s)} / n)$$

$$p(\mu | y) \approx \frac{1}{S} \sum_{s=1}^S \mathcal{N}(\mu | \bar{y}, (\sigma^2)^{(s)} / n)$$




- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian


Joint posterior



Cond. distr of μ



Marginal of sigma



Factorization

$$p(\mu, \sigma^2 | y) = p(\mu | \sigma^2, y) p(\sigma^2 | y)$$

$$(\sigma^2)^{(s)} \sim p(\sigma^2 | y)$$

$$p(\mu | (\sigma^2)^{(s)}, y) = \mathcal{N}(\mu | \bar{y}, (\sigma^2)^{(s)}/n)$$

$$p(\mu | y) \approx \frac{1}{S} \sum_{s=1}^S \mathcal{N}(\mu | \bar{y}, (\sigma^2)^{(s)})$$



Marginal posterior $p(\mu \mid y)$

$$p(\mu \mid y) = \int_0^\infty p(\mu, \sigma^2 \mid y) d\sigma^2$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu \mid y) &= \int_0^\infty p(\mu, \sigma^2 \mid y) d\sigma^2 \\ &\propto \int_0^\infty \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\sigma^2 \end{aligned}$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu \mid y) &= \int_0^\infty p(\mu, \sigma^2 \mid y) d\sigma^2 \\ &\propto \int_0^\infty \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\sigma^2 \end{aligned}$$

Transformation (integration by substitution)

$$A = (n-1)s^2 + n(\mu - \bar{y})^2$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu \mid y) &= \int_0^\infty p(\mu, \sigma^2 \mid y) d\sigma^2 \\ &\propto \int_0^\infty \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\sigma^2 \end{aligned}$$

Transformation (integration by substitution)

$$A = (n-1)s^2 + n(\mu - \bar{y})^2 \quad \text{and} \quad z = \frac{A}{2\sigma^2}$$

$$dz = \left(-\frac{A}{2(\sigma^2)^2}\right) d\sigma^2$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu \mid y) &= \int_0^\infty p(\mu, \sigma^2 \mid y) d\sigma^2 \\ &\propto \int_0^\infty \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\sigma^2 \end{aligned}$$

Transformation (integration by substitution)

$$A = (n-1)s^2 + n(\mu - \bar{y})^2 \quad \text{and} \quad z = \frac{A}{2\sigma^2}$$

$$dz = \left(-\frac{A}{2(\sigma^2)^2}\right) d\sigma^2$$

$$p(\mu \mid y) \propto A^{-n/2} \int_0^\infty z^{(n-2)/2} \exp(-z) dz$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal posterior $p(\mu | y)$

$$\begin{aligned} p(\mu | y) &= \int_0^\infty p(\mu, \sigma^2 | y) d\sigma^2 \\ &\propto \int_0^\infty \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\sigma^2 \end{aligned}$$

Transformation (integration by substitution)

$$A = (n-1)s^2 + n(\mu - \bar{y})^2 \quad \text{and} \quad z = \frac{A}{2\sigma^2}$$

$$dz = \left(-\frac{A}{2(\sigma^2)^2}\right) d\sigma^2$$

$$p(\mu | y) \propto A^{-n/2} \int_0^\infty z^{(n-2)/2} \exp(-z) dz$$

Recognize gamma integral $\Gamma(u) = \int_0^\infty x^{u-1} \exp(-x) dx$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu \mid y) &= \int_0^\infty p(\mu, \sigma^2 \mid y) d\sigma^2 \\ &\propto \int_0^\infty \sigma^{-n-2} \exp\left(-\frac{1}{2\sigma^2} [(n-1)s^2 + n(\bar{y} - \mu)^2]\right) d\sigma^2 \end{aligned}$$

Transformation (integration by substitution)

$$A = (n-1)s^2 + n(\mu - \bar{y})^2 \quad \text{and} \quad z = \frac{A}{2\sigma^2}$$

$$dz = \left(-\frac{A}{2(\sigma^2)^2}\right) d\sigma^2$$

$$p(\mu \mid y) \propto A^{-n/2} \int_0^\infty z^{(n-2)/2} \exp(-z) dz$$

Recognize gamma integral $\Gamma(u) = \int_0^\infty x^{u-1} \exp(-x) dx$

$$\propto [(n-1)s^2 + n(\mu - \bar{y})^2]^{-n/2}$$



$$p(\mu|y) \propto [(n-1)s^2 + n(\mu - \bar{y})^2]^{-n/2}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu|y) &\propto [(n-1)s^2 + n(\mu - \bar{y})^2]^{-n/2} \\ &\propto \left[1 + \frac{n(\mu - \bar{y})^2}{(n-1)s^2} \right]^{-n/2} \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Marginal posterior $p(\mu \mid y)$

$$\begin{aligned} p(\mu|y) &\propto [(n-1)s^2 + n(\mu - \bar{y})^2]^{-n/2} \\ &\propto \left[1 + \frac{n(\mu - \bar{y})^2}{(n-1)s^2} \right]^{-n/2} \end{aligned}$$

$$p(x|\mu_*, \sigma_*, \nu_*) = \frac{\Gamma(\frac{\nu_*+1}{2})}{\sqrt{\pi\nu_*\sigma_*^2} \Gamma(\frac{\nu_*}{2})} \left(1 + \frac{1}{\nu_*} \frac{(x - \mu_*)^2}{\sigma_*^2} \right)^{-\frac{\nu_*+1}{2}}$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Marginal posterior $p(\mu \mid y)$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

$$p(\mu|y) \propto [(n-1)s^2 + n(\mu - \bar{y})^2]^{-n/2}$$

$$\propto \left[1 + \frac{n(\mu - \bar{y})^2}{(n-1)s^2} \right]^{-n/2}$$

$$p(x|\mu_*, \sigma_*, \nu_*) = \frac{\Gamma(\frac{\nu_*+1}{2})}{\sqrt{\pi\nu_*\sigma_*^2} \Gamma(\frac{\nu_*}{2})} \left(1 + \frac{1}{\nu_*} \frac{(x - \mu_*)^2}{\sigma_*^2} \right)^{-\frac{\nu_*+1}{2}}$$

$$p(\mu \mid y) = \underbrace{t_{n-1}(\mu \mid \bar{y}, s^2/n)}_{\text{Student's } t}$$




- Introduction
 - Multiple parameter models
 - Marginalization
 - **Gaussian Model**
 - Example
 - Conjugate prior
 - Bioassay example
 - Additional examples
 - Multinomial model
 - Multivariate Gaussian
- For a new observation $\tilde{y}$, which uncertainties should be included?
 - uncertainty about μ ?
 - noise in a new observation?
 - both?
 - what are the two types of uncertainties called?




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior



Marginal of sigma




Predictive distribution for new $\tilde{y}$

$$p(\tilde{y}|y) = \int p(\tilde{y}|\mu, \sigma) p(\mu, \sigma|y) d\mu d\sigma$$




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior



Marginal of sigma



Predictive distribution for new $\tilde{y}$

$$p(\tilde{y}|y) = \int p(\tilde{y}|\mu, \sigma) p(\mu, \sigma|y) d\mu d\sigma$$


$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma|y)$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior

60
-Exact contour plot — Cond. distribution of μ
Sample from joint post. — Sample from the marg.




Predictive distribution for new $\tilde{y}$

$$p(\tilde{y}|y) = \int p(\tilde{y}|\mu, \sigma) p(\mu, \sigma|y) d\mu d\sigma$$

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma|y)$$


$$\tilde{y}^{(s)} \sim p(\tilde{y}|\mu^{(s)}, \sigma^{(s)})$$

Marginal of sigma



Posterior predictive distribution

- Sample from the predictive distribution
- Predictive distribution given the posterior sample







- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior

60
-Exact contour plot — Cond. distribution of μ
Sample from joint post. — Sample from the marg.



Marginal of sigma



Predictive distribution for new $\tilde{y}$


$$p(\tilde{y}|y) = \int p(\tilde{y}|\mu, \sigma) p(\mu, \sigma|y) d\mu d\sigma$$

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma|y)$$

$$\tilde{y}^{(s)} \sim p(\tilde{y}|\mu^{(s)}, \sigma^{(s)})$$

Posterior predictive distribution

- Sample from the predictive distribution
- Predictive distribution given the posterior sample






- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior

60
-Exact contour plot — Cond. distribution of μ
Sample from joint post. — Sample from the marg.




Predictive distribution for new $\tilde{y}$

$$p(\tilde{y}|y) = \int p(\tilde{y}|\mu, \sigma) p(\mu, \sigma|y) d\mu d\sigma$$

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma|y)$$


$$\tilde{y}^{(s)} \sim p(\tilde{y}|\mu^{(s)}, \sigma^{(s)})$$

Marginal of sigma



Posterior predictive distribution

- Sample from the predictive distribution
- Predictive distribution given the posterior sample







- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Joint posterior

60
-Exact contour plot — Cond. distribution of μ
Sample from joint post. — Sample from the marg.



Marginal of sigma



Predictive distribution for new $\tilde{y}$


$$p(\tilde{y}|y) = \int p(\tilde{y}|\mu, \sigma) p(\mu, \sigma|y) d\mu d\sigma$$

$$\mu^{(s)}, \sigma^{(s)} \sim p(\mu, \sigma|y)$$

$$\tilde{y}^{(s)} \sim p(\tilde{y}|\mu^{(s)}, \sigma^{(s)})$$

Posterior predictive distribution

- Sample from the predictive distribution
- Predictive distribution given the posterior sample
- Exact predictive distribution





UPPSALA
UNIVERSITET

Gaussian - posterior predictive distribution

Posterior predictive distribution given known variance

$$p(\tilde{y} \mid \sigma^2, y) = \int p(\tilde{y} \mid \mu, \sigma^2) p(\mu \mid \sigma^2, y) d\mu$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior predictive distribution given known variance

$$\begin{aligned} p(\tilde{y} \mid \sigma^2, y) &= \int p(\tilde{y} \mid \mu, \sigma^2) p(\mu \mid \sigma^2, y) d\mu \\ &= \int \mathcal{N}(\tilde{y} \mid \mu, \sigma^2) \mathcal{N}(\mu \mid \bar{y}, \sigma^2/n) d\mu \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior predictive distribution given known variance

$$\begin{aligned} p(\tilde{y} \mid \sigma^2, y) &= \int p(\tilde{y} \mid \mu, \sigma^2) p(\mu \mid \sigma^2, y) d\mu \\ &= \int \mathcal{N}(\tilde{y} \mid \mu, \sigma^2) \mathcal{N}(\mu \mid \bar{y}, \sigma^2/n) d\mu \\ &= \mathcal{N}(\tilde{y} \mid \bar{y}, (1 + \frac{1}{n})\sigma^2) \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior predictive distribution given known variance

$$\begin{aligned} p(\tilde{y} \mid \sigma^2, y) &= \int p(\tilde{y} \mid \mu, \sigma^2) p(\mu \mid \sigma^2, y) d\mu \\ &= \int \mathcal{N}(\tilde{y} \mid \mu, \sigma^2) \mathcal{N}(\mu \mid \bar{y}, \sigma^2/n) d\mu \\ &= \mathcal{N}(\tilde{y} \mid \bar{y}, (1 + \frac{1}{n})\sigma^2) \end{aligned}$$

this is up to scaling factor same as $p(\mu \mid \sigma^2, y)$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior predictive distribution given known variance

$$\begin{aligned} p(\tilde{y} \mid \sigma^2, y) &= \int p(\tilde{y} \mid \mu, \sigma^2) p(\mu \mid \sigma^2, y) d\mu \\ &= \int \mathcal{N}(\tilde{y} \mid \mu, \sigma^2) \mathcal{N}(\mu \mid \bar{y}, \sigma^2/n) d\mu \\ &= \mathcal{N}(\tilde{y} \mid \bar{y}, (1 + \frac{1}{n})\sigma^2) \end{aligned}$$

this is up to scaling factor same as $p(\mu \mid \sigma^2, y)$

$$p(\tilde{y} \mid y) = t_{n-1}(\tilde{y} \mid \bar{y}, (1 + \frac{1}{n})s^2)$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



With $p(\mu, \sigma^2) \propto \sigma^{-2}$ and $s^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$:

$$\sigma^2 \mid y \sim \text{Inv-}\chi^2(n-1, s^2)$$

$$\mu \mid y \sim t_{n-1}(\cdot \mid \bar{y}, s^2/n)$$

$$\tilde{y} \mid y \sim t_{n-1}(\cdot \mid \bar{y}, (1 + \frac{1}{n})s^2)$$

- Posterior uncertainty for μ is smaller than predictive uncertainty for a new $\tilde{y}$.



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- In these slides, $t_\nu(\cdot \mid m, \nu)$ uses ν as scale squared.
- If software expects a scale parameter, use $\sqrt{\nu}$.
- Examples:

$$\mu \mid y \sim t_{n-1}(\cdot \mid \bar{y}, s^2/n)$$

$$\tilde{y} \mid y \sim t_{n-1}(\cdot \mid \bar{y}, (1 + \frac{1}{n})s^2)$$



- Introduction
- Multiple parameter models
 - Marginalization
- **Gaussian Model**
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Often the quantity of interest is $q = g(\theta)$.
- If $\theta^{(s)} \sim p(\theta | y)$, then compute

$$q^{(s)} = g(\theta^{(s)}).$$

- The draws $q^{(s)}$ summarize $p(q | y)$.
- Examples: $\theta_1 - \theta_2$, $\mu_1 - \mu_2$, $x_{LD50} = -\alpha/\beta$.




- Introduction
 - Multiple parameter models
 - Marginalization
 - **Gaussian Model**
 - Example
 - Conjugate prior
 - Bioassay example
 - Additional examples
 - Multinomial model
 - Multivariate Gaussian
- Suppose $\mu_d = \mu_1 - \mu_2$ has a continuous posterior.
 - What is $\Pr(\mu_d = 0 \mid y)$?
 - What would be a more useful posterior summary?



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Simon Newcomb's speed of light experiment in 1882

Newcomb measured ($n = 66$) the time required for light to travel from his laboratory on the Potomac River to a mirror at the base of the Washington Monument and back, a total distance of 7442 meters.






- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Simon Newcomb's speed of light experiment in 1882


Newcomb measured ($n = 66$) the time required for light to travel from his laboratory on the Potomac River to a mirror at the base of the Washington Monument and back, a total distance of 7442 meters.





Simon Newcomb's speed of light experiment in 1882


Newcomb measured ($n = 66$) the time required for light to travel from his laboratory on the Potomac River to a mirror at the base of the Washington Monument and back, a total distance of 7442 meters.





Simon Newcomb's speed of light experiment in 1882

Newcomb measured ($n = 66$) the time required for light to travel from his laboratory on the Potomac River to a mirror at the base of the Washington Monument and back, a total distance of 7442 meters.





- Conjugate prior on the form $p(\sigma^2)p(\mu \mid \sigma^2)$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Conjugate prior on the form $p(\sigma^2)p(\mu \mid \sigma^2)$
- Handy parameterization

$$\mu \mid \sigma^2 \sim N(\mu_0, \sigma^2 / \kappa_0)$$

$$\sigma^2 \sim \text{Inv-}\chi^2(\nu_0, \sigma_0^2)$$

which can be written as

$$p(\mu, \sigma^2) = N\text{Inv-}\chi^2(\mu_0, \sigma_0^2 / \kappa_0; \nu_0, \sigma_0^2)$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - **Conjugate prior**
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Conjugate prior on the form $p(\sigma^2)p(\mu \mid \sigma^2)$
- Handy parameterization

$$\mu \mid \sigma^2 \sim N(\mu_0, \sigma^2/\kappa_0)$$

$$\sigma^2 \sim \text{Inv-}\chi^2(\nu_0, \sigma_0^2)$$

which can be written as

$$p(\mu, \sigma^2) = N\text{Inv-}\chi^2(\mu_0, \sigma_0^2/\kappa_0; \nu_0, \sigma_0^2)$$

- μ and σ^2 are a priori dependent
 - if σ^2 is large, then μ has wide prior



Joint posterior (exercise 3.9)

$$p(\mu, \sigma^2 | y) = \text{N} \text{H} \nu - \chi^2(\mu_n, \sigma_n^2 / \kappa_n; \nu_n, \sigma_n^2)$$

where

$$\mu_n = \frac{\kappa_0}{\kappa_0 + n} \mu_0 + \frac{n}{\kappa_0 + n} \bar{y}$$

$$\kappa_n = \kappa_0 + n$$

$$\nu_n = \nu_0 + n$$

$$\nu_n \sigma_n^2 = \nu_0 \sigma_0^2 + (n - 1) s^2 + \frac{\kappa_0 n}{\kappa_0 + n} (\bar{y} - \mu_0)^2$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Conditional $p(\mu \mid \sigma^2, y)$

$$\begin{aligned}\mu \mid \sigma^2, y &\sim N(\mu_n, \sigma^2 / \kappa_n) \\ &= N\left(\frac{\frac{\kappa_0}{\sigma^2} \mu_0 + \frac{n}{\sigma^2} \bar{y}}{\frac{\kappa_0}{\sigma^2} + \frac{n}{\sigma^2}}, \frac{1}{\frac{\kappa_0}{\sigma^2} + \frac{n}{\sigma^2}}\right)\end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Conditional $p(\mu \mid \sigma^2, y)$

$$\begin{aligned}\mu \mid \sigma^2, y &\sim N(\mu_n, \sigma^2 / \kappa_n) \\ &= N\left(\frac{\frac{\kappa_0}{\sigma^2} \mu_0 + \frac{n}{\sigma^2} \bar{y}}{\frac{\kappa_0}{\sigma^2} + \frac{n}{\sigma^2}}, \frac{1}{\frac{\kappa_0}{\sigma^2} + \frac{n}{\sigma^2}}\right)\end{aligned}$$

- Marginal $p(\sigma^2 \mid y)$

$$\sigma^2 \mid y \sim \text{Inv-}\chi^2(\nu_n, \sigma_n^2)$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Conditional $p(\mu \mid \sigma^2, y)$

$$\begin{aligned}\mu \mid \sigma^2, y &\sim N(\mu_n, \sigma^2 / \kappa_n) \\ &= N\left(\frac{\frac{\kappa_0}{\sigma^2} \mu_0 + \frac{n}{\sigma^2} \bar{y}}{\frac{\kappa_0}{\sigma^2} + \frac{n}{\sigma^2}}, \frac{1}{\frac{\kappa_0}{\sigma^2} + \frac{n}{\sigma^2}}\right)\end{aligned}$$

- Marginal $p(\sigma^2 \mid y)$

$$\sigma^2 \mid y \sim \text{Inv-}\chi^2(\nu_n, \sigma_n^2)$$


- Marginal $p(\mu \mid y)$

$$\mu \mid y \sim t_{\nu_n}(\mu \mid \mu_n, \sigma_n^2 / \kappa_n)$$



Bioassay

Dose, x_i (log g/ml)	Number of animals, n_i	Number of deaths, y_i
-0.86	5	0
-0.30	5	1
-0.05	5	3
0.73	5	5




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay

Dose, x_i (log g/ml)	Number of animals, n_i	Number of deaths, y_i
-0.86	5	0
-0.30	5	1
-0.05	5	3
0.73	5	5



Find out lethal dose 50% (LD50)


- used to classify how hazardous chemical is
- 1984 EEC directive has 4 levels



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay

Dose, x_i (log g/ml)	Number of animals, n_i	Number of deaths, y_i
-0.86	5	0
-0.30	5	1
-0.05	5	3
0.73	5	5



Find out lethal dose 50% (LD50)


- used to classify how hazardous chemical is
- 1984 EEC directive has 4 levels

Bayesian methods help to

- reduce the number of animals needed
- easy to make sequential experiment and stop as soon as desired accuracy is obtained




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian






- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian






- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian





- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian


Bioassay





- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay




Binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay





Binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i), \quad \text{logit}(\theta_i) = \log\left(\frac{\theta_i}{1 - \theta_i}\right) = \alpha + \beta x_i$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$
$$\text{logit}(\theta_i) = \log\left(\frac{\theta_i}{1 - \theta_i}\right)$$
$$= \alpha + \beta x_i$$






- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian


$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$
$$\text{logit}(\theta_i) = \log\left(\frac{\theta_i}{1 - \theta_i}\right)$$
$$= \alpha + \beta x_i$$

$$\theta_i = \frac{1}{1 + \exp(-(\alpha + \beta x_i))}$$

Logistic regression fit



Logistic regression in latent space





- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian


- What assumptions are needed for $y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$?
- Are animals within a dose group exchangeable and independent?
- Name one practical reason why the binomial model could fail.
- Why use a logit model instead of $\theta_i = \alpha + \beta x_i$?



- Introduction
 - Multiple parameter models
 - Marginalization
 - Gaussian Model
 - Example
 - Conjugate prior
 - Bioassay example
 - Additional examples
 - Multinomial model
 - Multivariate Gaussian
- In this lecture, we constrain toxicity to be monotone.
 - The grid is restricted to $\beta \geq 0$; increasing dose cannot decrease the probability of death.
 - Conceptually, the LD50 summary is interpreted under the positive-dose-effect assumption $\beta > 0$.
 - If $\beta \leq 0$ were allowed, we would also report $\Pr(\beta > 0 \mid y)$ and LD50 conditional on $\beta > 0$.




Logistic regression fit



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian





- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay: Lethal Dose 50%




$$E\left(\frac{y}{n}\right) = \text{logit}^{-1}(\alpha + \beta x) = 0.5$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay: Lethal Dose 50%




$$E\left(\frac{y}{n}\right) = \text{logit}^{-1}(\alpha + \beta x) = 0.5 \quad \Rightarrow \quad x_{\text{LD50}} = -\alpha/\beta$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay: Lethal Dose 50%




$$E\left(\frac{y}{n}\right) = \text{logit}^{-1}(\alpha + \beta x) = 0.5 \quad \Rightarrow \quad x_{\text{LD50}} = -\alpha/\beta$$
$$x_{\text{LD50}}^{(s)} = -\alpha^{(s)}/\beta^{(s)}$$



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay: Lethal Dose 50%



$$E\left(\frac{y}{n}\right) = \text{logit}^{-1}(\alpha + \beta x) = 0.5 \quad \Rightarrow \quad x_{\text{LD50}} = -\alpha/\beta$$
$$x_{\text{LD50}}^{(s)} = -\alpha^{(s)}/\beta^{(s)}$$



Bioassay posterior

Binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$

Link function

$$\text{logit}(\theta_i) = \alpha + \beta x_i$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay posterior

Binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$

Link function

$$\text{logit}(\theta_i) = \alpha + \beta x_i$$

Likelihood

$$p(y_i \mid \alpha, \beta, n_i, x_i) \propto \theta_i^{y_i} [1 - \theta_i]^{n_i - y_i}$$



Bioassay posterior

Binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$

Link function

$$\text{logit}(\theta_i) = \alpha + \beta x_i$$

Likelihood

$$\begin{aligned} p(y_i \mid \alpha, \beta, n_i, x_i) &\propto \theta_i^{y_i} [1 - \theta_i]^{n_i - y_i} \\ &\propto [\text{logit}^{-1}(\alpha + \beta x_i)]^{y_i} [1 - \text{logit}^{-1}(\alpha + \beta x_i)]^{n_i - y_i} \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay posterior

Binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$

Link function

$$\text{logit}(\theta_i) = \alpha + \beta x_i$$

Likelihood

$$\begin{aligned} p(y_i \mid \alpha, \beta, n_i, x_i) &\propto \theta_i^{y_i} [1 - \theta_i]^{n_i - y_i} \\ &\propto [\text{logit}^{-1}(\alpha + \beta x_i)]^{y_i} [1 - \text{logit}^{-1}(\alpha + \beta x_i)]^{n_i - y_i} \end{aligned}$$

Posterior

$$p(\alpha, \beta \mid y, n, x) \propto p(\alpha, \beta) \prod_{i=1}^k p(y_i \mid \alpha, \beta, n_i, x_i)$$

No analytic posterior distribution? What can we do?



1. Setup an area (can be hard) for α and β that capture most mass (here $\alpha = [-1, 5]$ and $\beta = [0, 30]$)

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Grid evaluation

1. Setup an area (can be hard) for α and β that capture most mass (here $\alpha = [-1, 5]$ and $\beta = [0, 30]$)
2. Compute unnormalized $p(\alpha^{(g)}, \beta^{(g)} \mid y, n, x)$, here $\tilde{p}$, at the grid points g



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Grid evaluation

1. Setup an area (can be hard) for α and β that capture most mass (here $\alpha = [-1, 5]$ and $\beta = [0, 30]$)
2. Compute unnormalized $p(\alpha^{(g)}, \beta^{(g)} \mid y, n, x)$, here $\tilde{p}$, at the grid points g
3. Sum up $\tilde{p}$ over the whole grid (for all $g \in \{1, \dots, G\}$)



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian


Grid evaluation

1. Setup an area (can be hard) for α and β that capture most mass (here $\alpha = [-1, 5]$ and $\beta = [0, 30]$)
2. Compute unnormalized $p(\alpha^{(g)}, \beta^{(g)} \mid y, n, x)$, here $\tilde{p}$, at the grid points g
3. Sum up $\tilde{p}$ over the whole grid (for all $g \in \{1, \dots, G\}$)
4. Compute (normalize) the pmf approximation of the posterior $\hat{p}$

g	(α, β)	$\tilde{p}$	$\hat{p}$
1	(0, -1)	0.02	0.0002
2	(0, -0.8)	0.03	0.0003
...	...	...	...
G	(30, 5)	0.001	0.00001
$\sum_g^G$	-	100	1



Posterior density evaluated in a grid




- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay (with uniform prior on α, β)




Density evaluated in grid, but plotted using interpolation



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay (with uniform prior on α, β)




Density evaluated in grid, and plotted without interpolation



Bioassay (with uniform prior on α, β)

Posterior density evaluated in a grid




Density evaluated in a coarser grid

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay (with uniform prior on α, β)




- Approximate the density as piecewise constant function
- Evaluate density in a grid over some finite region
- Density times cell area gives probability mass in each cell



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay (with uniform prior on α, β)




- Densities at 1, 2, and 3: 0.0027 0.0010 0.0001
- Probabilities of cells 1, 2, and 3: 0.0431 0.0166 0.0010
- Probabilities of cells sum to 1



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay (with uniform prior on α, β)

Posterior density and draws in a grid




- Sample according to grid cell probabilities



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- **Bioassay example**
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

Bioassay (with uniform prior on α, β)

Posterior density and draws in a grid




- Sample according to grid cell probabilities



Bioassay (with uniform prior on α, β)

Posterior density and draws in a grid




- Sample according to grid cell probabilities
- Several draws can be from the same grid cell



Bioassay (with uniform prior on α, β)

Posterior density in a grid and jittered draws



- Jitter can be added to improve visualization



- Draws can be used to estimate expectations, for example

$$E[x_{LD50}] = E[-\alpha/\beta] \approx \frac{1}{S} \sum_{s=1}^S \frac{-\alpha^{(s)}}{\beta^{(s)}}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Draws can be used to estimate expectations, for example

$$E[x_{LD50}] = E[-\alpha/\beta] \approx \frac{1}{S} \sum_{s=1}^S \frac{-\alpha^{(s)}}{\beta^{(s)}}$$

- Instead of sampling, grid could be used to evaluate functions directly, for example

$$E[-\alpha/\beta] \approx \sum_{t=1}^T w_{\text{cell}}^{(t)} \frac{-\alpha^{(t)}}{\beta^{(t)}},$$

where $w_{\text{cell}}^{(t)}$ is the normalized probability of a grid cell t , and $\alpha^{(t)}$ and $\beta^{(t)}$ are center locations of grid cells

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Draws can be used to estimate expectations, for example

$$E[x_{LD50}] = E[-\alpha/\beta] \approx \frac{1}{S} \sum_{s=1}^S \frac{-\alpha^{(s)}}{\beta^{(s)}}$$

- Instead of sampling, grid could be used to evaluate functions directly, for example

$$E[-\alpha/\beta] \approx \sum_{t=1}^T w_{\text{cell}}^{(t)} \frac{-\alpha^{(t)}}{\beta^{(t)}},$$

where $w_{\text{cell}}^{(t)}$ is the normalized probability of a grid cell t , and $\alpha^{(t)}$ and $\beta^{(t)}$ are center locations of grid cells

- Grid sampling gets computationally too expensive in high dimensions

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Multiparameter posteriors are joint distributions.
- Marginalization removes nuisance parameters.
- Predictive distributions include parameter uncertainty and new observation noise.
- Posterior draws make derived quantities easy to summarize.
- Bioassay: a nonconjugate logistic-binomial model, where grid evaluation works because the parameter dimension is small.



Multinomial model for categorical data

- Extension of binomial to K categories
- Observation model: **Multinomial model**

$$\mathbf{y} \mid \boldsymbol{\theta}, m \sim \text{Multinomial}(m, \boldsymbol{\theta})$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - **Multinomial model**
 - Multivariate Gaussian



- Extension of binomial to K categories
- Observation model: **Multinomial model**

$$\mathbf{y} \mid \boldsymbol{\theta}, m \sim \text{Multinomial}(m, \boldsymbol{\theta})$$

$$p(\mathbf{y} \mid \boldsymbol{\theta}, m) \propto \prod_{k=1}^K \theta_k^{y_k},$$

$$\text{where } \sum_{k=1}^K \theta_k = 1, \text{ and } \theta_k > 0$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - **Multinomial model**
 - Multivariate Gaussian



Multinomial model for categorical data

- Extension of binomial to K categories
- Observation model: **Multinomial model**

$$\mathbf{y} \mid \boldsymbol{\theta}, m \sim \text{Multinomial}(m, \boldsymbol{\theta})$$

$$p(\mathbf{y} \mid \boldsymbol{\theta}, m) \propto \prod_{k=1}^K \theta_k^{y_k},$$

$$\text{where } \sum_{k=1}^K \theta_k = 1, \text{ and } \theta_k > 0$$

- Example: $\mathbf{y} = (0, 1, 0, 6, 30)$ - **what is K here?**

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - **Multinomial model**
 - Multivariate Gaussian



Multinomial model for categorical data

- Extension of binomial to K categories
- Observation model: **Multinomial model**

$$\mathbf{y} \mid \boldsymbol{\theta}, m \sim \text{Multinomial}(m, \boldsymbol{\theta})$$

$$p(\mathbf{y} \mid \boldsymbol{\theta}, m) \propto \prod_{k=1}^K \theta_k^{y_k},$$

$$\text{where } \sum_{k=1}^K \theta_k = 1, \text{ and } \theta_k > 0$$

- Example: $\mathbf{y} = (0, 1, 0, 6, 30)$ - **what is K here?**
- Categorical distribution: $\text{Multinomial}(1, \boldsymbol{\theta})$



UPPSALA
UNIVERSITET

Prior for the multinomial model

- What is important when choosing the prior for θ ?

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- What is important when choosing the prior for θ ?
- Conjugate prior: The Dirichlet distribution

$$p(\theta) \propto \prod_{k=1}^K \theta_k^{\alpha_k - 1},$$

$$\text{where } \alpha_k > 0, \sum_{k=1}^K \theta_k = 1, \text{ and } \theta_k > 0$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Prior for the multinomial model

- What is important when choosing the prior for θ ?
- Conjugate prior: The Dirichlet distribution

$$p(\theta) \propto \prod_{k=1}^K \theta_k^{\alpha_k - 1},$$

$$\text{where } \alpha_k > 0, \sum_{k=1}^K \theta_k = 1, \text{ and } \theta_k > 0$$

- What is the connection to the Beta distribution?

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior for the multinomial model

Let $y_k = \sum_{i=1}^n y_{k,i}$ be the observed count in category k .

$$p(\theta \mid \mathbf{y}) \propto p(\mathbf{y} \mid \theta)p(\theta)$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior for the multinomial model

Let $y_k = \sum_{i=1}^n y_{k,i}$ be the observed count in category k .

$$p(\boldsymbol{\theta} \mid \mathbf{y}) \propto p(\mathbf{y} \mid \boldsymbol{\theta})p(\boldsymbol{\theta})$$

$$\propto \prod_{k=1}^K \theta_k^{\alpha_k - 1} \prod_{i=1}^n \prod_{k=1}^K \theta_k^{y_{k,i}}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior for the multinomial model

Let $y_k = \sum_{i=1}^n y_{k,i}$ be the observed count in category k .

$$p(\boldsymbol{\theta} \mid \mathbf{y}) \propto p(\mathbf{y} \mid \boldsymbol{\theta})p(\boldsymbol{\theta})$$

$$\begin{aligned} &\propto \prod_{k=1}^K \theta_k^{\alpha_k - 1} \prod_{i=1}^n \prod_{k=1}^K \theta_k^{y_{k,i}} \\ &= \prod_{k=1}^K \theta_k^{\alpha_k - 1} \prod_{k=1}^K \theta_k^{\sum_{i=1}^n y_{k,i}} \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior for the multinomial model

Let $y_k = \sum_{i=1}^n y_{k,i}$ be the observed count in category k .

$$p(\theta \mid \mathbf{y}) \propto p(\mathbf{y} \mid \theta) p(\theta)$$

$$\begin{aligned} &\propto \prod_{k=1}^K \theta_k^{\alpha_k - 1} \prod_{i=1}^n \prod_{k=1}^K \theta_k^{y_{k,i}} \\ &= \prod_{k=1}^K \theta_k^{\alpha_k - 1} \prod_{k=1}^K \theta_k^{\sum_{i=1}^n y_{k,i}} \\ &= \prod_{k=1}^K \theta_k^{\alpha_k - 1 + \sum_{i=1}^n y_{k,i}} \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



Posterior for the multinomial model

Let $y_k = \sum_{i=1}^n y_{k,i}$ be the observed count in category k .

$$p(\boldsymbol{\theta} \mid \mathbf{y}) \propto p(\mathbf{y} \mid \boldsymbol{\theta})p(\boldsymbol{\theta})$$

$$\begin{aligned} &\propto \prod_{k=1}^K \theta_k^{\alpha_k-1} \prod_{i=1}^n \prod_{k=1}^K \theta_k^{y_{k,i}} \\ &= \prod_{k=1}^K \theta_k^{\alpha_k-1} \prod_{k=1}^K \theta_k^{\sum_{i=1}^n y_{k,i}} \\ &= \prod_{k=1}^K \theta_k^{\alpha_k-1+\sum_{i=1}^n y_{k,i}} \\ &= \text{Dirichlet}(\alpha_1 + y_1, \dots, \alpha_K + y_K) \end{aligned}$$

- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian



- Introduction
- Multiple parameter models
 - Marginalization
- Gaussian Model
 - Example
 - Conjugate prior
- Bioassay example
- Additional examples
 - Multinomial model
 - Multivariate Gaussian

- Observation model

$$p(y \mid \mu, \Sigma) \propto |\Sigma|^{-1/2} \exp \left(-\frac{1}{2} (y - \mu)^T \Sigma^{-1} (y - \mu) \right),$$

where $y \in \mathcal{R}^D$

- See BDA3 p. 72–
- New recommended LKJ-prior mentioned in Appendix A, see more in Stan manual