

Assignment-based Subjective Questions

Question 1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: <Your answer for Question 1 goes below this line> (Do not edit)

-After analysis of the categorical variables from the dataset we could inferred that the bike rental rates are likely to be getting higher in summer every year, and the most demanding month is September. No impact of weekday or weekend. Approx. 3200-6500 users are renting bikes . Additionally we saw that bike rentals are higher on holidays.

Question 2. Why is it important to use **drop_first=True** during dummy variable creation? (Do not edit)

Total Marks: 2 marks (Do not edit)

Answer: <Your answer for Question 2 goes below this line> (Do not edit)

-drop_first=True reduces the extra column to avoid redundancy created during the dummy variable creation.

Question 3. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable? (Do not edit)

Total Marks: 1 mark (Do not edit)

Answer: <Your answer for Question 3 goes below this line> (Do not edit)

-Temp Variable.

Question 4. How did you validate the assumptions of Linear Regression after building the model on the training set? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: <Your answer for Question 4 goes below this line> (Do not edit)

-Validated by checking the VIF, error distribution of residuals and linear relationship between the dependent variable and a feature variable.

Question 5. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes? (Do not edit)

Total Marks: 2 marks (Do not edit)

Answer: <Your answer for Question 5 goes below this line> (Do not edit)

-Year, September month, season Spring.

General Subjective Questions

Question 6. Explain the linear regression algorithm in detail. (Do not edit)

Total Marks: 4 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<

Linear Regression is an ML algorithm. It helps in predicting a dependent variable(target) based on the given independent variable(s). The regression technique tends to establish a linear relationship between a dependent variable and the other given independent variables. There are two types of linear regression- simple linear regression and multiple linear regression. Simple linear regression is used when a single independent variable is used to predict the value of the target variable. Multiple Linear Regression is when multiple independent variables are used to predict the numerical value of the target variable. A linear line showing the relationship between the dependent and independent variables is called a regression line.>

Question 7. Explain the Anscombe's quartet in detail. (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<It consists of four data sets that have nearly identical simple descriptive statistics but have very different distributions and appear very different when presented graphically. Each and every dataset consists of eleven points. The main purpose of Anscombe's quartet is to illustrate the importance by looking at a set of data graphically before starting the analysis process as the statistics does not give an accurate representation of two datasets being compared. >

Question 8. What is Pearson's R? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<It is used to establish a linear relationship between two quantities. It gives an indication of the measure of strength between two variables and the value of the coefficient can be between -1 and +1. >

Question 9. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Scaling is a technique performed in pre-processing during building a machine learning model to standardize the independent feature variables in the dataset in a fixed range. The dataset could have several features which are highly ranging between high magnitudes and units. If there is no scaling performed on this data, it leads to incorrect modelling as there will be some mismatch in the units of all the features involved in the model. The difference between normalization and standardization is that while normalization brings all the data points in a range between 0 and 1,

standardization replaces the values with their Z scores>

Question 10. You might have observed that sometimes the value of VIF is infinite. Why does this happen? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<The value of VIF is infinite when there is a perfect correlation between the two independent variables. The R squared value is 1 in this case. This leads to VIF infinity as VIF equals to $1/(1-R^2)$. This concept suggests that there is a problem of multi-collinearity and one of these variables need to be dropped in order to define a working model for regression. >

Question 11. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression. (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<The quantile-quantile (Q-Q) plot are used to plot quantiles of a sample distribution with a theoretical distribution to determine if any dataset concerned follows any distribution such as normal, uniform or exponential distribution. It helps us determine if two datasets follow the same kind of distribution. It also helps to find out if the errors in dataset are normal in nature or not. >
