



UNIVERSITÀ
DI TORINO

Characterization of hadronic showers in the Belle II Electromagnetic Calorimeter

Metatesi exam

Emanuele Zanusso

Dipartimento di Fisica
Università degli studi di Torino

January 14 2026

Summary




1. Anti-neutron in physics experiment

2. Cluster variables

Anti-neutron in HEP experiments

The \bar{n} plays a key role in several physics measurements, such as:

- The neutron e.m. form factor study in $e^+ + e^- \rightarrow n + \bar{n}$ annihilation [1]



- The hyperon decay channels which involved \bar{n} , such as:

$$\bar{\Lambda}^0 \rightarrow \pi^0 + \bar{n}, \quad \bar{\Sigma}^- \rightarrow \pi^- + \bar{n}, \quad \bar{\Lambda}_c \rightarrow K_s^0 + \pi^0 + \bar{n}$$

- Other typical B-factories decay channels which involved \bar{n} , such as:

$$e^+ + e^- \rightarrow p + \bar{n} + X^- \quad (X^-: \text{combination of charged pions and kaons})$$

Anti-neutron in astrophysical experiments

\bar{n} also plays a key role in several astro-physics measurements, such as:

- Studying \bar{n} - anti-hyperon potential to improve the understanding of the equation of state of the neutron star [2]
- Investigating dark matter through anti-deuterons (\bar{D}) in cosmic rays, produced by dark matter annihilation or decay [3]:

$$A_{d.m.} + B_{d.m.} \rightarrow \text{hadrons } (n, \bar{n}, p, \bar{p} \text{ etc...})$$

$$X_{d.m.} \rightarrow \text{hadrons } (n, \bar{n}, p, \bar{p} \text{ etc...})$$

- \bar{D} is mainly produced through a coalescence mechanism:


$$\bar{n} + \bar{p} \rightarrow \bar{D}$$

Where \bar{p} and \bar{n} are nearby in the phase-space

Anti-neutron interactions in physics

The \bar{n} interacts with matter primarily via strong nuclear force

- It can annihilate with nucleons in the material, producing light mesons (mainly pions)
- Hadronic (π^+ , π^-) and electromagnetic (π^0) showers are generated within the detectors
- Since annihilation stars are produced, both forward and backward directions are involved




Electromagnetic and hadronic showers

Different processes occur for e.m. (1) and hadronic (2) showers:

1. $\pi^0 \rightarrow \gamma\gamma$ and bremsstrahlung process (e^+ , e^- , γ)
2. Strong interactions of hadrons with the material (p , n , $pions$, $kaons$...)

After \bar{n} annihilation, the produced pions can be detected via their showers:

- About the 95% of the hadronic shower is contained within a cylinder of radius λ_{had} (~ 44.12 cm in Csl)
- About the 90% of the e.m. shower is contained within a cylinder of radius R_M (~ 3.6 cm in Csl)





Anti-neutron interactions in physics

Several channels can be selected to look at \bar{n} annihilation, such as:

- $e^+ + e^- \rightarrow J/\Psi \rightarrow p + \bar{n} + \pi^-$ (BESIII)
- $e^+ + e^- + \gamma_{ISR} \rightarrow X \rightarrow p + \bar{n} + \pi^-$ (Belle II)
- $\bar{\Lambda}_c \rightarrow K_S^0 + \pi^0 + \bar{n}$

Several variables can be used to distinguish their clusters, such as:

- clusterLAT, clusterSecondMoment, clusterZernikeMoment, clusterNHits...



The MANTRA project

Measuring **A**nti-**N**eutron: **T**agging and **R**econstruction **A**lgorithm:

- A general method to measure the $E_{\bar{n}}$ up to 10 GeV, by combining information from:
 1. A detector with high time resolution ($< 100ps$), like a T.O.F. detector
 2. An electromagnetic calorimeter
 3. A muon system (alternating layers of active material and high-Z absorber)
- These features are common in modern general-purpose collider experiments such as **Belle II** and BESIII
- For MANTRA project, only signals from (1) and/or (2) are taken into account. In this thesis only (2) signals are studied

The MANTRA project

The measurement of the energy of the anti-neutrons is a two-step process:



1. \bar{n} identification via its induced shower vs ones from by photons or other hadrons
2. Combine the signals from (1) and (2) to reconstruct the \bar{n} 's energy
 - If π^0 ($\sim 5\%$): energy is all contained in the calorimeter, the shower is fully reconstructed
 - If π^\pm ($\sim 95\%$): their products may escape the crystals
→ the goal is to complement the calorimeter information with that from the adjacent detectors

The Belle II experiment



SuperKEKB is an asymmetric $e^+ e^-$ collider
(Tsukuba, Japan)

- 7 GeV electrons beam (HER)
- 4 GeV positrons beam (LER)
- Peak Lumi $\sim 5.1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ (Dec 2024)





It operates mainly around $\Upsilon(4S)$ resonance ($\sim 10.58 \text{ GeV}$):

- It decays almost exclusively into entangled couple of $B \bar{B} \rightarrow B - \text{factory}$
- Several goals: flavour physics, BSM, B and charm mesons etc...

The Electromagnetic CaLorimeter

The ECL plays a central role in this thesis

- Array of **Csl(Tl)** crystals (8376 $6\times6\times30\text{cm}^3$ crystals in total)
- It covers barrel and end-caps regions ($12^\circ \leq \theta \leq 155^\circ$)
- Energy resolution of 4% @100 MeV and 1.6% @8 GeV



n vs gamma list in cluster variables





Thank you for your attention

Emanuele Zanusso

Dipartimento di Fisica
Università degli studi di Torino

January 14 2026

References I



M. Ablikim et al.
Nature Physics, 17, 2021.



G.F. Burgio et al.
Prog. Part. Nucl. Phys., 10389:120, 2021.



F. Donato, N. Fornengo, and P. Salati.
Phys. Rev. D, 62:043003, 2000.