



UNIVERSITÀ
DI TORINO

Characterization of hadronic showers in the Belle II Electromagnetic Calorimeter

Metatesi exam

Emanuele Zanusso

Dipartimento di Fisica
Università degli studi di Torino

January 14 2026

Talk Outline




1. Anti-neutron in physics experiment

2. Preliminary study via clean MC sample

Anti-neutron in HEP experiments

The \bar{n} plays a key role in several physics measurements, such as:

- The neutron e.m. form factor studies in $e^+ + e^- \rightarrow n + \bar{n}$ process [1]



- The decay channels studied at B-factories which involved \bar{n} , such as:
 - The hyperons decay channel:

$$\bar{\Lambda}^0 \rightarrow \pi^0 + \bar{n}, \quad \bar{\Sigma}^- \rightarrow \pi^- + \bar{n}, \quad \bar{\Lambda}_c \rightarrow K_s^0 + \pi^0 + \bar{n}$$

- Other typical B-factories processes:

$$e^+ + e^- \rightarrow p + \bar{n} + X^- \quad (X^-: \text{combination of charged pions and kaons})$$

Anti-neutron in astrophysical experiments

\bar{n} also plays a key role in several astro-physics measurements, such as:

- Studying \bar{n} - anti-hyperon potential to improve the understanding of the equation of state of the neutron star [2]
- Investigating dark matter through anti-deuterons (\bar{D}) in cosmic rays, produced by dark matter annihilation or decay [3]:

$$A_{d.m.} + B_{d.m.} \rightarrow \text{hadrons } (n, \bar{n}, p, \bar{p} \text{ etc...})$$

$$X_{d.m.} \rightarrow \text{hadrons } (n, \bar{n}, p, \bar{p} \text{ etc...})$$

- \bar{D} is mainly produced through a coalescence mechanism:

$$\bar{n} + \bar{p} \rightarrow \bar{D}$$



Where \bar{p} and \bar{n} are nearby in the phase-space

The Belle II experiment



SuperKEKB is an asymmetric $e^+ e^-$ collider (Tsukuba, Japan)

- 7 GeV electrons beam (HER)
- 4 GeV positrons beam (LER)
- Peak Lumi $\sim 5.1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ 🏆
- Design Lumi $\sim 6.5 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$
→ x40 the Belle's one





It operates mainly around $\Upsilon(4S)$ resonance ($\sim 10.58 \text{ GeV}$):

- It decays almost exclusively into entangled couple of $B \bar{B} \rightarrow$ B-factory
- Several goals: flavour physics, BSM, B and charm mesons etc...

The Electromagnetic Calorimeter

The ECL plays a central role in this thesis


- Array of **CsI(Tl)** crystals ($8376\ 6\times6\times30\text{cm}^3$ crystals in total)
- It covers barrel and end-caps regions ($12^\circ \leq \theta \leq 155^\circ$)
- Energy resolution of 4% @100 Mev and 1.6% @8 GeV



Anti-neutron interactions in physics

The \bar{n} interacts with matter primarily via strong nuclear force

- It can annihilate with nucleons in the material, producing light mesons (mainly pions)
- Hadronic (π^+ , π^-) and electromagnetic (π^0) showers are generated within the ECL
- Since annihilation stars are produced, both backward (TOP) and forward (KLM) directions are involved




Electromagnetic and hadronic showers

Different processes occur for e.m. (1) and hadronic (2) showers:

1. Brems. and p.p. process (e^+ , e^- , γ) and $\pi^0 \rightarrow \gamma\gamma$
2. Strong interactions of hadrons with the material (p , n , *pions*, *kaons*...)

After \bar{n} annihilation, the produced pions can be detected via their showers:

- About the 95% of the hadronic shower is contained within a cylinder of radius λ_{had} (~ 44.12 cm in CsI(Tl))
- About the 90% of the e.m. shower is contained within a cylinder of radius R_M (~ 3.6 cm in CsI(Tl))




Anti-neutron interactions in physics

Several channels can be selected to look at \bar{n} annihilation, such as:

- $e^+ + e^- \rightarrow X \rightarrow p + \bar{n} + \pi^-$ (Mine)
- $\bar{\Lambda}_c \rightarrow K_S^0 + \pi^0 + \bar{n}$
- $\Lambda(\rightarrow p + \pi^-) + \bar{\Lambda}(\rightarrow \bar{n} + \pi^0)$

Several variables can be used to distinguish their clusters, such as:

- clusterZernikeMoment, clusterSecondMoment, clusterLAT etc...



The distributions for ECL variables for \bar{p} and \bar{n} do not in agree [4] $\rightarrow \bar{p}$ cannot be used as proxy for \bar{n}

The MANTRA project

Measuring Anti-Neutron: Tagging and Reconstruction Algorithm:

- A general method to measure the $E_{\bar{n}}$ up to 10 GeV, by combining information from:
 1. A detector with high time resolution ($< 100ps$), like a T.O.F. detector (TOP)
 2. An electromagnetic calorimeter (**ECL**)
 3. A muon system (alternating layers of active material and high-Z absorber) (KLM)
- These features are common in modern general-purpose collider experiments such as **Belle II** and BESIII
- For MANTRA project, only signals from ECL and TOP are taken into account. In this thesis only ECL signals are studied

The MANTRA project

Anti-neutrons do not interact with tracking sub-detector. The measurement of the energy is a two-step process:

1. \bar{n} identification via its induced ECL clusters and how they correlate to the initial energy
2. Combine the signals from (1) and (2) to reconstruct the \bar{n} 's energy, in cases of backscatter or pre-annihilation
 - If π^0 ($\sim 5\%$): energy is all contained in the calorimeter, the shower is fully reconstructed
 - If π^\pm ($\sim 95\%$): their products may escape the crystals
→ the goal is to complement the calorimeter information with that from the adjacent detectors

Analysis outline

1. Preliminary study of a clean selected channel via generators
 - (a) Recoil identification from the three-body system $p + \gamma_{ISR} + \pi^-$
 - (b) Study of the kinematic recoil variables (momentum, angles, energy, etc...)
 - (c) Study of ECL clusters
 - (d) Study of the effect of 1C kinematic fit over the $p + \gamma_{ISR} + \pi^-$ recoil mass
2. Study of MC cocktail events sample:
 - (a) Recoil identification from the three-body system $p + \gamma_{ISR} + \pi^-$
3. Study of real data events sample:
 - (a) Recoil identification from the three-body system $p + \gamma_{ISR} + \pi^-$
 - (b) Constraint with 1C kinematic fit over the $p + \gamma_{ISR} + \pi^-$ recoil mass
 - (c) Examine Data/MC agreement in ECL cluster shapes from \bar{n} channel

Analysis outline

- The analyzed channel is:

$$e^+ + e^- + \gamma_{ISR} \rightarrow X \rightarrow p + \bar{n} + \pi^- \text{ (Phokhara+evt_gen generator)}$$

The reconstructed particles are (cuts and selections in backup):

- $vpho \rightarrow p + \gamma_{ISR} + \pi^-$, where $vpho$ is a fake particle, mimicking the recoil system
- \bar{n} candidate list, used to compare its variables with those of the recoil

rowNo	decay tree	decay final state	iDcyTr	nEtr	nCEtr
1	$vpho \rightarrow \pi^- \bar{n} p$	$\pi^- \bar{n} p$	0	35291	35291
2	$e^+ e^- \rightarrow vpho \gamma^I \gamma^I, vpho \rightarrow \pi^- \bar{n} p$	$\pi^- \bar{n} p \gamma^I \gamma^I$	2	22971	58262
3	$e^+ e^- \rightarrow vpho \gamma^I, vpho \rightarrow \pi^- \bar{n} p$	$\pi^- \bar{n} p \gamma^I$	1	18735	76997
4	$vpho \rightarrow \pi^- \bar{n} p \gamma^F$	$\pi^- \bar{n} p \gamma^F$	3	10005	87002
5	$e^+ e^- \rightarrow vpho \gamma^I \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^I \gamma^F$	6	5274	92276
6	$e^+ e^- \rightarrow vpho \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^F$	4	4621	96897
7	$vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^F \gamma^F$	7	1503	98400
8	$e^+ e^- \rightarrow vpho \gamma^I \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^I \gamma^F \gamma^F$	8	700	99100
9	$e^+ e^- \rightarrow vpho \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^F \gamma^F$	5	597	99697
10	$vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^F \gamma^F \gamma^F$	9	167	99864
11	$e^+ e^- \rightarrow vpho \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^F \gamma^F \gamma^F$	12	63	99927
12	$e^+ e^- \rightarrow vpho \gamma^I \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^I \gamma^F \gamma^F \gamma^F$	10	61	99988
13	$e^+ e^- \rightarrow vpho \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^F \gamma^F \gamma^F \gamma^F$	11	4	99992
14	$e^+ e^- \rightarrow vpho \gamma^I \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^I \gamma^F \gamma^F \gamma^F \gamma^F$	15	4	99996
15	$vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F$	14	2	99998
16	$vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F \gamma^F$	13	1	99999
17	$e^+ e^- \rightarrow vpho \gamma^I \gamma^I, vpho \rightarrow \pi^- \bar{n} p \gamma^F \gamma^F \gamma^F \gamma^F \gamma^F$	$\pi^- \bar{n} p \gamma^I \gamma^I \gamma^F \gamma^F \gamma^F \gamma^F \gamma^F$	16	1	100000


- 100k events** were generated, **17525** candidates have been reconstructed
→ reconstruction efficiency:

$$\epsilon = \frac{\text{n}^\circ \text{ of reconstructed candidates}}{\text{n}^\circ \text{ of generated events}} \sim 18\%$$

The recoil mass (1a)


- The recoil three body system is well reconstructed as shown in the recoil mass distribution, where a peak emerges above the \bar{n} mass.

→ reconstructed \bar{n} variables can be compared with the reconstructed recoil variables (p , θ , ϕ)




The recoil and the \bar{n} momentum (1b)

- The reconstructed \bar{n} candidate list shows a discrepancy with the recoil momentum $\rightarrow \gamma$'s are mis-identified as \bar{n} in reconstruction
- MC selection $\bar{n}_{mcPDG!} = 22$ is applied in order to directly compare the recoil kinematic variables with the \bar{n} from MC truth







The recoil and the \bar{n} momentum (1b)

- The reconstructed \bar{n} candidate list shows a discrepancy with the recoil momentum $\rightarrow \gamma$'s are mis-identified as \bar{n} in reconstruction
- MC selection $\bar{n}_{mcPDG!} = 22$ is applied in order to directly compare the recoil kinematic variables with the \bar{n} from MC truth



\bar{n} vs recoil vector correlation (1b)


- Good correlation is observed at the generator level in both the momentum and θ distributions
- The reconstructed \bar{n} momentum in the ECL is not a reliable variable, since no high correlation is observed (energy loss)



\bar{n} ECL cluster variables (1c)







Cluster variables can be studied to distinguish \bar{n} from other neutral particles:

- **clusterE**, **clusterE1E9** and **clusterE9E21**
($E_{min} = 20$ GeV)
- **clusterAbsZernikeMoment51**: $|Z_{51}|$




- **clusterLAT**: lateral energy distribution, defined as:
$$S = \frac{\sum_{i=2}^n \omega_i E_i r_i^2}{\omega_0 E_0 r_0^2 + \omega_1 E_1 r_1^2 + \sum_{i=2}^n \omega_i E_i r_i^2}$$
- **clusterSecondMoment**: second moment S , defined as:
$$S = \frac{\sum_{i=0}^n \omega_i E_i r_i^2}{\sum_{i=0}^n \omega_i E_i}$$

\bar{n} ECL cluster variables (1c)




\bar{n} ECL cluster variables (1c)



Analysis outline

Analysis of a $\Lambda \rightarrow p + \pi^-$ ($\bar{\Lambda} \rightarrow \bar{p} + \pi^+$) sample shows that [5]:



Poor Data/MC agreement in $\bar{p} \rightarrow$ will it be the same for \bar{n} ?



Thank you for your attention

Emanuele Zanusso


Dipartimento di Fisica
Università degli studi di Torino

January 14 2026

Applied selections and cuts on clean MC sample




- (a) $protonID > 0.9$ and $dr < 1$ and $abs(dz) < 3$ and $pionID > 0.1$ (From IP?)
- (b) $p_PDG == 2212$ and $pi_PDG == -211$ and $gamma_PDG == 22$
- (c) Rec. \bar{n} in theta ECL Acceptance and From ECL
- (d) $mRecoil > 0GeV$ and $mRecoil < 2GeV$
- (e) $\alpha < 0.35$ rad (~ 20 deg), where α is the 3D angle between the recoil vector and the closest reconstructed \bar{n} candidate (rankByLowest)




\bar{n} mcPDG I

γ 's are mis-identified as \bar{n} in reconstruction:



$p_{\bar{n}}/p_{\text{Recoil}}$ I

\bar{n} is underrated in the most of cases (annihilation process + loss of energy)



References I

- [1] M. Ablikim et al. In: *Nature Physics* 17 (2021).
- [2] G.F. Burgio et al. In: *Prog. Part. Nucl. Phys.* 10389 (2021), p. 120.
- [3] F. Donato, N. Fornengo, and P. Salati. In: *Phys. Rev. D* 62 (2000), p. 043003.
- [4] Sanjeeda Das. In: *Belle II Italy May* (2025).
- [5] Shanette De La Motte. In: *Belle II Italy December* (2025).