CONCORDIA UNIVERSITY DEPARTMENT OF COMPUTER SCIENCE AND SOFTWARE ENGINEERING SOEN 6011: SOFTWARE ENGINEERING PROCESS: SUMMER 2019

DELIVERABLE 1: OPEN PROBLEMS

STUDENT NAME: MANUSHREE MALLARAJU STUDENT IDENTIFICATION NUMBER: 40082236

PROBLEM T1-OP5. Source(s): "Give a brief description, not exceeding one page, of your function, including the domain and co-domain of function, and the characteristics that make it unique. To ensure that you have attained sufficient background, Test 1 will have a problem related to your function."

INTRODUCTION

The exponential function ab^x is one of the power rules of math. As the name of an exponential function is described, it involves an exponent. This exponent is represented with a variable rather than a constant, and its base is represented with constant value rather than a variable. Let $(f(x) = ab^x)$ be an exponential function where "b" is its change factor (or a constant), the exponent "x" is the independent variable (or input of the function), the coefficient "a" is called the initial value of the function (or the y-intercept), and "f(x)" represent the dependent variable (or output of the function)

DOMAIN:

The domain is a set of all real numbers, R. where :b>0, x>0

CO DOMAIN

The co-domain is also a set of all real numbers, R.

CHARACTERISTICS

- 1. Exponential functions defined by an equation of the form ab^x are called exponential decay functions if the change factor "b" (fixed base value) is 0 < b < 1, or it is also called exponential growth functions if the change factor is b > 0
- 2. the y-intercept is (0,a) and it is located at the initial value "a". There is no x-intercept. The domain for an exponential decay function of this form is all real numbers and the range is y > 0

GRAPH

Representation of ab^x



