

Slides from FYS4480/9480 Lectures

Morten Hjorth-Jensen

Department of Physics and Center for Computing in Science Education
University of Oslo, N-0316 Oslo, Norway and
Department of Physics and Astronomy and Facility for Rare Isotope Beams,
Michigan State University, East Lansing, MI 48824, USA

Fall 2022

Diagram rules

- ▶ Draw all topologically distinct diagrams by linking up particle and hole lines with various interaction vertices. Two diagrams can be made topologically equivalent by deformation of fermion lines under the restriction that the ordering of the vertices is not changed and particle lines and hole lines remain particle and hole lines.
- ▶ For the explicit evaluation of a diagram: Sum freely over all internal indices and label all lines.
- ▶ Extract matrix elements for the one-body operators (if present) as $\langle \text{out} | \hat{f} | \text{in} \rangle$ and for the two-body operator (if present) as $\langle \text{left out}, \text{right out} | \hat{V} | \text{left in}, \text{right in} \rangle$.

Diagram rules

- ▶ Calculate the phase factor: $(-1)^{\text{holelines}+\text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent pair of lines (particle lines or hole lines) that begin at the same interaction vertex and end at the same (yet different from the first) interaction vertex.
- ▶ For each interval between successive interaction vertices with minimum one single-particle state above the Fermi level with n hole states and m particle states there is a factor

$$\frac{1}{\sum_i^n \epsilon_i - \sum_a^m \epsilon_a}.$$

Schrödinger picture

The time-dependent Schrödinger equation (or equation of motion) reads

$$i\hbar \frac{\partial}{\partial t} |\Psi_S(t)\rangle = \hat{H} \Psi_S(t)\rangle,$$

where the subscript S stands for Schrödinger here. A formal solution is given by

$$|\Psi_S(t)\rangle = \exp(-i\hat{H}(t - t_0)/\hbar) |\Psi_S(t_0)\rangle.$$

The Hamiltonian \hat{H} is hermitian and the exponent represents a unitary operator with an operation carried out on the wave function at a time t_0 .

Interaction picture

Our Hamiltonian is normally written out as the sum of an unperturbed part \hat{H}_0 and an interaction part \hat{H}_I , that is

$$\hat{H} = \hat{H}_0 + \hat{H}_I.$$

In general we have $[\hat{H}_0, \hat{H}_I] \neq 0$ since $[\hat{T}, \hat{V}] \neq 0$. We wish now to define a unitary transformation in terms of \hat{H}_0 by defining

$$|\Psi_I(t)\rangle = \exp(i\hat{H}_0 t/\hbar)|\Psi_S(t)\rangle,$$

which is again a unitary transformation carried out now at the time t on the wave function in the Schrödinger picture.

Interaction picture

We can easily find the equation of motion by taking the time derivative

$$i\hbar \frac{\partial}{\partial t} |\Psi_I(t)\rangle = -\hat{H}_0 \exp(i\hat{H}_0 t/\hbar) \Psi_S(t) + \exp(i\hat{H}_0 t/\hbar) i\hbar \frac{\partial}{\partial t} \Psi_S(t).$$

Interaction picture

Using the definition of the Schrödinger equation, we can rewrite the last equation as

$$i\hbar \frac{\partial}{\partial t} |\Psi_I(t)\rangle = \exp(i\hat{H}_0 t/\hbar) \left[-\hat{H}_0 + \hat{H}_0 + \hat{H}_I \right] \exp(-i\hat{H}_0 t/\hbar) \Psi_I(t),$$

which gives us

$$i\hbar \frac{\partial}{\partial t} |\Psi_I(t)\rangle = \hat{H}_I(t) \Psi_I(t),$$

with

$$\hat{H}_I(t) = \exp(i\hat{H}_0 t/\hbar) \hat{H}_I \exp(-i\hat{H}_0 t/\hbar).$$

Interaction picture

The order of the operators is important since \hat{H}_0 and \hat{H}_I do generally not commute. The expectation value of an arbitrary operator in the interaction picture can now be written as

$$\langle \Psi'_S(t) | \hat{O}_S | \Psi_S(t) \rangle = \langle \Psi'_I(t) | \exp(i\hat{H}_0 t/\hbar) \hat{O}_I \exp(-i\hat{H}_0 t/\hbar) | \Psi_I(t) \rangle,$$

and using the definition

$$\hat{O}_I(t) = \exp(i\hat{H}_0 t/\hbar) \hat{O}_I \exp(-i\hat{H}_0 t/\hbar),$$

we obtain

$$\langle \Psi'_S(t) | \hat{O}_S | \Psi_S(t) \rangle = \langle \Psi'_I(t) | \hat{O}_I(t) | \Psi_I(t) \rangle,$$

stating that a unitary transformation does not change expectation values!

Interaction picture

If we take the time derivative of the operator in the interaction picture we arrive at the following equation of motion

$$i\hbar \frac{\partial}{\partial t} \hat{O}_I(t) = \exp(i\hat{H}_0 t/\hbar) \left[\hat{O}_S \hat{H}_0 - \hat{H}_0 \hat{O}_S \right] \exp(-i\hat{H}_0 t/\hbar) = \left[\hat{O}_I(t), \hat{H}_0 \right].$$

Here we have used the time-independence of the Schrödinger equation together with the observation that any function of an operator commutes with the operator itself.

Interaction picture

In order to solve the equation of motion equation in the interaction picture, we define a unitary operator time-development operator $\hat{U}(t, t')$. Later we will derive its connection with the linked-diagram theorem, which yields a linked expression for the actual operator. The action of the operator on the wave function is

$$|\Psi_I(t)\rangle = \hat{U}(t, t_0)|\Psi_I(t_0)\rangle,$$

with the obvious value $\hat{U}(t_0, t_0) = 1$.

Interaction picture

The time-development operator U has the properties that

$$\hat{U}^\dagger(t, t')\hat{U}(t, t') = \hat{U}(t, t')\hat{U}^\dagger(t, t') = 1,$$

which implies that U is unitary

$$\hat{U}^\dagger(t, t') = \hat{U}^{-1}(t, t').$$

Further,

$$\hat{U}(t, t')\hat{U}(t', t'') = \hat{U}(t, t'')$$

and

$$\hat{U}(t, t')\hat{U}(t', t) = 1,$$

which leads to

$$\hat{U}(t, t') = \hat{U}^\dagger(t', t).$$

Interaction picture

Using our definition of Schrödinger's equation in the interaction picture, we can then construct the operator \hat{U} . We have defined

$$|\Psi_I(t)\rangle = \exp(i\hat{H}_0 t/\hbar)|\Psi_S(t)\rangle,$$

which can be rewritten as

$$|\Psi_I(t)\rangle = \exp(i\hat{H}_0 t/\hbar) \exp(-i\hat{H}(t - t_0)/\hbar)|\Psi_S(t_0)\rangle,$$

or

$$|\Psi_I(t)\rangle = \exp(i\hat{H}_0 t/\hbar) \exp(-i\hat{H}(t - t_0)/\hbar) \exp(-i\hat{H}_0 t_0/\hbar)|\Psi_I(t_0)\rangle.$$

Interaction picture

From the last expression we can define

$$\hat{U}(t, t_0) = \exp(i\hat{H}_0 t/\hbar) \exp(-i\hat{H}(t - t_0)/\hbar) \exp(-i\hat{H}_0 t_0/\hbar).$$

It is then easy to convince oneself that the properties defined above are satisfied by the definition of \hat{U} .

Interaction picture

We derive the equation of motion for \hat{U} using the above definition. This results in

$$i\hbar \frac{\partial}{\partial t} \hat{U}(t, t_0) = \hat{H}_I(t) \hat{U}(t, t_0),$$

which we integrate from t_0 to a time t resulting in

$$\hat{U}(t, t_0) - \hat{U}(t_0, t_0) = \hat{U}(t, t_0) - 1 = -\frac{i}{\hbar} \int_{t_0}^t dt' \hat{H}_I(t') \hat{U}(t', t_0),$$

which can be rewritten as

$$\hat{U}(t, t_0) = 1 - \frac{i}{\hbar} \int_{t_0}^t dt' \hat{H}_I(t') \hat{U}(t', t_0).$$

Interaction picture

We can solve this equation iteratively keeping in mind the time-ordering of the operators

$$\hat{U}(t, t_0) = 1 - \frac{i}{\hbar} \int_{t_0}^t dt' \hat{H}_I(t') + \left(\frac{-i}{\hbar}\right)^2 \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}_I(t') \hat{H}_I(t'') + \dots$$

The third term can be written as

$$\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}_I(t') \hat{H}_I(t'') = \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}_I(t') \hat{H}_I(t'') + \frac{1}{2} \int_{t_0}^t dt'' \int_{t''}^t dt' \hat{H}_I(t') \hat{H}_I(t'').$$

Interaction picture

We obtain this expression by changing the integration order in the second term via a change of the integration variables t' and t'' in

$$\frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}_I(t') \hat{H}_I(t'').$$

We can rewrite the terms which contain the double integral as

$$\begin{aligned} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}_I(t') \hat{H}_I(t'') = \\ \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \left[\hat{H}_I(t') \hat{H}_I(t'') \Theta(t' - t'') + \hat{H}_I(t') \hat{H}_I(t'') \Theta(t'' - t') \right], \end{aligned}$$

with $\Theta(t'' - t')$ being the standard Heavyside or step function. The step function allows us to give a specific time-ordering to the above expression.

Interaction picture

With the Θ -function we can rewrite the last expression as

$$\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}_I(t') \hat{H}_I(t'') = \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{T} [\hat{H}_I(t') \hat{H}_I(t'')] ,$$

where \hat{T} is the so-called time-ordering operator.

Interaction picture

With this definition, we can rewrite the expression for \hat{U} as

$$\hat{U}(t, t_0) = \sum_{n=0}^{\infty} \left(\frac{-i}{\hbar} \right)^n \frac{1}{n!} \int_{t_0}^t dt_1 \cdots \int_{t_0}^t dt_N \hat{T} [\hat{H}_I(t_1) \cdots \hat{H}_I(t_n)] = \hat{T} \exp \left[\frac{-i}{\hbar} \int_{t_0}^t dt' \hat{H}_I(t') \right].$$

The above time-evolution operator in the interaction picture will be used to derive various contributions to many-body perturbation theory. See also exercise 26 for a discussion of the various time orderings.

Heisenberg picture

We wish now to define a unitary transformation in terms of \hat{H} by defining

$$|\Psi_H(t)\rangle = \exp(i\hat{H}t/\hbar)|\Psi_S(t)\rangle,$$

which is again a unitary transformation carried out now at the time t on the wave function in the Schrödinger picture. If we combine this equation with Schrödinger's equation we obtain the following equation of motion

$$i\hbar \frac{\partial}{\partial t} |\Psi_H(t)\rangle = 0,$$

meaning that $|\Psi_H(t)\rangle$ is time independent. An operator in this picture is defined as

$$\hat{O}_H(t) = \exp(i\hat{H}t/\hbar)\hat{O}_S \exp(-i\hat{H}t/\hbar).$$

Heisenberg picture

The time dependence is then in the operator itself, and this yields in turn the following equation of motion

$$i\hbar \frac{\partial}{\partial t} \hat{O}_H(t) = \exp(i\hat{H}t/\hbar) [\hat{O}_H \hat{H} - \hat{H} \hat{O}_H] \exp(-i\hat{H}t/\hbar) = [\hat{O}_H(t), \hat{H}].$$

We note that an operator in the Heisenberg picture can be related to the corresponding operator in the interaction picture as

$$\begin{aligned} \hat{O}_H(t) &= \exp(i\hat{H}t/\hbar) \hat{O}_S \exp(-i\hat{H}t/\hbar) = \\ &\exp(i\hat{H}_I t/\hbar) \exp(-i\hat{H}_0 t/\hbar) \hat{O}_I \exp(i\hat{H}_0 t/\hbar) \exp(-i\hat{H}_I t/\hbar). \end{aligned}$$

Heisenberg picture

With our definition of the time evolution operator we see that

$$\hat{O}_H(t) = \hat{U}(0, t) \hat{O}_I \hat{U}(t, 0),$$

which in turn implies that $\hat{O}_S = \hat{O}_I(0) = \hat{O}_H(0)$, all operators are equal at $t = 0$. The wave function in the Heisenberg formalism is related to the other pictures as

$$|\Psi_H\rangle = |\Psi_S(0)\rangle = |\Psi_I(0)\rangle,$$

since the wave function in the Heisenberg picture is time independent. We can relate this wave function to that at a given time t via the time evolution operator as

$$|\Psi_H\rangle = \hat{U}(0, t) |\Psi_I(t)\rangle.$$

Adiabatic hypothesis

We assume that the interaction term is switched on gradually. Our wave function at time $t = -\infty$ and $t = \infty$ is supposed to represent a non-interacting system given by the solution to the unperturbed part of our Hamiltonian \hat{H}_0 . We assume the ground state is given by $|\Phi_0\rangle$, which could be a Slater determinant. We define our Hamiltonian as

$$\hat{H} = \hat{H}_0 + \exp(-\varepsilon t/\hbar)\hat{H}_I,$$

where ε is a small number. The way we write the Hamiltonian and its interaction term is meant to simulate the switching of the interaction.

Adiabatic hypothesis

The time evolution of the wave function in the interaction picture is then

$$|\Psi_I(t)\rangle = \hat{U}_\varepsilon(t, t_0)|\Psi_I(t_0)\rangle,$$

with

$$\hat{U}_\varepsilon(t, t_0) = \sum_{n=0}^{\infty} \left(\frac{-i}{\hbar} \right)^n \frac{1}{n!} \int_{t_0}^t dt_1 \cdots \int_{t_0}^t dt_N \exp(-\varepsilon(t_1 + \cdots + t_N)/\hbar) \hat{T} [\hat{H}_I(t_1) \cdots \hat{H}_I(t_N)]$$

Adiabatic hypothesis

In the limit $t_0 \rightarrow -\infty$, the solution of Schrödinger's equation is $|\Phi_0\rangle$, and the eigenenergies are given by

$$\hat{H}_0|\Phi_0\rangle = W_0|\Phi_0\rangle,$$

meaning that

$$|\Psi_S(t_0)\rangle = \exp(-iW_0 t_0/\hbar)|\Phi_0\rangle,$$

with the corresponding interaction picture wave function given by

$$|\Psi_I(t_0)\rangle = \exp(i\hat{H}_0 t_0/\hbar)|\Psi_S(t_0)\rangle = |\Phi_0\rangle.$$

Adiabatic hypothesis

The solution becomes time independent in the limit $t_0 \rightarrow -\infty$. The same conclusion can be reached by looking at

$$i\hbar \frac{\partial}{\partial t} |\Psi_I(t)\rangle = \exp(\varepsilon|t|/\hbar) \hat{H}_I |\Psi_I(t)\rangle$$

and taking the limit $t \rightarrow -\infty$. We can rewrite the equation for the wave function at a time $t = 0$ as

$$|\Psi_I(0)\rangle = \hat{U}_\varepsilon(0, -\infty) |\Phi_0\rangle.$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

Our wave function for ground state (after Gell-Mann and Low, see Phys. Rev. **84**, 350 (1951)) is then

$$\frac{|\Psi_0\rangle}{\langle\Phi_0|\Psi_0\rangle} = \lim_{\epsilon \rightarrow 0} \lim_{t' \rightarrow -\infty} \frac{U(0, -\infty)|\Phi_0\rangle}{\langle\Phi_0|U(0, -\infty)|\Phi_0\rangle},$$

and we ask whether this quantity exists to all orders in perturbation theory. Goldstone's theorem states that only linked diagrams enter the expression for the final binding energy. It means that energy difference reads now

$$\Delta E = \sum_{i=0}^{\infty} \langle\Phi_0|\hat{H}_I \left\{ \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I \right\}^i |\Phi_0\rangle_L,$$

where the subscript L indicates that only linked diagrams are included. In our Rayleigh-Schrödinger expansion, the energy difference included also unlinked diagrams.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

If it does, Gell-Mann and Low showed that it is an eigenstate of \hat{H} with eigenvalue

$$\hat{H} \frac{|\Psi_0\rangle}{\langle\Phi_0|\Psi_0\rangle} = E \frac{|\Psi_0\rangle}{\langle\Phi_0|\Psi_0\rangle}$$

and multiplying from the left with $\langle\Phi_0|$ we can rewrite the last equation as

$$E - W_0 = \frac{\langle\Phi_0|\hat{H}|\Psi_0\rangle}{\langle\Phi_0|\Psi_0\rangle},$$

since $\hat{H}_0|\Phi_0\rangle = W_0|\Phi_0\rangle$. The numerator and the denominators of the last equation do not exist separately. The theorem of Gell-Mann and Low asserts that this ratio exists.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

We note that also that the term D is nothing but the denominator of the equation for the energy. We obtain then the following expression for the energy

$$E - W_0 = \Delta E = N_L = \langle \Phi_0(0) | \hat{H}_I U_\epsilon(0, -\infty) | \Phi_0(-\infty) \rangle_L,$$

and Goldstone's theorem is then proved. The corresponding expression from Rayleigh-Schrödinger perturbation theory is given by

$$\Delta E = \langle \Phi_0 | \left(\hat{H}_I + \hat{H}_I \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I + \hat{H}_I \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I + \dots \right) | \Phi_0 \rangle_C.$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

An important point in the derivation of the Gell-Mann and Low theorem

$$E - W_0 = \frac{\langle \Phi_0 | \hat{H}_I | \Psi_0 \rangle}{\langle \Phi_0 | \Psi_0 \rangle},$$

is that the numerator and the denominators of the last equation do not exist separately. The theorem of Gell-Mann and Low asserts that this ratio exists. To prove it we proceed as follows. Consider the expression

$$(\hat{H}_0 - E)U_\epsilon(0, -\infty)|\Phi_0\rangle = [\hat{H}_0, U_\epsilon(0, -\infty)]|\Phi_0\rangle.$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

To evaluate the commutator

$$(\hat{H}_0 - E)U_\epsilon(0, -\infty)|\Phi_0\rangle = [\hat{H}_0, U_\epsilon(0, -\infty)]|\Phi_0\rangle.$$

we write the associate commutator as

$$\begin{aligned} [\hat{H}_0, \hat{H}_I(t_1)\hat{H}_I(t_2)\dots\hat{H}_I(t_n)] &= [\hat{H}_0, \hat{H}_I(t_1)]\hat{H}_I(t_2)\dots\hat{H}_I(t_n) + \\ &\dots + \hat{H}_I(t_1)[\hat{H}_0, \hat{H}_I(t_2)]\hat{H}_I(t_3)\dots\hat{H}_I(t_n) + \dots \end{aligned}$$

Using the equation of motion for an operator in the interaction picture we have

$$i\hbar\frac{\partial}{\partial t}\hat{H}_I(t) = [\hat{H}_I(t), \hat{H}_0].$$

Each of the above commutators yield then a time derivative!

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

We have then

$$\left[\hat{H}_0, \hat{H}_I(t_1) \hat{H}_I(t_2) \dots \hat{H}_I(t_n) \right] = i\hbar \left(\frac{\partial}{\partial t_n} + \frac{\partial}{\partial t_1} + \dots + \frac{\partial}{\partial t_n} \right) \hat{H}_I(t_1) \hat{H}_I(t_2) \dots \hat{H}_I(t_n),$$

meaning that we can rewrite

$$(\hat{H}_0 - E) U_\epsilon(0, -\infty) |\Phi_0\rangle = \left[\hat{H}_0, U_\epsilon(0, -\infty) \right] |\Phi_0\rangle,$$

as

$$\begin{aligned} (\hat{H}_0 - E) U_\epsilon(0, -\infty) |\Phi_0\rangle &= - \sum_{n=1}^{\infty} \left(\frac{-i}{\hbar} \right)^{n-1} \frac{1}{n!} \int_{t_0}^t dt_1 \dots \int_{t_0}^t dt_n \exp(-\epsilon(t_1 + \dots + t_n)/\hbar) \\ &\quad \times \sum_{i=1}^n \left(\frac{\partial}{\partial t_i} \right) \hat{T} \left[\hat{H}_I(t_1) \dots \hat{H}_I(t_n) \right]. \end{aligned}$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

All the time derivatives in this equation

$$(\hat{H}_0 - E)U_\epsilon(0, -\infty)|\Phi_0\rangle = -\sum_{n=1}^{\infty} \left(\frac{-i}{\hbar}\right)^{n-1} \frac{1}{n!} \int_{t_0}^t dt_1 \cdots \int_{t_0}^t dt_N \exp(-\epsilon(t_1 + \cdots + t_n)/\hbar) \\ \times \sum_{i=1}^n \left(\frac{\partial}{\partial t_i}\right) \hat{T} [\hat{H}_I(t_1) \cdots \hat{H}_I(t_n)],$$

make the same contribution, as can be seen by changing dummy variables. We can therefore retain just one time derivative $\partial/\partial t$ and multiply with n . Integrating by parts wrt t_1 we obtain two terms.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

Integrating by parts wrt t_1 one can finally show that

$$\frac{|\Psi_0\rangle}{\langle\Phi_0|\Psi_0\rangle} = \lim_{\epsilon\rightarrow 0} \lim_{t'\rightarrow -\infty} \frac{U(0, -\infty)|\Phi_0\rangle}{\langle\Phi_0|U(0, -\infty)|\Phi_0\rangle},$$

For more details about the derivation, see Gell-Mann and Low, Phys. Rev. **84**, 350 (1951). See also chapter 6.2 of Raimis or Fetter and Walecka, chapter 3.6.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

In the present discussion of the time-dependent theory we will make use of the so-called complex-time approach to describe the time evolution operator U . This means that we allow the time t to be rotated by a small angle ϵ relative to the real time axis. The complex time t is then related to the real time \tilde{t} by

$$t = \tilde{t}(1 - i\epsilon).$$

Let us first study the true eigenvector Ψ_α which evolves from the unperturbed eigenvectors Φ_α through the action of the time development operator

$$U_\epsilon(t, t') = \lim_{\epsilon \rightarrow 0} \lim_{t' \rightarrow -\infty} \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_{t'}^t dt_1 \int_{t'}^{t_1} dt_2 \cdots \int_{t'}^{t_{n-1}} dt_n \\ \times T [H_1(t_1) H_1(t_2) \cdots H_1(t_n)],$$

where T stands for the correct time-ordering.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

In time-dependent perturbation theory we let Ψ_α develop from Φ_α in the remote past to a given time t

$$\frac{|\Psi_\alpha\rangle}{\langle\psi_\alpha|\Psi_\alpha\rangle} = \lim_{\epsilon \rightarrow 0} \lim_{t' \rightarrow -\infty} \frac{U_\epsilon(t, t')|\psi_\alpha\rangle}{\langle\psi_\alpha|U(t, t')|\Phi_\alpha\rangle},$$

and similarly, we let Ψ_β develop from Φ_β in the remote future

$$\frac{\langle\Psi_\beta|}{\langle\psi_\beta|\Psi_\beta\rangle} = \lim_{\epsilon \rightarrow 0} \lim_{t' \rightarrow \infty} \frac{\langle\psi_\beta|U_\epsilon(t', t)}{\langle\psi_\beta|U_\epsilon(t', t)|\Phi_\beta\rangle}.$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

Here we are interested in the expectation value of a given operator \mathcal{O} acting at a time $t = 0$. This can be achieved from the two previous equations defining

$$|\Psi'_{\alpha,\beta}\rangle = \frac{|\Psi_{\alpha,\beta}\rangle}{\langle\Phi_{\alpha,\beta}|\Psi_{\alpha,\beta}\rangle}$$

we have

$$\mathcal{O}_{\alpha\beta} = \frac{N_{\beta\alpha}}{D_{\beta}D_{\alpha}},$$

where we have introduced

$$N_{\beta\alpha} = \langle\Phi_{\beta}|U_{\varepsilon}(\infty, 0)\mathcal{O}U_{\varepsilon}(0, -\infty)|\Phi_{\alpha}\rangle,$$

and

$$D_{\alpha,\beta} = \sqrt{\langle\psi_{\alpha,\beta}|U_{\varepsilon}(\infty, 0)U_{\varepsilon}(0, -\infty)|\Phi_{\alpha,\beta}\rangle}.$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

If the operator \mathcal{O} stands for the hamiltonian H we obtain

$$\frac{\langle \Psi'_\lambda | H | \Psi'_\lambda \rangle}{\langle \Psi'_\lambda | \Psi'_\lambda \rangle}$$

At this stage, *it is important to observe* that our expression for the expectation value of a given operator \mathcal{O} *is hermitian* insofar $\mathcal{O}^\dagger = \mathcal{O}$. This is readily demonstrated. The above equation is of the general form

$$U(t, t_0) \mathcal{O} U(t_0, -t),$$

and noting that

$$U^\dagger(t, t_0) = \left(e^{iH_0 t} e^{-iH(t-t_0)} e^{-iH_0 t} \right)^\dagger = U(t_0, -t),$$

since $H^\dagger = H$ and $H_0^\dagger = H_0$, we have that

$$(U(t, t_0) \mathcal{O} U(t_0, -t))^\dagger = U(t, t_0) \mathcal{O} U(t_0, -t).$$

The question we pose now is what happens in the limit $\varepsilon \rightarrow 0$? Do we get results which are meaningful?

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

Our wave function for ground state is then

$$\frac{|\Psi_0\rangle}{\langle\Phi_0|\Psi_0\rangle} = \lim_{\epsilon \rightarrow 0} \lim_{t' \rightarrow -\infty} \frac{U(0, -\infty)|\Phi_0\rangle}{\langle\Phi_0|U(0, -\infty)|\Phi_0\rangle},$$

meaning that the energy difference is given by

$$E_0 - W_0 = \Delta E_0 = \lim_{\epsilon \rightarrow 0} \lim_{t' \rightarrow -\infty} \frac{\langle\Phi_0|\hat{H}_I U_\epsilon(0, -\infty)|\Phi_0\rangle}{\langle\Phi_0|U_\epsilon(0, -\infty)|\Phi_0\rangle},$$

and we ask whether this quantity exists to all orders in perturbation theory.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

If it does, Gell-Mann and Low showed that it is an eigenstate of \hat{H} with eigenvalue

$$\hat{H} \frac{|\Psi_0\rangle}{\langle \Phi_0 | \Psi_0 \rangle} = E_0 \frac{|\Psi_0\rangle}{\langle \Phi_0 | \Psi_0 \rangle}$$

and multiplying from the left with $\langle \Phi_0 |$ we can rewrite the last equation as

$$E_0 - W_0 = \frac{\langle \Phi_0 | \hat{H}_I | \Psi_0 \rangle}{\langle \Phi_0 | \Psi_0 \rangle},$$

since $\hat{H}_0 |\Phi_0\rangle = W_0 |\Phi_0\rangle$. The numerator and the denominators of the last equation do not exist separately. The theorem of Gell-Mann and Low asserts that this ratio exists.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

Goldstone's theorem states that only linked diagrams enter the expression for the final binding energy. It means that energy difference reads now

$$\Delta E_0 = \sum_{i=0}^{\infty} \langle \Phi_0 | \hat{H}_I \left\{ \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I \right\}^i | \Phi_0 \rangle_L,$$

where the subscript L indicates that only linked diagrams are included. In our Rayleigh-Schrödinger expansion, the energy difference included also unlinked diagrams.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

From this term we can obtain both linked and unlinked contributions. Goldstone's theorem states that only linked diagrams enter the expression for the final binding energy. A linked diagram (or connected diagram) is a diagram which is linked to the last interaction vertex at $t = 0$.

We label the number of linked diagrams with the variable ν and the number of unlinked with μ with $n = \nu + \mu$. The number of unlinked diagrams is then $\mu = n - \nu$.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

In general, the way we can distribute μ unlinked diagrams among the total of n diagrams is given by the combinatorial factor

$$\binom{n}{\mu} = \frac{n!}{\mu! \nu!},$$

and using the following relation

$$\sum_{n=0}^{\infty} \frac{1}{n!} \sum_{\mu+\nu=n}^{\infty} \frac{n!}{\mu! \nu!} = \sum_{\mu=0}^{\infty} \frac{1}{\mu!} \sum_{\nu}^{\infty} \frac{1}{\nu!},$$

we can rewrite the numerator N as

$$N = \langle \Phi_0(0) | \hat{H}_I U_\epsilon(0, -\infty) | \Phi_0(-\infty) \rangle_L \langle \Phi_0(0) | U_\epsilon(0, -\infty) | \Phi_0(-\infty) \rangle = N_L D.$$

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

We define N_L to contain only linked terms with the subscript L indicating that only linked diagrams appear, that is those diagrams which are linked to the last interaction vertex.

Goldstone's theorem and Gell-Mann and Low theorem on the ground state

We note that also that the term D is nothing but the denominator of the equation for the energy. We obtain then the following expression for the energy

$$E_0 - W_0 = \Delta E_0 = N_L = \langle \Phi_0(0) | \hat{H}_I U_\epsilon(0, -\infty) | \Phi_0(-\infty) \rangle_L,$$

and Goldstone's theorem is then proved. The corresponding expression from Rayleigh-Schrödinger perturbation theory is given by

$$\Delta E_0 = \langle \Phi_0 | \left(\hat{H}_I + \hat{H}_I \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I + \hat{H}_I \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I \frac{\hat{Q}}{W_0 - \hat{H}_0} \hat{H}_I + \dots \right) | \Phi_0 \rangle_C.$$