

Slides from FYS4480/9480 Lectures

Morten Hjorth-Jensen

Department of Physics and Center for Computing in Science Education
University of Oslo, N-0316 Oslo, Norway and
Facility for Rare Isotope Beams and Department of Physics and Astronomy,
Michigan State University, East Lansing, MI 48824, USA

Weeks 46 and 47, Fall 2022

CCSD with twobody Hamiltonian

Truncating the cluster operator \hat{T} at the $n = 2$ level, defines CCSD approximation to the Coupled Cluster wavefunction. The coupled cluster wavefunction is now given by

$$|\Psi_{CC}\rangle = e^{\hat{T}_1 + \hat{T}_2} |\Phi_0\rangle$$

where

$$\hat{T}_1 = \sum_{ia} t_i^a a_a^\dagger a_i$$
$$\hat{T}_2 = \frac{1}{4} \sum_{ijab} t_{ij}^{ab} a_a^\dagger a_b^\dagger a_j a_i.$$

CCSD with twobody Hamiltonian cont.

Normal ordered Hamiltonian

$$\begin{aligned}\hat{H} &= \sum_{pq} f_q^p \{ a_p^\dagger a_q \} + \frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \} \\ &\quad + E_0 \\ &= \hat{F}_N + \hat{V}_N + E_0 = \hat{H}_N + E_0\end{aligned}$$

where

$$\begin{aligned}f_q^p &= \langle p | \hat{t} | q \rangle + \sum_i \langle pi | \hat{v} | qi \rangle \\ \langle pq || rs \rangle &= \langle pq | \hat{v} | rs \rangle \\ E_0 &= \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle\end{aligned}$$

Diagram equations - Derivation

Contract \hat{H}_N with \hat{T} in all possible unique combinations that satisfy a given form. The diagram equation is the sum of all these diagrams.

- ▶ Contract one \hat{H}_N element with 0, 1 or multiple \hat{T} elements.
- ▶ All \hat{T} elements must have **atleast** one contraction with \hat{H}_N .
- ▶ No contractions between \hat{T} elements are allowed.
- ▶ A single \hat{T} element can contract with a single element of \hat{H}_N in different ways.

Diagram equations - Derivation

Contract \hat{H}_N with \hat{T} in all possible unique combinations that satisfy a given form. The diagram equation is the sum of all these diagrams.

- ▶ Contract one \hat{H}_N element with 0, 1 or multiple \hat{T} elements.
- ▶ All \hat{T} elements must have **atleast** one contraction with \hat{H}_N .
- ▶ No contractions between \hat{T} elements are allowed.
- ▶ A single \hat{T} element can contract with a single element of \hat{H}_N in different ways.

Diagram equations - Derivation

Contract \hat{H}_N with \hat{T} in all possible unique combinations that satisfy a given form. The diagram equation is the sum of all these diagrams.

- ▶ Contract one \hat{H}_N element with 0, 1 or multiple \hat{T} elements.
- ▶ All \hat{T} elements must have **atleast** one contraction with \hat{H}_N .
- ▶ No contractions between \hat{T} elements are allowed.
- ▶ A single \hat{T} element can contract with a single element of \hat{H}_N in different ways.

Diagram equations - Derivation

Contract \hat{H}_N with \hat{T} in all possible unique combinations that satisfy a given form. The diagram equation is the sum of all these diagrams.

- ▶ Contract one \hat{H}_N element with 0, 1 or multiple \hat{T} elements.
- ▶ All \hat{T} elements must have **atleast** one contraction with \hat{H}_N .
- ▶ No contractions between \hat{T} elements are allowed.
- ▶ A single \hat{T} element can contract with a single element of \hat{H}_N in different ways.

Diagram equations - Derivation

Contract \hat{H}_N with \hat{T} in all possible unique combinations that satisfy a given form. The diagram equation is the sum of all these diagrams.

- ▶ Contract one \hat{H}_N element with 0, 1 or multiple \hat{T} elements.
- ▶ All \hat{T} elements must have **atleast** one contraction with \hat{H}_N .
- ▶ No contractions between \hat{T} elements are allowed.
- ▶ A single \hat{T} element can contract with a single element of \hat{H}_N in different ways.

Diagram elements - Directed lines




Figure: Particle line



Figure: Hole line

- ▶ Represents a contraction between second quantized operators.
- ▶ External lines are connected to one operator vertex and infinity.
- ▶ Internal lines are connected to operator vertices in both ends.

Diagram elements - Onebody Hamiltonian



- ▶ Horizontal dashed line segment with one vertex.
- ▶ Excitation level identify the number of particle/hole pairs created by the operator.

Diagram elements - Twobody Hamiltonian




Level: -2



Level: -1




Level: -1



Level: 0




Level: 0



Level: 0



Level: +1



Level: +1



Level: +2

Diagram elements - Onebody cluster operator



Level: +1

- ▶ Horizontal line segment with one vertex.
- ▶ Excitation level of +1.

Diagram elements - Twobody cluster operator




Level: +2

- ▶ Horizontal line segment with two vertices.
- ▶ Excitation level of +2.


CCSD energy equation - Derivation

$$E_{\text{CCSD}} = \langle \Phi_0 || \Phi_0 \rangle$$


- ▶ No external lines.
- ▶ Final excitation level: 0



Elements: \hat{H}_N



Elements: \hat{T}



CCSD energy equation

$$E_{CCSD} = \text{[Diagram 1]} + \text{[Diagram 2]} + \text{[Diagram 3]}$$

The equation represents the CCSD energy equation using Feynman diagrams. The first term is a single vertex with two incoming lines (one solid, one dashed) and two outgoing lines (one solid, one dashed), with a dashed line extending to the right ending in an 'x'. The second term is a two-vertex diagram with a solid line connecting the two vertices. The third term is a two-vertex diagram with a dashed line connecting the two vertices.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

CCSD energy equation


$$E_{CCSD} = f_a^i t_i^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

Note the implicit sum over repeated indices.


CCSD \hat{T}_1 amplitude equation - Derivation

$$0 = \langle \Phi_i^a || \Phi_0 \rangle$$


- ▶ One pair of particle/hole external lines.
- ▶ Final excitation level: +1



Elements: \hat{H}_N



Elements: \hat{T}



CCSD \hat{T}_1 amplitude equation

$$0 =$$

The equation is represented as $0 =$ followed by a sum of 14 Feynman diagrams. The diagrams are arranged in four rows:

- Row 1: 4 diagrams
- Row 2: 4 diagrams
- Row 3: 4 diagrams
- Row 4: 2 diagrams

Each diagram consists of solid lines with arrows and dashed lines with 'x' marks. The diagrams represent various terms in the CCSD \hat{T}_1 amplitude equation, including terms with one, two, and three loops.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} | | \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout}, \text{rout} || \text{lin}, \text{rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin}, \text{rin}}^{\text{lout}, \text{rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.


CCSD \hat{T}_1 amplitude equation

$$\begin{aligned} 0 = & f_i^a + f_e^a t_i^e - f_i^m t_m^a + \langle ma || ei \rangle t_m^e + f_e^m t_{im}^{ae} + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} \\ & - \frac{1}{2} \langle mn || ei \rangle t_{mn}^{ea} - f_e^m t_i^e t_m^a + \langle am || ef \rangle t_i^e t_m^f - \langle mn || ei \rangle t_m^e t_n^a \\ & + \langle mn || ef \rangle t_m^e t_{ni}^{fa} - \frac{1}{2} \langle mn || ef \rangle t_i^e t_{mn}^{af} - \frac{1}{2} \langle mn || ef \rangle t_n^a t_{mi}^{ef} \\ & - \langle mn || ef \rangle t_i^e t_m^a t_n^f \end{aligned}$$


CCSD \hat{T}_2 amplitude equation - Derivation

$$0 = \langle \Phi_{ij}^{ab} || \Phi_0 \rangle$$


- ▶ Two pairs of particle/hole external lines.
- ▶ Final excitation level: +2



Elements: \hat{H}_N



Elements: \hat{T}



CCSD \hat{T}_2 amplitude equation

$$\begin{aligned}
 0 = & \text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \text{diagram 4} \times + \times \text{diagram 5} + \text{diagram 6} \\
 & + \text{diagram 7} + \text{diagram 8} + \text{diagram 9} + \text{diagram 10} + \text{diagram 11} \\
 & + \text{diagram 12} + \text{diagram 13} + \text{diagram 14} + \text{diagram 15} + \text{diagram 16} \\
 & + \text{diagram 17} \times + \text{diagram 18} + \text{diagram 19} + \text{diagram 20} + \text{diagram 21} \\
 & + \text{diagram 22} + \text{diagram 23} + \text{diagram 24} + \text{diagram 25} + \text{diagram 26} \\
 & + \text{diagram 27} + \text{diagram 28} + \text{diagram 29} + \text{diagram 30} + \text{diagram 31}
 \end{aligned}$$

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} | | \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} | | \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} || \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} | | \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} | | \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} | | \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

Diagram rules

- ▶ Label all lines.
- ▶ Sum over all internal indices.
- ▶ Extract matrix elements. $(f_{\text{in}}^{\text{out}}, \langle \text{lout, rout} | | \text{lin, rin} \rangle)$
- ▶ Extract cluster amplitudes with indices in the order left to right. Incoming lines are subscripts, while outgoing lines are superscripts. $(t_{\text{in}}^{\text{out}}, t_{\text{lin, rin}}^{\text{lout, rout}})$
- ▶ Calculate the phase: $(-1)^{\text{holelines} + \text{loops}}$
- ▶ Multiply by a factor of $\frac{1}{2}$ for each equivalent line and each equivalent vertex.
- ▶ Antisymmetrize a pair of external particle/hole line that does not connect to the same operator.

CCSD \hat{T}_2 amplitude equation

$$\begin{aligned}
 0 = & \langle ab||ij \rangle + P(ij)\langle ab||ej \rangle t_i^e - P(ab)\langle am||ij \rangle t_m^b + P(ab)t_e^b t_{ij}^{ae} - P(ij)f_i^m t_{mj}^{ab} \\
 & + \frac{1}{2}\langle ab||ef \rangle t_{ij}^{ef} + \frac{1}{2}\langle mn||ij \rangle t_{mn}^{ab} + P(ij)P(ab)\langle mb||ej \rangle t_{im}^{ae} \\
 & + \frac{1}{2}P(ij)\langle ab||ef \rangle t_i^e t_j^f + \frac{1}{2}P(ab)\langle mn||ij \rangle t_m^a t_n^b - P(ij)P(ab)\langle mb||ej \rangle t_i^e t_m^a \\
 & + \frac{1}{4}\langle mn||ef \rangle t_{ij}^{ef} t_{mn}^{ab} + \frac{1}{2}P(ij)P(ab)\langle mn||ef \rangle t_{im}^{ae} t_{nj}^{fb} - \frac{1}{2}P(ab)\langle mn||ef \rangle t_{ij}^{ae} t_{mn}^{bf} \\
 & - \frac{1}{2}P(ij)\langle mn||ef \rangle t_{mi}^{ef} t_{nj}^{ab} - P(ij)f_e^m t_i^e t_{mj}^{ab} - P(ab)f_e^m t_{ij}^{ae} t_m^b \\
 & + P(ij)P(ab)\langle am||ef \rangle t_i^e t_{mj}^{fb} - \frac{1}{2}P(ab)\langle am||ef \rangle t_{ij}^{ef} t_m^b + P(ab)\langle bm||ef \rangle t_{ij}^{ae} t_m^f \\
 & - P(ij)P(ab)\langle mn||ej \rangle t_{im}^{ae} t_n^b + \frac{1}{2}P(ij)\langle mn||ej \rangle t_i^e t_{mn}^{ab} - P(ij)\langle mn||ei \rangle t_m^e t_{nj}^{ab} \\
 & - \frac{1}{2}P(ij)P(ab)\langle am||ef \rangle t_i^e t_j^f t_m^b + \frac{1}{2}P(ij)P(ab)\langle mn||ej \rangle t_i^e t_m^a t_n^b \\
 & + \frac{1}{4}P(ij)\langle mn||ef \rangle t_i^e t_{mn}^{ab} t_j^f - P(ij)P(ab)\langle mn||ef \rangle t_i^e t_m^a t_{nj}^{fb} \\
 & + \frac{1}{4}P(ab)\langle mn||ef \rangle t_m^a t_{ij}^{ef} t_n^b - P(ij)\langle mn||ef \rangle t_m^e t_i^f t_{nj}^{ab} - P(ab)\langle mn||ef \rangle t_{ij}^{ae} t_m^b t_n^f \\
 & + \frac{1}{4}P(ij)P(ab)\langle mn||ef \rangle t_i^e t_m^a t_j^f t_n^b
 \end{aligned}$$

The expansion

$$E_{CC} = \langle \psi_0 | \left(\hat{H}_N + [\hat{H}_N, \hat{T}] + \frac{1}{2} [[\hat{H}_N, \hat{T}], \hat{T}] + \frac{1}{3!} [[[\hat{H}_N, \hat{T}], \hat{T}], \hat{T}] \right. \\ \left. + \frac{1}{4!} [[[[\hat{H}_N, \hat{T}], \hat{T}], \hat{T}], \hat{T}] + \dots \right) | \psi_0 \rangle$$

$$0 = \langle \psi_{ij\dots}^{ab\dots} | \left(\hat{H}_N + [\hat{H}_N, \hat{T}] + \frac{1}{2} [[\hat{H}_N, \hat{T}], \hat{T}] + \frac{1}{3!} [[[\hat{H}_N, \hat{T}], \hat{T}], \hat{T}] \right. \\ \left. + \frac{1}{4!} [[[[\hat{H}_N, \hat{T}], \hat{T}], \hat{T}], \hat{T}] + \dots \right) | \psi_0 \rangle$$

The CCSD energy equation revisited

The expanded CC energy equation involves an infinite sum over nested commutators

$$\begin{aligned} E_{CC} = \langle \Psi_0 | & \left(\hat{H}_N + [\hat{H}_N, \hat{T}] + \frac{1}{2} [[\hat{H}_N, \hat{T}], \hat{T}] \right. \\ & + \frac{1}{3!} [[[\hat{H}_N, \hat{T}], \hat{T}], \hat{T}] \\ & \left. + \frac{1}{4!} [[[[\hat{H}_N, \hat{T}], \hat{T}], \hat{T}], \hat{T}] + \dots \right) | \Psi_0 \rangle, \end{aligned}$$

but fortunately we can show that it truncates naturally, depending on the Hamiltonian.

The first term is zero by construction.

$$\langle \Psi_0 | \hat{H}_N | \Psi_0 \rangle = 0$$

The CCSD energy equation revisited.

The second term can be split up into different pieces

$$\langle \Psi_0 | [\hat{H}_N, \hat{T}] | \Psi_0 \rangle = \langle \Psi_0 | \left([\hat{F}_N, \hat{T}_1] + [\hat{F}_N, \hat{T}_2] + [\hat{V}_N, \hat{T}_1] + [\hat{V}_N, \hat{T}_2] \right) | \Psi_0 \rangle$$

Since we need the explicit expressions for the commutators both in the next term and in the amplitude equations, we calculate them separately.

The expansion - $[\hat{F}_N, \hat{T}_1]$

$$\begin{aligned}
 [\hat{F}_N, \hat{T}_1] &= \sum_{pqia} \left(f_q^p \{a_p^\dagger a_q\} t_i^a \{a_a^\dagger a_i\} - t_i^a \{a_a^\dagger a_i\} f_q^p \{a_p^\dagger a_q\} \right) \\
 &= \sum_{pqia} f_q^p t_i^a \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} \right)
 \end{aligned}$$

$$\{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \{a_a^\dagger a_i a_p^\dagger a_q\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \{a_p^\dagger a_q a_a^\dagger a_i\} + \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$= \{a_p^\dagger a_q a_a^\dagger a_i\} + \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}$$

The expansion - $\left[\hat{F}_N, \hat{T}_1 \right]$

$$\begin{aligned} \left[\hat{F}_N, \hat{T}_1 \right] &= \sum_{pqia} \left(f_q^p \left\{ a_p^\dagger a_q \right\} t_i^a \left\{ a_a^\dagger a_i \right\} - t_i^a \left\{ a_a^\dagger a_i \right\} f_q^p \left\{ a_p^\dagger a_q \right\} \right) \\ &= \sum_{pqia} f_q^p t_i^a \left(\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_i \right\} - \left\{ a_a^\dagger a_i \right\} \left\{ a_p^\dagger a_q \right\} \right) \end{aligned}$$

$$\left\{ a_a^\dagger a_i \right\} \left\{ a_p^\dagger a_q \right\} = \left\{ a_a^\dagger a_i a_p^\dagger a_q \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_i \right\}$$

$$\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_i \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_i \right\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_i} \right\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\}$$

$$= \left\{ a_p^\dagger a_q a_a^\dagger a_i \right\} + \delta_{qa} \left\{ a_p^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger \right\} + \delta_{qa} \delta_{pi}$$

The expansion - $[\hat{F}_N, \hat{T}_1]$

$$\begin{aligned}
 [\hat{F}_N, \hat{T}_1] &= \sum_{pqia} \left(f_q^p \{a_p^\dagger a_q\} t_i^a \{a_a^\dagger a_i\} - t_i^a \{a_a^\dagger a_i\} f_q^p \{a_p^\dagger a_q\} \right) \\
 &= \sum_{pqia} f_q^p t_i^a \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} \right)
 \end{aligned}$$

$$\{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \{a_a^\dagger a_i a_p^\dagger a_q\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \} + \{ a_p^\dagger \overline{a_q a_a^\dagger a_i} \}$$

$$+ \{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \}$$

$$= \{a_p^\dagger a_q a_a^\dagger a_i\} + \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}$$

The expansion - $[\hat{F}_N, \hat{T}_1]$

$$\begin{aligned}
 [\hat{F}_N, \hat{T}_1] &= \sum_{pqia} \left(f_q^p \{a_p^\dagger a_q\} t_i^a \{a_a^\dagger a_i\} - t_i^a \{a_a^\dagger a_i\} f_q^p \{a_p^\dagger a_q\} \right) \\
 &= \sum_{pqia} f_q^p t_i^a \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} \right)
 \end{aligned}$$

$$\{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \{a_a^\dagger a_i a_p^\dagger a_q\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_i} \right\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\}$$

$$= \{a_p^\dagger a_q a_a^\dagger a_i\} + \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}$$

The expansion - $[\hat{F}_N, \hat{T}_1]$

$$\begin{aligned}
 [\hat{F}_N, \hat{T}_1] &= \sum_{pqia} \left(f_q^p \{a_p^\dagger a_q\} t_i^a \{a_a^\dagger a_i\} - t_i^a \{a_a^\dagger a_i\} f_q^p \{a_p^\dagger a_q\} \right) \\
 &= \sum_{pqia} f_q^p t_i^a \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} \right)
 \end{aligned}$$

$$\{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \{a_a^\dagger a_i a_p^\dagger a_q\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_i} \right\}$$

$$+ \left\{ \overline{\overline{a_p^\dagger a_q a_a^\dagger a_i}} \right\}$$

$$= \{a_p^\dagger a_q a_a^\dagger a_i\} + \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}$$

The expansion - $[\hat{F}_N, \hat{T}_1]$

$$\begin{aligned}
 [\hat{F}_N, \hat{T}_1] &= \sum_{pqia} \left(f_q^p \{a_p^\dagger a_q\} t_i^a \{a_a^\dagger a_i\} - t_i^a \{a_a^\dagger a_i\} f_q^p \{a_p^\dagger a_q\} \right) \\
 &= \sum_{pqia} f_q^p t_i^a \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} \right)
 \end{aligned}$$

$$\{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \{a_a^\dagger a_i a_p^\dagger a_q\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_i} \right\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\}$$

$$= \{a_p^\dagger a_q a_a^\dagger a_i\} + \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}$$

The expansion - $[\hat{F}_N, \hat{T}_1]$

$$\begin{aligned}
 [\hat{F}_N, \hat{T}_1] &= \sum_{pqia} \left(f_q^p \{a_p^\dagger a_q\} t_i^a \{a_a^\dagger a_i\} - t_i^a \{a_a^\dagger a_i\} f_q^p \{a_p^\dagger a_q\} \right) \\
 &= \sum_{pqia} f_q^p t_i^a \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} \right)
 \end{aligned}$$

$$\{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \{a_a^\dagger a_i a_p^\dagger a_q\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} = \{a_p^\dagger a_q a_a^\dagger a_i\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_i} \right\}$$

$$+ \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_i} \right\}$$

$$= \{a_p^\dagger a_q a_a^\dagger a_i\} + \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}$$

The expansion - $[\hat{F}_N, \hat{T}_1]$

Wicks theorem gives us

$$\{a_p^\dagger a_q\} \{a_a^\dagger a_i\} - \{a_a^\dagger a_i\} \{a_p^\dagger a_q\} = \delta_{qa} \{a_p^\dagger a_i\} + \delta_{pi} \{a_q a_a^\dagger\} + \delta_{qa} \delta_{pi}.$$

Inserted into the original expression, we arrive at the explicit value of the commutator

$$\begin{aligned} [\hat{F}_N, \hat{T}_1] &= \sum_{pai} f_a^p t_i^a \{a_p^\dagger a_i\} + \sum_{qai} f_q^i t_i^a \{a_q a_a^\dagger\} + \sum_{ai} f_a^i t_i^a \\ &= (\hat{F}_N \hat{T}_1)_c. \end{aligned}$$

The subscript means that the product only includes terms where the operators are connected by atleast one shared index.

The expansion - $[\hat{F}_N, \hat{T}_2]$

$$\begin{aligned} [\hat{F}_N, \hat{T}_2] &= \left[\sum_{pq} f_q^p \{a_p^\dagger a_q\}, \frac{1}{4} \sum_{ijab} t_{ij}^{ab} \{a_a^\dagger a_b^\dagger a_j a_i\} \right] \\ &= \frac{1}{4} \sum_{\substack{pq \\ ijab}} [\{a_p^\dagger a_q\}, \{a_a^\dagger a_b^\dagger a_j a_i\}] \\ &= \frac{1}{4} \sum_{\substack{pq \\ ijab}} f_q^p t_{ij}^{ab} \left(\{a_p^\dagger a_q\} \{a_a^\dagger a_b^\dagger a_j a_i\} - \{a_a^\dagger a_b^\dagger a_j a_i\} \{a_p^\dagger a_q\} \right) \end{aligned}$$

The expansion - $\left[\hat{F}_N, \hat{T}_2 \right]$

$$\left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} = \left\{ a_a^\dagger a_b^\dagger a_j a_i a_p^\dagger a_q \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\}$$

$$\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \\ + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\}$$

The expansion - $\left[\hat{F}_N, \hat{T}_2 \right]$

$$\left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} = \left\{ a_a^\dagger a_b^\dagger a_j a_i a_p^\dagger a_q \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\}$$

$$\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \\ + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\}$$

The expansion - $\left[\hat{F}_N, \hat{T}_2 \right]$

$$\left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} = \left\{ a_a^\dagger a_b^\dagger a_j a_i a_p^\dagger a_q \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\}$$

$$\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \\ + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\}$$

The expansion - $\left[\hat{F}_N, \hat{T}_2 \right]$

$$\left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} = \left\{ a_a^\dagger a_b^\dagger a_j a_i a_p^\dagger a_q \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\}$$

$$\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \\ + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\}$$

The expansion - $\left[\hat{F}_N, \hat{T}_2 \right]$

$$\left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} = \left\{ a_a^\dagger a_b^\dagger a_j a_i a_p^\dagger a_q \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\}$$

$$\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ = \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \\ + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\}$$

The expansion - $\left[\hat{F}_N, \hat{T}_2 \right]$

$$\begin{aligned} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} &= \left\{ a_a^\dagger a_b^\dagger a_j a_i a_p^\dagger a_q \right\} \\ &= \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} \end{aligned}$$

$$\begin{aligned} \left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} &= \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ &\quad + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ a_p^\dagger \overline{a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ &\quad + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} + \left\{ \overline{a_p^\dagger a_q a_a^\dagger a_b^\dagger} a_j a_i \right\} \\ &= \left\{ a_p^\dagger a_q a_a^\dagger a_b^\dagger a_j a_i \right\} - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \\ &\quad + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ &\quad + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\} \end{aligned}$$

The expansion - $[\hat{F}_N, \hat{T}_2]$

Wicks theorem gives us

$$\begin{aligned} & \left(\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} - \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} \right) = \\ & - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} \\ & - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} \\ & - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\} \end{aligned}$$

Inserted into the original expression, we arrive at

$$\begin{aligned} [\hat{F}_N, \hat{T}_2] &= \frac{1}{4} \sum_{\substack{pq \\ abij}} f_q^p t_{ij}^{ab} \left(-\delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \right. \\ &+ \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ &\left. + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\} \right). \end{aligned}$$

The expansion - $[\hat{F}_N, \hat{T}_2]$

Wicks theorem gives us

$$\begin{aligned} & \left(\left\{ a_p^\dagger a_q \right\} \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} - \left\{ a_a^\dagger a_b^\dagger a_j a_i \right\} \left\{ a_p^\dagger a_q \right\} \right) = \\ & - \delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} + \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} \\ & - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} \\ & - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\} \end{aligned}$$

Inserted into the original expression, we arrive at

$$\begin{aligned} [\hat{F}_N, \hat{T}_2] &= \frac{1}{4} \sum_{\substack{pq \\ abij}} f_{ij}^p t_{ij}^{ab} \left(-\delta_{pj} \left\{ a_q a_a^\dagger a_b^\dagger a_i \right\} + \delta_{pi} \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \right. \\ &+ \delta_{qa} \left\{ a_p^\dagger a_b^\dagger a_j a_i \right\} - \delta_{qb} \left\{ a_p^\dagger a_a^\dagger a_j a_i \right\} - \delta_{pj} \delta_{qa} \left\{ a_b^\dagger a_i \right\} \\ &\left. + \delta_{pi} \delta_{qa} \left\{ a_b^\dagger a_j \right\} + \delta_{pj} \delta_{qb} \left\{ a_a^\dagger a_i \right\} - \delta_{pi} \delta_{qb} \left\{ a_a^\dagger a_j \right\} \right). \end{aligned}$$

The expansion - $[\hat{F}_N, \hat{T}_2]$

After renaming indices and changing the order of operators, we arrive at the explicit expression

$$\begin{aligned} [\hat{F}_N, \hat{T}_2] &= \frac{1}{2} \sum_{qijab} f_q^i t_{ij}^{ab} \{a_q a_a^\dagger a_b^\dagger a_j\} + \frac{1}{2} \sum_{pijab} f_a^p t_{ij}^{ab} \{a_p^\dagger a_b^\dagger a_j a_i\} \\ &\quad + \sum_{ijab} f_a^i t_{ij}^{ab} \{a_b^\dagger a_j\} \\ &= \left(\hat{F}_N \hat{T}_2 \right)_c. \end{aligned}$$

The subscript implies that only the connected terms from the product contribute.

The expansion - $\frac{1}{2} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right]$

$$\left[\hat{F}_N, \hat{T}_1 \right] = \sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a$$

$$\begin{aligned} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right] &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\}, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \sum_{pabij} f_a^p t_i^a t_j^b \left[\left\{ a_p^\dagger a_i \right\}, \left\{ a_b^\dagger a_j \right\} \right] + \sum_{qabij} f_q^i t_i^a t_j^b \left[\left\{ a_q a_a^\dagger \right\}, \left\{ a_b^\dagger a_j \right\} \right] \end{aligned}$$

$$\left\{ a_b^\dagger a_j \right\} \left\{ a_p^\dagger a_i \right\} = \left\{ a_b^\dagger a_j a_p^\dagger a_i \right\} = \left\{ a_p^\dagger a_i a_b^\dagger a_j \right\}$$

$$\left\{ a_b^\dagger a_j \right\} \left\{ a_q a_a^\dagger \right\} = \left\{ a_b^\dagger a_j a_q a_a^\dagger \right\} = \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\}$$

The expansion - $\frac{1}{2} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right]$

$$\left[\hat{F}_N, \hat{T}_1 \right] = \sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a$$

$$\begin{aligned} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right] &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\}, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \sum_{pabij} f_a^p t_i^a t_j^b \left[\left\{ a_p^\dagger a_i \right\}, \left\{ a_b^\dagger a_j \right\} \right] + \sum_{qabij} f_q^i t_i^a t_j^b \left[\left\{ a_q a_a^\dagger \right\}, \left\{ a_b^\dagger a_j \right\} \right] \end{aligned}$$

$$\left\{ a_b^\dagger a_j \right\} \left\{ a_p^\dagger a_i \right\} = \left\{ a_b^\dagger a_j a_p^\dagger a_i \right\} = \left\{ a_p^\dagger a_i a_b^\dagger a_j \right\}$$

$$\left\{ a_b^\dagger a_j \right\} \left\{ a_q a_a^\dagger \right\} = \left\{ a_b^\dagger a_j a_q a_a^\dagger \right\} = \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\}$$

The expansion - $\frac{1}{2} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right]$

$$\left[\hat{F}_N, \hat{T}_1 \right] = \sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a$$

$$\begin{aligned} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right] &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\}, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \sum_{pabij} f_a^p t_i^a t_j^b \left[\left\{ a_p^\dagger a_i \right\}, \left\{ a_b^\dagger a_j \right\} \right] + \sum_{qabij} f_q^i t_i^a t_j^b \left[\left\{ a_q a_a^\dagger \right\}, \left\{ a_b^\dagger a_j \right\} \right] \end{aligned}$$

$$\begin{aligned} \left\{ a_b^\dagger a_j \right\} \left\{ a_p^\dagger a_i \right\} &= \left\{ a_b^\dagger a_j a_p^\dagger a_i \right\} = \left\{ a_p^\dagger a_i a_b^\dagger a_j \right\} \\ \left\{ a_b^\dagger a_j \right\} \left\{ a_q a_a^\dagger \right\} &= \left\{ a_b^\dagger a_j a_q a_a^\dagger \right\} = \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \end{aligned}$$

The expansion - $\frac{1}{2} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right]$

$$\left[\hat{F}_N, \hat{T}_1 \right] = \sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a$$

$$\begin{aligned} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right] &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\} + \sum_{ai} f_a^i t_i^a, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \left[\sum_{pai} f_a^p t_i^a \left\{ a_p^\dagger a_i \right\} + \sum_{qai} f_q^i t_i^a \left\{ a_q a_a^\dagger \right\}, \sum_{jb} t_j^b \left\{ a_b^\dagger a_j \right\} \right] \\ &= \sum_{pabij} f_a^p t_i^a t_j^b \left[\left\{ a_p^\dagger a_i \right\}, \left\{ a_b^\dagger a_j \right\} \right] + \sum_{qabij} f_q^i t_i^a t_j^b \left[\left\{ a_q a_a^\dagger \right\}, \left\{ a_b^\dagger a_j \right\} \right] \end{aligned}$$

$$\begin{aligned} \left\{ a_b^\dagger a_j \right\} \left\{ a_p^\dagger a_i \right\} &= \left\{ a_b^\dagger a_j a_p^\dagger a_i \right\} = \left\{ a_p^\dagger a_i a_b^\dagger a_j \right\} \\ \left\{ a_b^\dagger a_j \right\} \left\{ a_q a_a^\dagger \right\} &= \left\{ a_b^\dagger a_j a_q a_a^\dagger \right\} = \left\{ a_q a_a^\dagger a_b^\dagger a_j \right\} \end{aligned}$$

The expansion - $\left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right]$

$$\begin{aligned} \frac{1}{2} \left[\left[\hat{F}_N, \hat{T}_1 \right], \hat{T}_1 \right] &= \frac{1}{2} \left(\sum_{pabij} f_a^p t_i^a t_j^b \delta_{pj} \left\{ a_i a_b^\dagger \right\} - \sum_{qabij} f_q^j t_i^a t_j^b \delta_{qb} \left\{ a_a^\dagger a_j \right\} \right) \\ &= -\frac{1}{2} 2 \sum_{abij} f_b^j t_j^a t_i^b \left\{ a_a^\dagger a_i \right\} \\ &= - \sum_{abij} f_b^j t_j^a t_i^b \left\{ a_a^\dagger a_i \right\} \\ &= \frac{1}{2} \left(\hat{F}_N \hat{T}_1^2 \right)_c \end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}\langle \Phi_0 | [\hat{V}_N, \hat{T}_1] | \Phi_0 \rangle &= \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \sum_{ia} t_i^a \{ a_a^\dagger a_i \} \right] | \Phi_0 \rangle \\ &= \frac{1}{4} \sum_{\substack{pqrs \\ sia}} \langle pq || rs \rangle t_i^a \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_i \} \right] | \Phi_0 \rangle \\ &= 0\end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}\langle \Phi_0 | [\hat{V}_N, \hat{T}_1] | \Phi_0 \rangle &= \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \sum_{ia} t_i^a \{ a_a^\dagger a_i \} \right] | \Phi_0 \rangle \\ &= \frac{1}{4} \sum_{\substack{pqrs \\ sia}} \langle pq || rs \rangle t_i^a \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_i \} \right] | \Phi_0 \rangle \\ &= 0\end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}\langle \Phi_0 | [\hat{V}_N, \hat{T}_1] | \Phi_0 \rangle &= \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \sum_{ia} t_i^a \{ a_a^\dagger a_i \} \right] | \Phi_0 \rangle \\ &= \frac{1}{4} \sum_{\substack{pqrs \\ sia}} \langle pq || rs \rangle t_i^a \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_i \} \right] | \Phi_0 \rangle \\ &= 0\end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}
 \langle \Phi_0 | [\hat{V}_N, \hat{T}_2] | \Phi_0 \rangle &= \\
 \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \frac{1}{4} \sum_{ijab} t_{ij}^{ab} \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left(\left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right. &= \\
 \left. \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right) | \Phi_0 \rangle &= \\
 = \frac{1}{4} \sum_{ijab} \langle ij || ab \rangle t_{ij}^{ab} &
 \end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}
 \langle \Phi_0 | [\hat{V}_N, \hat{T}_2] | \Phi_0 \rangle &= \\
 \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \frac{1}{4} \sum_{ijab} t_{ij}^{ab} \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left(\left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right. & \\
 \left. \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right) | \Phi_0 \rangle &= \\
 = \frac{1}{4} \sum_{ijab} \langle ij || ab \rangle t_{ij}^{ab} &
 \end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}
 \langle \Phi_0 | [\hat{V}_N, \hat{T}_2] | \Phi_0 \rangle &= \\
 \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \frac{1}{4} \sum_{ijab} t_{ij}^{ab} \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left(\left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right. &= \\
 \left. \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right) | \Phi_0 \rangle &= \\
 = \frac{1}{4} \sum_{ijab} \langle ij || ab \rangle t_{ij}^{ab} &
 \end{aligned}$$

The CCSD energy equation revisited

$$\begin{aligned}
 \langle \Phi_0 | [\hat{V}_N, \hat{T}_2] | \Phi_0 \rangle &= \\
 \langle \Phi_0 | \left[\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle \{ a_p^\dagger a_q^\dagger a_s a_r \}, \frac{1}{4} \sum_{ijab} t_{ij}^{ab} \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left[\{ a_p^\dagger a_q^\dagger a_s a_r \}, \{ a_a^\dagger a_b^\dagger a_j a_i \} \right] | \Phi_0 \rangle &= \\
 = \frac{1}{16} \sum_{\substack{pqr \\ sijab}} \langle pq || rs \rangle t_{ij}^{ab} \langle \Phi_0 | \left(\left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right. &= \\
 \left. \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} + \left\{ \overline{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_b^\dagger a_j a_i} \right\} \right) | \Phi_0 \rangle &= \\
 = \frac{1}{4} \sum_{ijab} \langle ij || ab \rangle t_{ij}^{ab} &
 \end{aligned}$$

The CCSD energy equation revisited

The CCSD energy get two contributions from $(\hat{H}_N \hat{T})_c$

$$\begin{aligned} E_{CC} &\Leftarrow \langle \Phi_0 | [\hat{H}_N, \hat{T}] | \Phi_0 \rangle \\ &= \sum_{ia} f_a^i t_i^a + \frac{1}{4} \sum_{ijab} \langle ij || ab \rangle t_{ij}^{ab} \end{aligned}$$

The CCSD energy equation revisited

$$E_{CC} \Leftarrow \langle \Phi_0 | \frac{1}{2} \left(\hat{H}_N \hat{T}^2 \right)_c | \Phi_0 \rangle$$

$$\begin{aligned} \langle \Phi_0 | \frac{1}{2} \left(\hat{V}_N \hat{T}_1^2 \right)_c | \Phi_0 \rangle &= \\ \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \left(\left\{ a_p^\dagger a_q^\dagger a_s a_r \right\} \left\{ a_a^\dagger a_i \right\} \left\{ a_b^\dagger a_j \right\} \right)_c | \Phi_0 \rangle \\ &= \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \\ &\quad \left(\left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{\text{diag 1}} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{\text{diag 2}} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{\text{diag 3}} \right\} \right. \\ &\quad \left. + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{\text{diag 4}} \right\} \right) | \Phi_0 \rangle \\ &= \frac{1}{2} \sum_{ijab} \langle ij || ab \rangle t_i^a t_j^b \end{aligned}$$

The CCSD energy equation revisited

$$E_{CC} \Leftarrow \langle \Phi_0 | \frac{1}{2} \left(\hat{H}_N \hat{T}^2 \right)_c | \Phi_0 \rangle$$

$$\langle \Phi_0 | \frac{1}{2} \left(\hat{V}_N \hat{T}_1^2 \right)_c | \Phi_0 \rangle =$$

$$\frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \left(\left\{ a_p^\dagger a_q^\dagger a_s a_r \right\} \left\{ a_a^\dagger a_i \right\} \left\{ a_b^\dagger a_j \right\} \right)_c | \Phi_0 \rangle$$

$$= \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 |$$

$$\left(\left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} \right. \\ \left. + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} \right) | \Phi_0 \rangle$$

$$= \frac{1}{2} \sum_{ijab} \langle ij || ab \rangle t_i^a t_j^b$$

The CCSD energy equation revisited

$$E_{CC} \Leftarrow \langle \Phi_0 | \frac{1}{2} \left(\hat{H}_N \hat{T}^2 \right)_c | \Phi_0 \rangle$$

$$\begin{aligned} \langle \Phi_0 | \frac{1}{2} \left(\hat{V}_N \hat{T}_1^2 \right)_c | \Phi_0 \rangle &= \\ \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \left(\left\{ a_p^\dagger a_q^\dagger a_s a_r \right\} \left\{ a_a^\dagger a_i \right\} \left\{ a_b^\dagger a_j \right\} \right)_c | \Phi_0 \rangle \\ &= \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \\ &\quad \left(\left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} \right. \\ &\quad \left. + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j} \right\} \right) | \Phi_0 \rangle \\ &= \frac{1}{2} \sum_{ijab} \langle ij || ab \rangle t_i^a t_j^b \end{aligned}$$

The CCSD energy equation revisited

$$E_{CC} \Leftarrow \langle \Phi_0 | \frac{1}{2} \left(\hat{H}_N \hat{T}^2 \right)_c | \Phi_0 \rangle$$

$$\begin{aligned} \langle \Phi_0 | \frac{1}{2} \left(\hat{V}_N \hat{T}_1^2 \right)_c | \Phi_0 \rangle &= \\ \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \left(\left\{ a_p^\dagger a_q^\dagger a_s a_r \right\} \left\{ a_a^\dagger a_i \right\} \left\{ a_b^\dagger a_j \right\} \right)_c | \Phi_0 \rangle \\ &= \frac{1}{8} \sum_{pqrs} \sum_{ijab} \langle pq || rs \rangle t_i^a t_j^b \langle \Phi_0 | \\ &\quad \left(\left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{(pqrs)(ab)} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{(pqrs)(ab)} \right\} + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{(pqrs)(ab)} \right\} \right. \\ &\quad \left. + \left\{ \overbrace{a_p^\dagger a_q^\dagger a_s a_r a_a^\dagger a_i a_b^\dagger a_j}^{(pqrs)(ab)} \right\} \right) | \Phi_0 \rangle \\ &= \frac{1}{2} \sum_{ijab} \langle ij || ab \rangle t_i^a t_j^b \end{aligned}$$

The CCSD energy equation revisited

- ▶ No contractions possible between cluster operators.
- ▶ Cluster operators need to contract with free indices to the left.
- ▶ Disconnected parts automatically cancel in the commutator.
- ▶ Onebody operators can connect to maximum two cluster operators.
- ▶ Twobody operators can connect to maximum four cluster operators.
- ▶ Different terms in the expansion contributes to different equations.

The CCSD energy equation revisited

- ▶ No contractions possible between cluster operators.
- ▶ Cluster operators need to contract with free indices to the left.
- ▶ Disconnected parts automatically cancel in the commutator.
- ▶ Onebody operators can connect to maximum two cluster operators.
- ▶ Twobody operators can connect to maximum four cluster operators.
- ▶ Different terms in the expansion contributes to different equations.

The CCSD energy equation revisited

- ▶ No contractions possible between cluster operators.
- ▶ Cluster operators need to contract with free indices to the left.
- ▶ Disconnected parts automatically cancel in the commutator.
- ▶ Onebody operators can connect to maximum two cluster operators.
- ▶ Twobody operators can connect to maximum four cluster operators.
- ▶ Different terms in the expansion contributes to different equations.

The CCSD energy equation revisited

- ▶ No contractions possible between cluster operators.
- ▶ Cluster operators need to contract with free indices to the left.
- ▶ Disconnected parts automatically cancel in the commutator.
- ▶ Onebody operators can connect to maximum two cluster operators.
- ▶ Twobody operators can connect to maximum four cluster operators.
- ▶ Different terms in the expansion contributes to different equations.

The CCSD energy equation revisited


- ▶ No contractions possible between cluster operators.
- ▶ Cluster operators need to contract with free indices to the left.
- ▶ Disconnected parts automatically cancel in the commutator.
- ▶ Onebody operators can connect to maximum two cluster operators.
- ▶ Twobody operators can connect to maximum four cluster operators.
- ▶ Different terms in the expansion contributes to different equations.

The CCSD energy equation revisited

- ▶ No contractions possible between cluster operators.
- ▶ Cluster operators need to contract with free indices to the left.
- ▶ Disconnected parts automatically cancel in the commutator.
- ▶ Onebody operators can connect to maximum two cluster operators.
- ▶ Twobody operators can connect to maximum four cluster operators.
- ▶ Different terms in the expansion contributes to different equations.

Factoring, motivation

Diagram (2.12)



$$= \frac{1}{4} \langle mn || ef \rangle t_{ij}^{ef} t_{mn}^{ab}$$

Diagram (2.26)



$$= \frac{1}{4} P(ij) \langle mn || ef \rangle t_i^e t_{mn}^{ab} t_j^f$$


Diagram (2.31)



$$= \frac{1}{4} P(ij) P(ab) \langle mn || ef \rangle t_i^e t_m^a t_j^f t_n^b$$

Factoring, motivation


Diagram (2.12)



$$= \frac{1}{4} \langle mn || ef \rangle t_{ij}^{ef} t_{mn}^{ab}$$

Diagram cost: $n_p^4 n_h^4$

Diagram (2.13) - Factored



$$= \frac{1}{4} \langle mn || ef \rangle t_{ij}^{ef} t_{mn}^{ab}$$

$$= \frac{1}{4} \left(\langle mn || ef \rangle t_{ij}^{ef} \right) t_{mn}^{ab}$$

$$= \frac{1}{4} X_{ij}^{mn} t_{mn}^{ab}$$

Factoring, motivation

Diagram (2.26)



$$= \frac{1}{4} P(ij) \langle mn || ef \rangle t_i^e t_{mn}^{ab} t_j^f$$

Diagram cost: $n_p^4 n_h^4$

Diagram (2.26) - Factored




$$= \frac{1}{4} P(ij) \langle mn || ef \rangle t_i^e t_{mn}^{ab} t_j^f$$

$$= \frac{1}{4} P(ij) t_{mn}^{ab} t_i^e X_{ej}^{mn}$$

$$= \frac{1}{4} P(ij) t_{mn}^{ab} Y_{ij}^{mn}$$

Factoring, motivation


Diagram (2.31)



$$= \frac{1}{4} P(ij) P(ab) \langle mn || ef \rangle t_i^e t_m^a t_j^f t_n^b$$

Diagram cost: $n_p^4 n_h^4$

Diagram (2.31) - Factored




$$\begin{aligned}
 &= \frac{1}{4} P(ij) P(ab) \langle mn || ef \rangle t_i^e t_m^a t_j^f t_n^b \\
 &= \frac{1}{4} P(ij) P(ab) t_m^a t_n^b t_i^e X_{ej}^{mn} \\
 &= \frac{1}{4} P(ij) P(ab) t_m^a t_n^b Y_{ij}^{mn} \\
 &= \frac{1}{4} P(ij) P(ab) t_m^a Z_{ij}^{mb}
 \end{aligned}$$

Factoring, Classification

A diagram is classified by how many hole and particle lines between a \hat{T}_i operator and the interaction ($T_i(p^{np}h^{nh})$).

Diagram (2.12) Classification




$$= \frac{1}{4} \langle mn || ef \rangle t_{ij}^{ef} t_{mn}^{ab}$$

This diagram is classified as $T_2(p^2) \times T_2(h^2)$

Factoring, Classification


Diagram (2.26)



$$= \frac{1}{4} P(ij) \langle mn || ef \rangle t_i^e t_{mn}^{ab} t_j^f$$

This diagram is classified as $T_2(h^2) \times T_1(p) \times T_1(p)$

Diagram (2.31)



$$= \frac{1}{4} P(ij) P(ab) \langle mn || ef \rangle t_i^e t_m^a t_j^f t_n^b$$

This diagram is classified as $T_1(p) \times T_1(p) \times T_1(h) \times T_1(h)$

Factoring, Classification

Cost of making intermediates

Object	CPU cost	Memory cost
$T_2(h)$	$n_p^2 n_h$	n_p^2
$T_2(h^2)$	n_p^2	$n_h^{-2} n_p^2$
$T_2(p)$	$n_p n_h^2$	n_h^2
$T_2(ph)$	$n_p n_h$	1
$T_1(h)$	n_p	$n_h^{-1} n_p$
$T_2(ph^2)$	n_p	n_h^{-2}
$T_2(p^2)$	n_h^2	$n_p^{-2} n_h^2$
$T_1(p)$	n_h	$n_p^{-1} n_h$
$T_2(p^2 h)$	n_h	n_p^{-2}
$T_1(ph)$	1	$n_p^{-1} n_h^{-1}$

Factoring, Classification

Classification of \hat{T}_1 diagrams

Object	Expression id
$T_2(ph)$	5, 11
$T_1(h)$	3, 8, 10, 13, 14
$T_2(ph^2)$	7, 12
$T_1(p)$	2, 8, 9, 12, 14
$T_2(p^2h)$	6, 13
$T_1(ph)$	4, 9, 10, 11, 14

Factoring, Classification

Classification of \hat{T}_2 diagrams

Object	Expression id
$T_2(h)$	5, 15, 16, 23, 29
$T_2(h^2)$	7, 12, 22, 26
$T_2(p)$	4, 14, 17, 20, 30
$T_2(ph)$	8, 13, 13, 18, 21, 27
$T_1(h)$	3, 10, 10, 11, 17, 19, 21, 24, 25, 25, 27, 28, 28, 30, 31, 31
$T_2(ph^2)$	14
$T_2(p^2)$	6, 12, 19, 28
$T_1(p)$	2, 9, 9, 11, 16, 18, 22, 24, 24, 25, 26, 26, 27, 29, 31, 31
$T_2(p^2h)$	15
$T_1(ph)$	20, 23, 29, 30

Factoring, $T_2(h)$

Contribution to the \hat{T}_2 amplitude equation from $T_2(h)$

$$\begin{aligned} T_2(h) &\Leftarrow -P(ij)f_i^m t_{mj}^{ab} - \frac{1}{2}P(ij)\langle mn||ef\rangle t_{mi}^{ef} t_{nj}^{ab} - P(ij)f_e^m t_i^e t_{mj}^{ab} \\ &\quad - P(ij)\langle mn||ei\rangle t_m^e t_{nj}^{ab} - P(ij)\langle mn||ef\rangle t_m^e t_i^f t_{nj}^{ab} \\ &= -P(ij)t_{im}^{ab} \left[f_j^m + \langle mn||je\rangle t_n^e + \frac{1}{2}\langle mn||ef\rangle t_{jn}^{ef} \right. \\ &\quad \left. + t_j^e \left(f_e^m + \langle mn||ef\rangle t_n^f \right) \right] \\ &= -P(ij)t_{im}^{ab} (\bar{\text{H3}})_j^m \end{aligned}$$

Factoring, $T_2(h^2)$

Contribution to the \hat{T}_2 amplitude equation from $T_2(h^2)$

$$\begin{aligned} T_2(h^2) &\Leftarrow \frac{1}{2} \langle mn || ij \rangle t_{mn}^{ab} + \frac{1}{4} \langle mn || ef \rangle t_{ij}^{ef} t_{mn}^{ab} + \frac{1}{2} P(ij) \langle mn || ej \rangle t_i^e t_{mn}^{ab} \\ &\quad + \frac{1}{4} P(ij) \langle mn || ef \rangle t_i^e t_{mn}^{ab} t_j^f \\ &= \frac{1}{2} t_{mn}^{ab} \left[\langle mn || ij \rangle + \frac{1}{2} \langle mn || ef \rangle t_{ij}^{ef} \right. \\ &\quad \left. + P(ij) t_j^e \left(\langle mn || ie \rangle + \frac{1}{2} \langle mn || fe \rangle t_i^f \right) \right] \\ &= \frac{1}{2} t_{mn}^{ab} (\bar{H}9)_{ij}^{mn} \end{aligned}$$

Factored T_1 amplitude equations

$$0 = f_i^a + \langle ma || ei \rangle t_m^e + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + t_i^e (\text{I2a})_e^a - t_m^a (\bar{\text{H3}})_i^m \\ + \frac{1}{2} t_{mn}^{ea} (\bar{\text{H7}})_{ie}^{mn} + t_{im}^{ae} (\bar{\text{H1}})_e^m$$

Can be solved by

1. Matrix inversion for each iteration ($n_p^3 n_h^3$)
2. Extracting diagonal elements ($n_p^3 n_h^2$)

Factored T_1 amplitude equations

$$\begin{aligned}
 0 &= f_i^a + \langle ma || ei \rangle t_m^e + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + t_i^e (I2a)_e^a - t_m^a (\bar{H}3)_i^m \\
 &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m \\
 &= f_i^a + \langle ma || ei \rangle t_m^e + t_i^a (I2a)_a^a + (1 - \delta_{ea}) t_i^e (I2a)_e^a \\
 &\quad - t_i^a (\bar{H}3)_i^i - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} \\
 &\quad + t_{im}^{ae} (\bar{H}1)_e^m \\
 &= f_i^a + t_i^a \left((I2a)_a^a - (\bar{H}3)_i^i \right) + \langle ma || ei \rangle t_m^e \\
 &\quad + (1 - \delta_{ea}) t_i^e (I2a)_e^a - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} \\
 &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m
 \end{aligned}$$

Factored T_1 amplitude equations

$$\begin{aligned}
 0 &= f_i^a + \langle ma || ei \rangle t_m^e + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + t_i^e (I2a)_e^a - t_m^a (\bar{H}3)_i^m \\
 &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m \\
 &= f_i^a + \langle ma || ei \rangle t_m^e + t_i^a (I2a)_a^a + (1 - \delta_{ea}) t_i^e (I2a)_e^a \\
 &\quad - t_i^a (\bar{H}3)_i^i - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} \\
 &\quad + t_{im}^{ae} (\bar{H}1)_e^m \\
 &= f_i^a + t_i^a \left((I2a)_a^a - (\bar{H}3)_i^i \right) + \langle ma || ei \rangle t_m^e \\
 &\quad + (1 - \delta_{ea}) t_i^e (I2a)_e^a - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} \\
 &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m
 \end{aligned}$$

Factored T_1 amplitude equations

$$\begin{aligned}
 0 &= f_i^a + \langle ma || ei \rangle t_m^e + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + t_i^e (I2a)_e^a - t_m^a (\bar{H}3)_i^m \\
 &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m \\
 &= f_i^a + \langle ma || ei \rangle t_m^e + t_i^a (I2a)_a^a + (1 - \delta_{ea}) t_i^e (I2a)_e^a \\
 &\quad - t_i^a (\bar{H}3)_i^i - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} \\
 &\quad + t_{im}^{ae} (\bar{H}1)_e^m \\
 &= f_i^a + t_i^a \left((I2a)_a^a - (\bar{H}3)_i^i \right) + \langle ma || ei \rangle t_m^e \\
 &\quad + (1 - \delta_{ea}) t_i^e (I2a)_e^a - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} \\
 &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m
 \end{aligned}$$

Factored T_1 amplitude equations

Define

$$D_i^a = (\bar{H}3)_i^i - (I2a)_a^a,$$

and we get the T_1 amplitude equations

$$\begin{aligned} D_i^a t_i^a &= f_i^a + \langle ma || ei \rangle t_m^e + (1 - \delta_{ea}) t_i^e (I2a)_e^a \\ &\quad - (1 - \delta_{mi}) t_m^a (\bar{H}3)_i^m + \frac{1}{2} \langle am || ef \rangle t_{im}^{ef} \\ &\quad + \frac{1}{2} t_{mn}^{ea} (\bar{H}7)_{ie}^{mn} + t_{im}^{ae} (\bar{H}1)_e^m. \end{aligned}$$

Factored T_2 amplitude equations

$$\begin{aligned} 0 = & \langle ab || ij \rangle + \frac{1}{2} \langle ab || ef \rangle t_{ij}^{ef} - P(ij) t_{im}^{ab} (\bar{H}3)_j^m + \frac{1}{2} t_{mn}^{ab} (\bar{H}9)_{ij}^{mn} \\ & + P(ab) t_{ij}^{ae} (\bar{H}2)_e^b + P(ij) P(ab) t_{im}^{ae} (\text{I10c})_{ej}^{mb} - P(ab) t_m^a (\text{I12a})_{ij}^{mb} \\ & + P(ij) t_i^e (\text{I11a})_{ej}^{ab} \end{aligned}$$

Can be solved by

1. Matrix inversion for each iteration ($n_p^6 n_h^6$)
2. Extracting diagonal elements ($n_p^4 n_h^2$)

Factored T_2 amplitude equations

Similarly we define

$$D_{ij}^{ab} = (\bar{H}3)_i^j + (\bar{H}3)_j^i - (\bar{H}2)_a^a - (\bar{H}2)_b^b$$

and get the T_2 amplitude equations

$$\begin{aligned} D_{ij}^{ab} t_{ij}^{ab} = & \langle ab || ij \rangle + \frac{1}{2} \langle ab || ef \rangle t_{ij}^{ef} - P(ij)(1 - \delta_{jm}) t_{im}^{ab} (\bar{H}3)_j^m \\ & + \frac{1}{2} t_{mn}^{ab} (\bar{H}9)_{ij}^{mn} + P(ab)(1 - \delta_{be}) t_{ij}^{ae} (\bar{H}2)_e^b \\ & + P(ij)P(ab) t_{im}^{ae} (I10c)_{ej}^{mb} - P(ab) t_m^a (I12a)_{ij}^{mb} \\ & + P(ij) t_i^e (I11a)_{ej}^{ab} \end{aligned}$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

 Calculate intermediates

$$t_j^a \leftarrow \text{calculated value}$$

$$t_{ll}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow t_a^i t_j^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

 Calculate intermediates

$$t_i^a \leftarrow \text{calculated value}$$

$$t_{ij}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow t_a^i t_j^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

Calculate intermediates

$$t_i^a \leftarrow \text{calculated value}$$

$$t_{ij}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow f_a^i t_i^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

 Calculate intermediates

$$t_i^a \leftarrow \text{calculated value}$$

$$t_{ij}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow f_a^i t_i^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

 Calculate intermediates

$$t_i^a \leftarrow \text{calculated value}$$

$$t_{ij}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow f_a^i t_i^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

 Calculate intermediates

$$t_i^a \leftarrow \text{calculated value}$$

$$t_{ij}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow f_a^i t_i^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Setup modelspace

Calculate f and v amplitudes

$$t_i^a \leftarrow 0; t_{ij}^{ab} \leftarrow 0$$

$$E \leftarrow 1; E_{old} \leftarrow 0$$

$$E_{ref} \leftarrow \sum_i \langle i | \hat{t} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle$$

while not converged ($E - E_{old} > \epsilon$)

 Calculate intermediates

$$t_i^a \leftarrow \text{calculated value}$$

$$t_{ij}^{ab} \leftarrow \text{calculated value}$$

$$E_{old} \leftarrow E$$

$$E \leftarrow f_a^i t_i^a + \frac{1}{4} \langle ij || ab \rangle t_{ij}^{ab} + \frac{1}{2} \langle ij || ab \rangle t_i^a t_j^b$$

end while

$$E_{GS} \leftarrow E_{ref} + E$$

Coupled Cluster algorithm

Typical convergence of the T_2 amplitudes