

MD1 - CDL data analysis

February 21, 2021

Hypothesis

ManyDogs 1 aims to investigate how dogs perceive human pointing gestures. Our specific question is: Do dogs understand and act on human pointing as though it is a social communicative gesture? While studies of dogs following pointing are numerous, there is ongoing debate in the field of Canine Science about whether dogs perceive pointing gestures as peremptory and follow them much like they would a verbal command, or as socially communicative cues. We will address this question and investigate whether dogs understand and act on pointing as a social communicative gesture by comparing a condition in which pointing is prefaced by ostensive cues (intentional gaze, name calling verbal cues) to a condition in which it is not (gaze avoidance, no verbal cues).

Condition predictions

- i) We predict that if dogs perceive pointing gestures as commands, as a group they would follow the gesture above chance in both experimental conditions. Moreover, we would expect there to be no significant differences between point-following by dogs in the two experimental conditions.
- ii) We predict that if dogs perceive pointing gestures as socially communicative cues, they would perform significantly better in the ostensive condition than in the non-ostensive condition.
- iii) We predict that in the odor control task that dogs will not differ significantly from the hypothetical chance level of 0.5.

Analysis

We pre-registered the study design, procedure, predictions, and confirmatory analysis prior to data collection at the Open Science Framework (<https://osf.io/gz5pj/>). The data and analysis script is available in an associated online repository at https://github.com/ManyDogsProject/manydogs1_vienna.git. To evaluate whether the dogs performance deviated significantly from the hypothetical chance level of 0.5 in the ostensive, non-ostensive, and odor control condition we first aggregated the data across trials for each individual and condition. We then conducted one-sample t-tests to compare the performance against chance. To compare the performance between the test conditions (ostensive, non-ostensive) we fitted a Generalised Linear Mixed Model (GLMM) with binomial error distribution and logit link function using the function `glmer` of the `lme4` package [Bates.2015]. We included the predictor variables condition, order of condition (ostensive first, non-ostensive first), the trial number within condition, sex, age (in years), dogs' trainability score according to the c-BARQ questionnaire [Hsu.2003]. Additionally, we included the random intercept of subject ID and the random slopes of condition and trial number within subject ID. All covariates were centered and scaled to a standard deviation of 1. The random slope components of the factors were centered (to ensure that the results are not conditional on the choice of the reference category). Confidence intervals for the predictors are derived based on 1000 parametric bootstraps using a function kindly provided by Roger Mundry (based on the `bootMer` function of the package `lme4`). In order to check for collinearity issues, we determined Variance Inflation Factors (VIF) using the function `vif` (R package `car`, [Fox.2019]). Collinearity was no issue (maximum VIF: 1.02). To evaluate the model stability we dropped one level of subject ID random effect at a time and compared the model estimates of the resulting models (using a function kindly provided by Roger Mundry). This procedure revealed the model to be stable with respect to the fixed effects.

As inference criterion, we used p-values below .05 (two-tailed) for one sample t-tests. For the GLMM, we used likelihood ratio tests (R function drop1 with argument ‘test’ set to “Chisq”, @Barr.2013) with p-values smaller than .05 as criterion to make inferences about fixed effects.

Results

One-sample t-test to compare against chance level

The dogs ($N = 61$) performed significantly better than expected by chance in the ostensive condition (mean = 0.6, 95% CI [0.55, 0.65], $t(60)=4.41$, $p=0$, $BF=459.91$) but not in the non-ostensive condition (mean = 0.53, 95% CI [0.49, 0.57], $t(60)=1.47$, $p=0.146$, $BF=0.39$) or the odor control condition (mean = 0.46, 95% CI [0.41, 0.51], $t(60)=-1.45$, $p=0.151$, $BF=0.38$).

Generalised linear mixed model

The dogs were significantly more likely to choose the baited cup in the ostensive condition compared to the non-ostensive condition ($\chi^2(1)=5.107$, $p = 0.024$). The control predictors order of condition, trial number (within condition), sex, age, or the C-BARQ trainability score had no significant effect on dogs’ choices.

Table 1: Results of GLMM of the dogs’ choice performance in Experiment 1

	Estimate	SE	95% CI	χ^2	df	P
(Intercept)	0.122	0.128	-0.134 0.372			
Condition	0.298	0.132	0.034 0.575	5.107	1	0.024
Order of condition	0.057	0.133	-0.216 0.324	0.185	1	0.667
Trial number	-0.100	0.066	-0.227 0.032	2.322	1	0.128
Sex	-0.084	0.137	-0.361 0.184	0.371	1	0.543
Age	-0.012	0.067	-0.140 0.121	0.033	1	0.855
C-BARQ trainability score	0.095	0.067	-0.034 0.236	1.957	1	0.162

Notes: Reference categories: condition: non-ostensive condition; order of condition: non-ostensive condition first; sex: female; covariates trial number, age, and training experienced centered and scaled to a standard deviation of 1.