


## PAPER



Cite this: *Photochem. Photobiol. Sci.*, 2017, **16**, 1424

## The SOS Chromotest applied for screening plant antigenotoxic agents against ultraviolet radiation

J. L. Fuentes, <sup>a,b</sup> A. García Forero,<sup>a,b</sup> N. Quintero Ruiz,<sup>a,b</sup> C. A. Prada Medina,<sup>a</sup> N. Rey Castellanos,<sup>a</sup> D. A. Franco Niño,<sup>a</sup> D. A. Contreras García,<sup>a</sup> Y. Córdoba Campo<sup>b</sup> and E. E. Stashenko<sup>b</sup>

In this work, we investigated the usefulness of the SOS Chromotest for screening plant antigenotoxic agents against ultraviolet radiation (UV). Fifty Colombian plant extracts obtained by supercritical fluid (CO<sub>2</sub>) extraction, twelve plant extract constituents (apigenin, carvacrol,  $\beta$ -caryophyllene, 1,8-cineole, citral, *p*-cymene, geraniol, naringenin, pinocembrin, quercetin, squalene, and thymol) and five standard antioxidant and/or photoprotective agents (curcumin, epigallocatechin gallate, resveratrol,  $\alpha$ -tocopherol, and Trolox®) were evaluated for their genotoxicity and antigenotoxicity against UV using the SOS Chromotest. None of the plant extracts, constituents or agents were genotoxic in the SOS Chromotest at tested concentrations. Based on the minimal extract concentration that significantly inhibited UV-genotoxicity (CIG), five plant extracts were antigenotoxic against UV as follows: *Baccharis nitida* (16  $\mu\text{g mL}^{-1}$ ) = *Solanum crotonifolium* (16  $\mu\text{g mL}^{-1}$ ) > *Hyptis suaveolens* (31  $\mu\text{g mL}^{-1}$ ) = *Persea caerulea* (31  $\mu\text{g mL}^{-1}$ ) > *Lippia origanoides* (62  $\mu\text{g mL}^{-1}$ ). Based on CIG values, the flavonoid compounds showed the highest antigenotoxic potential as follows: apigenin (7  $\mu\text{M}$ ) > pinocembrin (15  $\mu\text{M}$ ) > quercetin (26  $\mu\text{M}$ ) > naringenin (38  $\mu\text{M}$ ) > epigallocatechin gallate (108  $\mu\text{M}$ ) > resveratrol (642  $\mu\text{M}$ ). UV-genotoxicity inhibition with epigallocatechin gallate, naringenin and resveratrol was related to its capability for inhibiting protein synthesis. A correlation analysis between compound antigenotoxicity estimates and antioxidant activity evaluated by the oxygen radical absorbance capacity (ORAC) assay showed that these activities were not related. The usefulness of the SOS Chromotest for bioprospecting of plant antigenotoxic agents against UV was discussed.

Received 23rd January 2017,  
Accepted 5th July 2017

DOI: 10.1039/c7pp00024c

rsc.li/pps

## Introduction

Skin exposure to solar radiation has been a major public concern because of its carcinogenicity.<sup>1–3</sup> UV induces DNA damage,<sup>4</sup> principally by producing the cyclobutane pyrimidine dimer (CPD) and (6–4) pyrimidine pyrimidone photoproducts (6–4PP), which cause mutations<sup>5</sup> that can lead to inflammatory responses, immunosuppression and skin cancer.<sup>6–8</sup> Therefore, preventive protection strategies using photoprotective agents are successful to reduce the harmful effect of solar radiation on healthy human populations.<sup>9</sup>

Phytochemicals are emerging as a class of genoprotective agents less toxic than synthetic photoprotective compounds.<sup>10</sup> Plants have metabolic pathways that naturally produce medicinal compounds;<sup>11</sup> therefore, they can be sources of antige-

toxic/antimutagenic compounds that prevent cellular mutations involved in skin cancer and aging by regulating UV-induced mutability.<sup>10,12,13</sup> These plant compounds operate by different mechanisms such as: absorbing photons directly, strengthening the endogenous antioxidant system, stimulating DNA repair mechanisms or melanogenesis, among others.<sup>14</sup>

The risk of UV exposure to human skin has been mainly assessed using empirical indices such as the minimum erythema dose (MED) and other UV indices based on erythema.<sup>15</sup> The index based on erythema has also been used to measure the ability of sunscreens to protect against UV. For example, the solar protection factor (SPF) indicates how much a sunscreen increases the capacity of natural skin defense against erythema or redness before burn. However, erythema is a risk indicator with the threshold UV dose of MED. The effect of UV doses lower than MED cannot be evaluated using erythema; whereas UV genotoxicity would be expected for the skin even at those low dose ranges.<sup>16</sup> In addition, compounds used as the active principle of sunscreens are selected mostly based on their absorbance UV spectra.<sup>17</sup> This criterion is not applicable to those phytochemicals that protect against UV by

<sup>a</sup>Laboratorio de Microbiología y Mutagénesis Ambiental, Grupo de Investigación en Microbiología y Genética, Escuela de Biología, Universidad Industrial de Santander (UIS), Bucaramanga, Colombia. E-mail: jfuentes@uis.edu.co

<sup>b</sup>Centro de Investigación en Biomoléculas, CIBIMOL, Centro de Investigación de Excelencia, CENIVAM, UIS, Bucaramanga, Colombia

stimulating cellular capability to repair DNA damage.<sup>12</sup> To overcome these limitations, the use of the DNA dosimeter has been proposed as a complement to improve the actual system that measures sunscreen genoprotective efficacy.<sup>18</sup>

*In vitro* bacterial assays are useful tools to evaluate the anti-genotoxic properties of natural compounds due to their sensitivity, simplicity and flexibility to design experimental procedures to study protective mechanisms.<sup>19</sup> The SOS Chromotest,<sup>20</sup> a colorimetric assay developed for carcinogen detection at Pasteur Institute (France), has been used to study antigenotoxic compounds against UV.<sup>21</sup> The assay exploits the inducible DNA replication–cell division coupling system (SOS response) present in *E. coli*<sup>22</sup> (Fig. 1). When *E. coli* cells are exposed to UV, CPD and 6-4PP are produced, which introduce structural distortions in DNA.<sup>23</sup> Since these photoproducts are an obstacle for chromosome replication,<sup>24</sup> they induce the SOS response.<sup>22</sup> Thus, the SOS Chromotest is highly sensitive to UV.<sup>25</sup>

In spite of the genetic differences between bacteria and human cells, there are several reasons that support the use of the SOS Chromotest to identify compounds with potential DNA protective activity in humans: (i) a DNA damage inducible system like bacterial SOS response that involves CPD repair, cell cycle arrest and increased antioxidant defenses and melanogenesis has been described in human skin cells.<sup>26</sup> Therefore, the predictive capability for DNA protection in human skin cells of the studied samples could be expected when the SOS Chromotest is used, because this assay is highly sensitive to UV-induced DNA damage.<sup>25</sup> (ii) It has been hypothesized that the suppression of SOS repair in *E. coli* comprises a possible antimutagenicity mechanism and protective effects of common vegetables.<sup>27</sup> (iii) Using UV-treated isogenic *E. coli* strains, we have previously identified critical genes for cell survival and SOS induction.<sup>28</sup> The simultaneous use of these

strains could provide preliminary information about the protection mechanisms by which natural compounds act against UV-induced genotoxicity. (iv) Plant extract cytotoxicity can produce false positive antigenotoxicity results with an *in vitro* assay.<sup>29</sup> The SOS Chromotest overcomes this limitation by measuring alkaline phosphatase activity in PQ37 cells, which reflects constitutive protein synthesis and bacteria cell density.<sup>30</sup> Therefore, samples that showed a UV-genotoxicity inhibitory effect related to their cytotoxic activity can be discarded during screening. (v) The SOS Chromotest is a validated genotoxicity assay<sup>31</sup> that compared to other methods allows greater throughput screening.<sup>32</sup>

We are developing a project aimed at identifying plants from Colombian flora as potential sources of compounds with genoprotective activity against UV. In this work, the usefulness of a modified protocol of the SOS Chromotest to detect the anti-genotoxic activity of phytochemical extracts against UV has been investigated. This is proposed as a reliable procedure for large-scale and rapid evaluation of plants that can be a source of genoprotective compounds against UV-induced genotoxicity.


## Material and methods

### Chemicals

Reference mutagen 4-nitroquinoline-1-oxide (4-NQO) and the standard compounds apigenin (95%), carvacrol (98%),  $\beta$ -caryophyllene (98%), citral (95%), 1,8-cineole (99%), curcumin (94%), *p*-cymene (99%), epigallocatechin gallate, EGCG (95%), fluorescein disodium (98%), geraniol (98%), naringenin (95%), pinocembrin (95%), quercetin (95%), resveratrol (99%), squalene (98%),  $\alpha$ -tocopherol (97%), thymol (99.5%), and Trolox® (97%) were obtained from Sigma Aldrich Co. (St Louis, Missouri, USA). 2,2'-Azobis(2-amidinopropane) dihydrochloride (AAPH) was purchased from Wako Chemicals USA, Inc. (Richmond, VA, USA). Luria-Bertani (LB) and the substrates for  $\beta$ -galactosidase (2-nitrophenyl- $\beta$ -D-galactopyranoside, ONPG) and alkaline phosphatase (4-nitrophenyl phosphate, PNPP) were purchased from Amresco (Solon, Ohio, USA). The other reagents were obtained from commercial houses J. T. Baker (Phillipsburg, NJ, USA) or Merck (Kenilworth, NJ, USA).

### Plant material and extracts

Fifty extracts were obtained from thirty-five Colombian plant species collected in different collection sites (Table 1), which were taxonomically identified by Dr José Luis Fernández Alonso at Colombian National Herbarium (National University, Bogotá, Colombia). Plant species collection during botanical expeditions was supported by the official collection permit granted by the Colombian Ministry of Housing and Environment (Agreement No. 101, Resolution No. 0812). For each specimen, fresh leaves and flowers were used to obtain the extract using supercritical fluid (CO<sub>2</sub>) extraction.<sup>33</sup> The extractions were developed at a pressure of 500 bar, while the



**Fig. 1** SOS Chromotest basis. The assay exploits the SOS response system in *E. coli* by fusion of a *lacZ* reporter gene to *sfiA* (*sulA*), a SOS gene responsible for temporarily inhibiting the cell division after DNA damage. Following DNA damage, the *lacZ* reporter gene is expressed when RecA binds to ssDNA at stalled replication forks, cleaving LexA enabling the transcription of the *lacZ* gene.  $\beta$ -Galactosidase, the gene product of *lacZ*, cleaves an indicator compound to produce a colored or fluorescent product that can be monitored using a plate reader. The quantity of colored/fluorescent product is directly proportional to the amount of damage induced.

**Table 1** List of the studied plants and collection sites is shown. For each plant extract, the minimal concentration that produces significant ( $p \leq 0.05$ ) constitutive alkaline phosphatase synthesis inhibition (CAPSI) and the minimal concentration that produces significant ( $p \leq 0.05$ ) genotoxicity inhibition (CGI) in PQ37 cells are shown. Only CAPSI and CGI values  $\leq 100 \mu\text{g mL}^{-1}$  were considered as biologically relevant cytotoxic and antigenotoxic concentrations, respectively.<sup>34</sup> The samples with relevant cytotoxic and antigenotoxic activities appear in bold letters

Family	Species name (CNH voucher)	Collection site	CAPSI ( $\mu\text{g mL}^{-1}$ )	CGI ( $\mu\text{g mL}^{-1}$ )
Asteraceae	<i>Achyrocline satureioides</i> (COL 579420)	Zapatoca (Santander)	1000 $\pm$ 0	125 $\pm$ 0
	<i>Ageratina popayanensis</i> (COL 579422)	Zapatoca (Santander)	250 $\pm$ 0	500 $\pm$ 0
	<i>Ambrosia peruviana</i> (COL 579246)	Zapatoca (Santander)	1000 $\pm$ 0	500 $\pm$ 0
	<i>Austroeupatorium inulifolium</i> (COL 559438)	Curití (Santander)	500 $\pm$ 0	500 $\pm$ 0
	<i>Baccharis nitida</i> (COL 559436)	Villanueva (Santander)	NI	<b>16 <math>\pm</math> 0</b>
	<i>Chromolaena peltia</i> (COL 559437)	Los Santos (Santander)	NI	125 $\pm$ 0
	<i>Lourteigia</i> ssp. (COL 559444)	Los Santos (Santander)	250 $\pm$ 0	250 $\pm$ 0
	<i>Tagetes caracasana</i> (COL 559441)	Zapatoca (Santander)	1000 $\pm$ 0	125 $\pm$ 0
	<i>Wedelia calycina</i> (COL 559439)	Barichara (Santander)	<b>4 <math>\pm</math> 0</b>	<b>16 <math>\pm</math> 0</b>
	<i>Wedelia calycina</i> (COL 578353)	Zapatoca (Santander)	1000 $\pm$ 0	250 $\pm$ 0
Burseraceae	<i>Bursera tomentosa</i> (COL 560236)	Los Santos (Santander)	<b>4 <math>\pm</math> 0</b>	NI
Capparaceae	<i>Capparis flexuosa</i> (COL 574225)-F1	Puerto Colombia (Atlántico)	NI	500 $\pm$ 0
Ericaceae	<i>Bejaria aestuans</i> (COL 560990)	Los Santos (Santander)	NI	1000 $\pm$ 0
Escalloniaceae	<i>Escallonia pendula</i> (COL 560238)	Curití (Santander)	<b>63 <math>\pm</math> 0</b>	1000 $\pm$ 0
Labiatae	<i>Hyptis brachiata</i> (COL 582531)-F1	Tame (Arauca)	NI	1000 $\pm$ 0
	<i>Hyptis brachiata</i> (COL 582531)-F2	Tame (Arauca)	500 $\pm$ 0	1000 $\pm$ 0
	<i>Hyptis sinuata</i> (COL 578965)-F1	Tame (Arauca)	NI	1000 $\pm$ 0
	<i>Hyptis sinuata</i> (COL 578965)-F2	Tame (Arauca)	NI	1000 $\pm$ 0
	<i>Hyptis suaveolens</i> (COL 560241)	Umpalá (Santander)	NI	<b>31 <math>\pm</math> 0</b>
	<i>Hyptis suaveolens</i> (COL 578964)-F1	Tame (Arauca)	1000 $\pm$ 0	250 $\pm$ 0
	<i>Minthostachys septentrionalis</i> (COL 560244)	Curití (Santander)	<b>4 <math>\pm</math> 0</b>	500 $\pm$ 0
	<i>Plectranthus amboinicus</i> (COL 560245)	Umpalá (Santander)	NI	500 $\pm$ 0
	<i>Salvia aratocensis</i> (COL 560246)	Los Santos (Santander)	NI	125 $\pm$ 0
	<i>Satureja</i> ssp. (COL 560973)	Jordán (Santander)	<b>8 <math>\pm</math> 0</b>	500 $\pm$ 0
Lauraceae	<i>Persea caerulea</i> (COL 560247)	Charalá (Santander)	250 $\pm$ 0	<b>31 <math>\pm</math> 0</b>
Malvaceae	<i>Sida aggregate</i> (COL 560249)	Los Santos (Santander)	NI	1000 $\pm$ 0
Myrtaceae	<i>Calycolpus</i> ssp. (COL 560976)	Los Santos (Santander)	250 $\pm$ 0	250 $\pm$ 0
	<i>Calycolpus moritzianus</i> (COL 578360)	Zapatoca (Santander)	1000 $\pm$ 0	250 $\pm$ 0
	<i>Eugenia puniceifolia</i> (COL 560250)	Los Santos (Santander)	<b>63 <math>\pm</math> 0</b>	1000 $\pm$ 0
	<i>Psidium sartorianum</i> (COL 578359)	Zapatoca (Santander)	NI	125 $\pm$ 0
Piperaceae	<i>Piper cumanense</i> (COL 578977)	Zapatoca (Santander)	1000 $\pm$ 0	250 $\pm$ 0
	<i>Piper dilatatifolium</i> (COL 578975)	Zapatoca (Santander)	250 $\pm$ 0	125 $\pm$ 0
	<i>Piper eriopodon</i> (COL 578364)	Zapatoca (Santander)	NI	125 $\pm$ 0
	<i>Piper eriopodon</i> (COL 578974)-F2	Tame (Arauca)	500 $\pm$ 0	125 $\pm$ 0
	<i>Piper subflavum</i> (COL 578976)	Zapatoca (Santander)	125 $\pm$ 0	500 $\pm$ 0
	<i>Piper subflavum</i> (COL 582361)-F2	Dagua (Valle del Cauca)	500 $\pm$ 0	1000 $\pm$ 0
	<i>Solanum crotonifolium</i> (COL 560253)	Umpalá (Santander)	NI	<b>16 <math>\pm</math> 0</b>
	<i>Turnera aff. diffusa</i> (COL 560255)	Los Santos (Santander)	NI	250 $\pm$ 0
Turneraceae	<i>Turnera diffusa</i> (COL 578361)	Girón (Santander)	250 $\pm$ 0	500 $\pm$ 0
	<i>Turnera diffusa</i> (COL 578361)-F1	Girón (Santander)	NI	125 $\pm$ 0
	<i>Turnera diffusa</i> (COL 578361)-F2	Girón (Santander)	1000 $\pm$ 0	500 $\pm$ 0
Verbenaceae	<i>Lippia americana</i> (COL 578970)	Zapatoca (Santander)	1000 $\pm$ 0	1000 $\pm$ 0
	<i>Lippia canescens</i> (COL 578969)	Zapatoca (Santander)	1000 $\pm$ 0	NI
	<i>Lippia micromera</i> (COL 578971)	Zapatoca (Santander)	NI	500 $\pm$ 0
	<i>Lippia origanoides</i> (COL 560267)	Umpalá (Santander)	125 $\pm$ 0	250 $\pm$ 0
	<i>Lippia origanoides</i> (COL 560259)	Los Santos (Santander)	NI	<b>62 <math>\pm</math> 0</b>
	<i>Lippia origanoides</i> (COL 560260)	Barichara (Santander)	NI	250 $\pm$ 0
	<i>Lippia origanoides</i> (COL 560980)	Los Santos (Santander)	<b>4 <math>\pm</math> 0</b>	250 $\pm$ 0
	<i>Lippia origanoides</i> (COL 582599)-F1	Tame (Arauca)	1000 $\pm$ 0	1000 $\pm$ 0
	<i>Lippia origanoides</i> (COL 582599)-F2	Tame (Arauca)	1000 $\pm$ 0	1000 $\pm$ 0

CNH: Colombian National Herbarium, NI: no inhibition.

collection of F1 and F2 fractions was done at pressures of 40 and 80 bar, respectively.

### Bacterial strains and culture

*Escherichia coli* PQ37 strain [ $F^-$  *thr leu his-4 pyrD thi galE galK or galT lacΔU169 srl300::Tn10 rpoB rpsL uvrA rfa trp::Muc<sup>+</sup> sfiA::Mud(Ap,lac)ts*], proposed for detecting genotoxic carcinogens,<sup>20</sup> was used. This strain carried the *suA::lacZ* gene fusion on the chromosome as a reporter of the primary DNA damage

induced during the SOS response. The cells were grown overnight at 37 °C and were shaken at 100 rpm in Luria-Bertani (LB) media (10 g tryptone per L, 5 g yeast extract per L, 10 g sodium chloride per L, pH 7.4), supplemented with ampicillin (50  $\mu\text{g mL}^{-1}$ ) and tetracycline (17  $\mu\text{g mL}^{-1}$ ).

### Cell irradiation

Cultures (1 mL) with an optical density ( $\text{OD}_{600}$ ) of 0.4 ( $\sim 2 \times 10^8$  cells per mL) were distributed into Petri plates with 5 cm-

diameter for irradiation during assays. For UVA (320–400 nm) and UVB (280–315 nm), the Petri plates with the cells were irradiated in darkness using a UVA/UVB irradiation chamber BS-02 equipped with a radiation controller UV-MAT (Opsytec Dr Groebel, Ettlingen, Baden-Württemberg, Germany). This radiation controller continuously measured the irradiance, calculated the irradiation dose and switched the lamps after reaching the target dose. The radiation dose used for the antigenotoxicity assay was  $10 \text{ J m}^{-2}$ , which induces adequately SOS functions in *E. coli* PQ37 cells without significantly affecting their survival.<sup>28</sup>

### Genotoxicity assay

Genotoxicity was evaluated using the SOS Chromotest as previously described.<sup>30</sup> Since organic solvents can affect genotoxicity and antigenotoxicity estimates in the SOS Chromotest,<sup>35</sup> we always used distilled water as the dissolvent and an ultrasonic bath E30H Elmasonic (Elma-Hans Schmidbauer GmbH & Co., Singen, Germany) to dissolve the extract and pure compounds. Overnight cultures were grown in LB medium to an optical density ( $\text{OD}_{600 \text{ nm}}$ ) of 0.4, diluted 10-fold in fresh LB medium and distributed (150  $\mu\text{L}$ ) into a series of Eppendorf tubes containing 5  $\mu\text{L}$  of the extract or compound to be tested. The bacteria were exposed for 30 min at 4 °C to different extract or compound concentrations, and then were cultured for 2 h at 37 °C with shaking at 300 rpm in a Termomixer apparatus (Eppendorf, Sao Paulo, Brazil). Negative (distilled water) and positive (2.34  $\mu\text{M}$  of 4-NQO) controls were always included in each assay. A minimum of four independent experiments per treatment with two replicates each were carried out.

$\beta$ -Galactosidase ( $\beta\text{G}$ ) and alkaline phosphatase (AP) activities were assayed in 96-well plates (Brand GMBH, Germany). For  $\beta\text{G}$  activity, cell membranes were disrupted by mixing 135  $\mu\text{L}$  of Z buffer (60 mM  $\text{Na}_2\text{HPO}_4$ , 40 mM  $\text{NaH}_2\text{PO}_4$ , 10 mM KCl, 1 mM  $\text{Mg}_2\text{SO}_4$ , 0.1% SDS, and 40 mM  $\beta$ -mercaptoethanol, pH 7.0) with 15  $\mu\text{L}$  of cell culture for 20 min at room temperature. The reaction was started by adding 30  $\mu\text{L}$  of ONPG (4 mg  $\text{mL}^{-1}$  in 0.1 M phosphate buffer, pH 7.0). After 40 min, the enzymatic reaction was stopped by adding 100  $\mu\text{L}$  of  $\text{Na}_2\text{CO}_3$  (1 M). For AP activity, cell membranes were disrupted by adding 135  $\mu\text{L}$  of T buffer (0.1% sodium dodecyl sulfate, 1 M Tris HCl, pH 8.8) to 15  $\mu\text{L}$  of the cell culture. The enzyme reaction was started by adding 30  $\mu\text{L}$  of PNPP solution (4 mg  $\text{mL}^{-1}$  in T buffer). After 40 min, the reaction was stopped by adding 50  $\mu\text{L}$  of HCl (2.5 M). After 5 min, 50  $\mu\text{L}$  of Tris (2 M) were added to restore the color. The final absorbances of the  $\beta\text{G}$  and AP assays were measured at  $\lambda = 420 \text{ nm}$ , using a microplate reader Multiskan GO (Thermo Scientific, MA, USA).  $\beta\text{G}$  and AP activities were calculated using the relationship: enzyme units =  $(1000 \times A_{420})/t$ , where  $A_{420}$  is the optical density at  $\lambda = 420 \text{ nm}$  and  $t$  is the length of incubation (min) with the substrate (ONPG or PNPP). The ratio of  $\beta\text{G}$  units to AP units ( $R = \beta\text{G}/\text{AP}$ ) reflects the induction of the *sulA* gene, even when there is some inhibition of protein synthesis. The genotoxicity criterion used was the SOS induction factor ( $I$ ) that represents

the normalized induction data of the *sulA* gene in each treatment (extract or compound) and was therefore considered to be an indirect measure of the primary DNA damage (genotoxicity) induced by the treatments. This parameter was calculated as  $I = R_t/R_{nt}$ , where  $t$  and  $nt$  were the treated and non-treated cells, respectively. The samples were classified as not genotoxic, if  $I$  was  $< 1.5$ , inconclusive, if  $I$  was between 1.5 and 2.0, and genotoxic, if  $I$  was  $> 2.0$ , and a clear concentration-response relationship was observed.

### Cytotoxicity assay

As we indicated above, plant extract cytotoxicity can produce false positive antigenotoxicity results with the SOS Chromotest.<sup>29</sup> In order to overcome this limitation, we considered the protein (AP) synthesis inhibition as a cytotoxicity criterion during screening. For each extract or compound, the concentration that produces significant ( $p \leq 0.05$ ) AP synthesis inhibition (CAPSI) in PQ37 cells was estimated. CAPSI values  $\leq 100 \mu\text{g mL}^{-1}$  for extracts and  $\leq 25 \mu\text{M}$  for compounds were considered as biologically relevant cytotoxic concentrations.<sup>34</sup> A minimum of four independent experiments per treatment with two replicates were carried out.

### Antigenotoxicity assay

Antigenotoxicity was assayed using a co-incubation procedure as previously described.<sup>36</sup> The cells were simultaneously treated with the extract or compound at different concentrations and  $10 \text{ J m}^{-2}$  of UVB, and then were cultured for 2 h at 37 °C with shaking at 300 rpm in a Termomixer apparatus (Eppendorf, Sao Paulo, Brazil). Negative (distilled water) and positive ( $10 \text{ J m}^{-2}$  of UVB) controls were always included in each assay. A minimum of four independent experiments per treatment with two replicates were carried out. The  $\beta\text{G}$  and AP activities were assayed as indicated above. Antigenotoxicity (the ability of the test extract or compound to protect against UV-induced genotoxicity) was measured as a significant reduction in the induction factor ( $I$ ) in the co-incubation procedure and expressed as a percentage of the genotoxicity inhibition:  $\% \text{GI} = 1 - (I_{ct} - I_{nt}/I_{UVB} - I_{nt}) \times 100$ , where  $I_{ct}$  is the SOS induction factor in the co-incubation procedure;  $I_{nt}$  is the SOS induction factor in non-treated cells, and  $I_{UVB}$  is the SOS induction factor in UVB-treated cells. Negative values of  $\% \text{GI}$  were considered as zero; therefore, this parameter ranged from a minimum of 0% to a maximum of 100%. The minimal concentration that produces significant ( $p \leq 0.05$ ) genotoxicity inhibition (CGI) in PQ37 cells was used for comparison of the genoprotective potential of the tested samples.

### Oxygen radical absorbance capacity (ORAC) assay

The ORAC assay was performed as previously described<sup>37,38</sup> with minor modifications. Pure compounds were prepared in potassium phosphate buffer (75 mM, pH 7.4), mixed with fluorescein prepared at 8.16  $\mu\text{M}$  in a 96-well microplate (Brand GmbH, Wertheim, Germany), and the mixture was incubated at 37 °C for 18 min. After that, an AAPH solution prepared at 153 mM in potassium phosphate buffer was added.



Fluorescence was measured on a Modulus™ II Microplate Multimode Reader (Turner Biosystems Inc., Sunnyvale, CA, USA), equipped with absorbance (UV-Vis) and fluorescence modules. An excitation band ( $\lambda = 490$  nm) and emission range  $\lambda$  from 510 to 579 nm were used for fluorescence measurement. All measurements were done in triplicate and their results were expressed as average  $\pm$  standard deviation. The ORAC values were calculated by plotting the Trolox® concentration ( $\mu\text{M}$ ) versus the remaining area under the fluorescence curve (AUC) for each compound. The remaining AUC was calculated subtracting the blank's AUC from the AUC of the sample. The results were expressed as micromoles of Trolox® per gram (g) ( $n = 3$ ).

### Statistical analysis

The  $\beta\text{G}$ , AP,  $I$ , %GI, CGI, CAPSI, and ORAC average values and the corresponding standard errors were calculated. The normality of the data was tested using the Kolmogorov–Smirnov test. Variance homogeneity and analysis of variance (ANOVA) tests were also conducted. Average values per treatment were compared with controls using a  $t$ -test for independent samples. Product–moment (Pearson) correlation analysis was used to examine the dose–response relationship in genotoxic and antigenotoxic studies and between the CGI and ORAC values. For all statistical analyses,  $p < 0.05$  was considered significant. The R program<sup>39</sup> was used for all analyses.

### Live subject statement

The project RC-0572-2012 was approved by the Operational Research and Extension Committee (Record No. 36-2011, File No. 1110) from UIS. The experiments and the chemical management were done according to the National law (Resolution No. 008430-1993) from the Ministry of Health of Colombia and Institutional Manual of Integrated Management and Processes (PGIR-PGGA.05).

## Results

### Induction of the *suIA* gene by UV

Table 2 shows the induction of the *suIA* gene in *E. coli* PQ37 following treatment with UVB (280–315 nm) and UVA (320–400 nm).

The extent of *suIA* gene induction by UV varied with the radiation type and the dose. A significant increase of  $I$  values was observed from doses 10 (UVB) and 5000 (UVA)  $\text{J m}^{-2}$ , respectively; although, smaller doses were not evaluated. UVB showed a biphasic induction of the *suIA* gene. Doses from 10 to 100  $\text{J m}^{-2}$  showed  $I$  values between 8.7 and 9.8 and then, this value increased abruptly to 21.6 at a dose of 280  $\text{J m}^{-2}$ . At this dose, a significant inhibitory effect on protein synthesis was detected. Finally, UVA reached the highest induction level ( $I = 10$ ) at a dose of 20.000  $\text{J m}^{-2}$  at the expense of a significant inhibition of protein synthesis.

**Table 2** Induction of the *suIA* gene in *E. coli* PQ37 after treatment with UVB and UVA. For each treatment, the  $\beta\text{G}$ , AP and  $I$  values were calculated from a minimum of four independent experiments with four replicates and the corresponding standard errors are given

UV doses ( $\text{J m}^{-2}$ )	SOS induction parameters		
	$\beta\text{G}$ value	AP value	$I$ values
<b>UVB</b>			
0.0	$2.7 \pm 0.2$	$2.9 \pm 0.3$	$1.0 \pm 0.0$
10.0	$22 \pm 1.6^*$	$2.8 \pm 0.1$ n.s.	$8.7 \pm 0.1^*$
20.0	$22 \pm 2.8^*$	$2.8 \pm 0.1$ n.s.	$8.4 \pm 0.6^*$
40.0	$24 \pm 2.8^*$	$2.7 \pm 0.1$ n.s.	$9.2 \pm 0.6^*$
50.0	$23 \pm 3.9^*$	$2.7 \pm 0.1$ n.s.	$9.0 \pm 0.7^*$
100.0	$23 \pm 1.3^*$	$2.6 \pm 0.1$ n.s.	$9.8 \pm 0.3^*$
180.0	$28 \pm 4.8^*$	$1.6 \pm 0.7^{**}$	$19 \pm 2.3^*$
280.0	$24 \pm 4.3^*$	$1.1 \pm 0.4^{**}$	$22 \pm 5.4^*$
390.0	$17 \pm 3.4^*$	$1.2 \pm 0.5^{**}$	$16 \pm 2.9^*$
500.0	$16 \pm 1.7^*$	$1.0 \pm 0.4^{**}$	$16 \pm 1.9^*$
<b>UVA</b>			
0.0	$2.4 \pm 0.3$	$1.6 \pm 0.3$	$1.0 \pm 0.0$
5000.0	$8.9 \pm 0.3^*$	$1.4 \pm 0.2$ n.s.	$4.0 \pm 0.5^*$
10 000.0	$11 \pm 0.5^*$	$1.3 \pm 0.1$ n.s.	$6 \pm 1.0^*$
15 000.0	$10 \pm 0.4^*$	$0.9 \pm 0.1^{**}$	$8 \pm 1.5^*$
20 000.0	$10 \pm 1.0^*$	$0.7 \pm 0.0^{**}$	$10 \pm 2.0^*$
40 000.0	$1.2 \pm 0.1$ n.s.	$0.2 \pm 0.0^{**}$	$4.1 \pm 0.8^*$
60 000.0	$1.2 \pm 0.0$ n.s.	$0.2 \pm 0.0^{**}$	$4.2 \pm 0.8^*$

\*Significant increase ( $p < 0.05$ ) with respect to zero was found using the  $t$ -test. \*\*Significant reduction ( $p < 0.05$ ) with respect to zero was found using the  $t$ -test. n.s. – no significant increase or reduction was found.

### Plant extract cytotoxicity, genotoxicity and antigenotoxicity against UV

Table 1 shows cytotoxicity estimates for each plant extract during antigenotoxicity screening. Based on CAPSI values, the plant extracts with relevant cytotoxic effects in *E. coli* (CAPSI values  $\leq 100 \mu\text{g mL}^{-1}$ )<sup>35</sup> were as follows: *W. calycina* COL559439 ( $4 \mu\text{g mL}^{-1}$ ) = *B. tomentosa* COL560236 ( $4 \mu\text{g mL}^{-1}$ ) = *M. septentrionalis* COL560244 ( $4 \mu\text{g mL}^{-1}$ ) = *C. moritzianus* COL560976 ( $4 \mu\text{g mL}^{-1}$ ) = *L. origanoides* COL560980 ( $4 \mu\text{g mL}^{-1}$ ) > *Satureja* ssp. COL560973 ( $8 \mu\text{g mL}^{-1}$ ) > *E. pendula* COL560238 ( $63 \mu\text{g mL}^{-1}$ ) = *E. puniceifolia* COL560250 ( $63 \mu\text{g mL}^{-1}$ ). The results suggest that these plants possess antimicrobial activity against *E. coli* cells.

The extract direct genotoxicity was assayed before its antigenotoxic effect was investigated. At tested doses, the extracts did not increase the  $I$  values in PQ37 *E. coli* strain; therefore, they were considered not genotoxic in *E. coli* cells (data not shown).

Table 1 shows the CGI values for each plant extract. Based on CGI values, the plant extracts with relevant antigenotoxic effects (CGI values  $\leq 100 \mu\text{g mL}^{-1}$ ) were as follows: *B. nitida* COL559436 ( $16 \mu\text{g mL}^{-1}$ ) = *S. crotonifolium* COL560253 ( $16 \mu\text{g mL}^{-1}$ ) = *W. calycina* COL559439 ( $16 \mu\text{g mL}^{-1}$ ) > *H. suaveolens* COL560241 ( $31 \mu\text{g mL}^{-1}$ ) = *P. caerulea* COL560247 ( $31 \mu\text{g mL}^{-1}$ ) > *L. origanoides* COL560259 ( $62 \mu\text{g mL}^{-1}$ ). The *W. calycina* (COL559439) extract showed CAPSI  $\leq$  CGI values, which suggested that its antigenotoxicity was possibly associated with the inhibition of protein synthesis. Thus, *B. nitida*

COL559436, *S. crotonifolium* COL560253, *H. suaveolens* COL560241, *P. caerulea* COL560247 and *L. origanoides* COL560259 are not cytotoxic extracts that significantly reduce UV-induced genotoxicity depending on the concentration (Fig. 2).




Fig. 2 Antigenotoxic effect of promissory plant extracts against UV expressed as %GI. Values above the dotted line indicate significant inhibition of UV-genotoxicity. Error bars indicate the standard error of the mean computed from four independent experiments ( $n = 4$ ).

### Plant constituent cytotoxicity, genotoxicity and antigenotoxicity against UV

Table 3 shows cytotoxicity estimates for plant constituents and standard compounds during antigenotoxicity screening.

Based on CAPSI values of each constituent and standard compounds, the relative inhibitory effect was as follows: naringenin ( $38 \mu\text{M}$ ) > pinocembrin ( $62 \mu\text{M}$ ) > resveratrol ( $80 \mu\text{M}$ ) > EGCG ( $108 \mu\text{M}$ ) > thymol ( $523 \mu\text{M}$ ) > geraniol ( $22\,454 \mu\text{M}$ ) > carvacrol ( $25\,414 \mu\text{M}$ ) > citral ( $47\,625 \mu\text{M}$ ). The remaining compounds did not inhibit protein synthesis. Since only com-

pounds with CAPSI values  $\leq 25 \mu\text{M}$  are considered as relevantly cytotoxic;<sup>35</sup> none of the compounds studied was cytotoxic in the SOS Chromotest.

The genotoxicity of plant constituents was also assayed before their antigenotoxic effect had been investigated. At tested doses, the compounds did not increase *I* values in PQ37 *E. coli* strain; therefore, these were considered as not genotoxic in *E. coli* cells (data not shown).

Table 3 also shows the CGI values estimated for each plant constituent and standard compound. In general, flavonoids showed CGI values lower than terpene compounds.

Based on CGI values, the relative antigenotoxic potential of compounds was as follows: apigenin ( $7 \mu\text{M}$ ) > pinocembrin ( $15 \mu\text{M}$ ) > quercetin ( $26 \mu\text{M}$ ) > curcumin ( $33 \mu\text{M}$ ) > naringenin ( $38 \mu\text{M}$ ) > EGCG ( $108 \mu\text{M}$ ) > resveratrol ( $642 \mu\text{M}$ ) >  $\beta$ -caryophyllene ( $1911 \mu\text{M}$ ) > Trolox® ( $3130 \mu\text{M}$ ) > thymol ( $8373 \mu\text{M}$ ) > citral ( $23\,813 \mu\text{M}$ ) > carvacrol ( $25\,414 \mu\text{M}$ ) > geraniol ( $89\,816 \mu\text{M}$ ) > 1,8-cineol ( $94\,679 \mu\text{M}$ ) > squalene ( $102\,925 \mu\text{M}$ ) > *p*-cymene ( $104\,151 \mu\text{M}$ ). The compounds EGCG, naringenin and resveratrol showed CAPSI values lower than or equal to the CGI values, suggesting that its antigenotoxicity was possibly associated with the inhibitory effect of protein synthesis. Considering CGI values, the compounds with the highest antigenotoxic potential were apigenin, pinocembrin, and quercetin (CGI values  $\leq 25 \mu\text{M}$ ).

### Plant constituent antioxidant activity

In Table 3, the antioxidant activity of plant constituents is also shown. Based on ORAC values ( $\mu\text{mol Trolox}^\circ$  per g substance) of each constituent and standard compound, the relative antioxidant potential was as follows: resveratrol ( $35\,300 \pm 150$ ) > quercetin ( $34\,000 \pm 1531$ ) > naringenin ( $27\,600 \pm 429$ ) > apigenin ( $24\,500 \pm 615$ ) > pinocembrin ( $20\,500 \pm 128$ ) > EGCG ( $13\,200 \pm 531$ ) > geraniol ( $12\,300 \pm 119$ ) > carvacrol ( $3410 \pm 50$ ) > thymol ( $2900 \pm 103$ ) = citral ( $2900 \pm 103$ ) >  $\beta$ -caryophyllene ( $2800 \pm 109$ ) >  $\alpha$ -tocopherol ( $550 \pm 13$ ) > 1,8-cineol ( $299 \pm 5$ ) > *p*-cymene ( $219 \pm 2$ ). Product-moment correlation analysis using CGI and ORAC data from these compounds showed moderate and negative correlation ( $R = -0.55$ ,  $p < 0.009$ ), suggesting that the mechanisms by which these compounds exert their antigenotoxic and antioxidant activities are possibly different and unconnected.

## Discussion

We showed that the SOS Chromotest was highly sensitive to UV-induced DNA damage. The dose detection limits depended on the radiation type as follows: UVB ( $10 \text{ J m}^{-2}$ ) > UVA ( $5000 \text{ J m}^{-2}$ ). Previous works showed the differences in dose detection limits and they depend on the UV dosimeter and radiation type used. Using the SOS Chromotest, Quillardet and Hofnung<sup>25</sup> showed significant SOS induction with  $0.02 \text{ J m}^{-2}$  of UVC. Differences found in this study could be explained based on the different energies of UVC, UVB and UVA radiation. Using a DNA dosimeter,<sup>40</sup> detection of UVB-induced

Table 3 Promissory plant constituents and their CAPSI and CGI values in PQ37 cells

Plant constituents	CAPSI ( $\mu\text{M}$ )	CGI ( $\mu\text{M}$ )	ORAC values ( $\mu\text{mol Trolox}^\circ$ per g substance)
Apigenin	NI	$7 \pm 0$	$24\,500 \pm 615$
Carvacrol	$25\,414 \pm 0$	$25\,414 \pm 0$	$3410 \pm 50$
$\beta$ -Caryophyllene	NI	$1911 \pm 0$	$2800 \pm 109$
1,8-Cineol	NI	$94\,679 \pm 0$	$299 \pm 5$
Citral	$47\,625 \pm 0$	$23\,813 \pm 0$	$2900 \pm 103$
Curcumin	NI	$33 \pm 0$	$4350 \pm 77$
<i>p</i> -Cymene	$6509 \pm 0$	$104\,151 \pm 0$	$219 \pm 2$
EGCG	$108 \pm 0$	$108 \pm 0$	$13\,200 \pm 531$
Geraniol	$22\,454 \pm 0$	$89\,816 \pm 0$	$12\,300 \pm 119$
Naringenin	$38 \pm 0$	$38 \pm 0$	$27\,600 \pm 429$
Pinocembrin	$62 \pm 0$	$15 \pm 0$	$20\,500 \pm 128$
Quercetin	NI	$26 \pm 0$	$34\,000 \pm 1531$
Resveratrol	$80 \pm 0$	$642 \pm 0$	$35\,300 \pm 150$
Squalene	NI	$3220 \pm 0$	$0 \pm 0$
Thymol	$523 \pm 0$	$8373 \pm 0$	$2900 \pm 103$
$\alpha$ -Tocopherol	NI	NI	$550 \pm 13$
Trolox®	NI	$3130 \pm 0$	—

NI: no significant inhibition.

photoproducts occurs at a dose of  $400 \text{ J m}^{-2}$ ; while the Comet assay detected UVC- and UVA-induced DNA damage at doses of 1000 and  $50\,000 \text{ J m}^{-2}$ , respectively.<sup>41,42</sup> Recently, Schuch and Menck<sup>43</sup> showed that a DNA dosimeter system using bacteriophage T4 endonuclease V was sensitive to the detection of CPDs at doses of  $50 \text{ J m}^{-2}$  (UVC),  $2000 \text{ J m}^{-2}$  (UVB) and  $50\,000 \text{ J m}^{-2}$  (UVA). All these results indicate that the SOS Chromotest stands among the most sensitive UV biological dosimeters currently known.

In this work, we showed that a proper use of the SOS Chromotest can provide useful information on the cytotoxic and antigenotoxic activities of plant extracts and their constituents. As we showed, extracts with high antigenotoxic activity were not cytotoxic with the only exception of the *W. calycina* COL 559439 extract. In this way, our study supports the use of the SOS Chromotest for antigenotoxic compound screening that might be used as a supplement in sunscreen formulation.

As indicated in the Introduction section, compounds used as the active principle of sunscreens are selected mainly based on their UV absorbance capability to be used as solar filters.<sup>17</sup> Sunscreen efficacy has been extensively evaluated using *in vivo* measuring MED indices.<sup>15</sup> However, UV genotoxicity can occur at UV doses lower than MED.<sup>16</sup> Thus, cell and DNA dosimeter assays have been proposed as an alternative to overcome limitations of sunscreen efficacy testing on humans.<sup>18,44</sup> The SOS Chromotest exploits the inducible DNA replication–cell division coupling system (SOS response) present in *E. coli*. Since photoproducts such as CPD and 6-4PP are an obstacle for chromosome replication, detection of antigenotoxic effects with this assay could be linked to the inhibition of CPD or 6-4PP formation, photoproducts DNA repair, and cell division restoration events. These modes of action have been described for several photoprotective agents (Table 4). Since PQ37 strain (*uvrA*) is defective for nucleotide excision repair and the *sulA* gene in this strain was not turned off after UV treatment,<sup>28</sup> we believe that the antigenotoxicity of extracts detected with PQ37 strain was linked to the inhibition of CPD or 6-4PP formation due to their UV absorbance capability. However, the stimulation of the recombination pathway, which is also involved in CPD repair,<sup>28</sup> cannot be discarded as a possible DNA protection mechanism of these plant extracts. The extracts or compounds with such properties could be a useful supplement in sunscreen formulation, because they can prevent mutations that lead to skin cancer.

Although we proposed the SOS Chromotest for bioprospecting of plant genoprotective agents, the simultaneous use of isogenic strains with defective DNA repair functions and wild types with this assay could provide preliminary information about the protection mechanisms by which compounds act against UV-induced genotoxicity as previously showed.<sup>54,58,61</sup> The SOS Chromotest can also overcome some limitations of the cell-based assays. For example, many plant compounds do not penetrate through cytoplasmic membranes in cell-based assays.<sup>29</sup> To overcome this limitation, the SOS Chromotest contains a *rfa* mutation that alters outer membrane lipopolysaccharides allowing better diffusion of compounds into the

cell.<sup>30</sup> In addition, plant extract cytotoxicity can produce false positive antigenotoxicity results with an *in vitro* assay.<sup>29</sup> The SOS Chromotest overcomes this limitation measuring AP activity in PQ37 cells, which reflects constitutive protein synthesis and bacteria cell density.<sup>30</sup> Thus, samples that showed the UV-genotoxicity inhibitory effect related to their cytotoxic activity can be discarded during screening. Here, the measurements of AP activity inhibition provide additional and preliminary information about the antimicrobial potential of plant extracts against *E. coli* cells.

However, some critical points must be taken into account when using the SOS Chromotest for antigenotoxicity studies. Secondary interactions between the solvent used to dissolve plant extracts or compounds may result in synergistic effects in the cells; this interference can cause antigenotoxicity overestimates with the SOS chromotest.<sup>34</sup> The relevance in humans of the tested concentrations with *in vitro* cell-based assay is little known. Testing antimicrobial activity, Cos *et al.*<sup>34</sup> indicated  $\text{IC}_{50}$  values below  $100 \mu\text{g mL}^{-1}$  as biologically relevant for extracts and below  $25 \mu\text{M}$  for pure compounds. However, the relevance of this criterion when the cell-based bioassays use different endpoints to measure bioactivities is unknown. Since the SOS Chromotest is an *in vitro* cell-based assay, additional studies using skin human cell and animal models should be addressed to clarify the protection mode of action of extract constituents and to support their use for sunscreen formulation.

The present work evidenced that Colombia flora contains plants useful as a source of antigenotoxic compounds against UV. We showed for the first time that extracts from *B. nitida*, *S. crotonifolium*, *H. suaveolens*, *P. caerulea*, and *L. origanoides* have high genoprotective potential against UV. A preliminary chemical characterization study of these extracts using gas chromatography–mass spectrometry (data not shown) showed the major constituents (relative amount higher than 10%) as follows: *B. nitida* (17%  $\alpha$ -eudesmol, 56% non-identified flavonoid compounds), *H. suaveolens* (14% fenchone, 13% caryophyllene oxide, 52% non-identified flavonoid compounds), *P. caerulea* (13%  $\alpha$ -copaene, 14%  $\beta$ -caryophyllene, 33% non-identified flavonoid compounds) and *L. origanoides* (11%  $\beta$ -caryophyllene, 55% pinocembrin). In addition, using high performance liquid chromatography (HPLC) coupled with a diode-array detector (DAD), Stashenko *et al.*<sup>33</sup> identified as the major constituents of *Lippia* extracts the flavonoid compounds apigenin, naringenin, pinocembrin and quercetin; the flavonoids naringenin and pinocembrin being the major constituents in *L. origanoides* extract. We evaluated several of these plant constituents as pure compounds and these resulted in antigenotoxicity against UV (Table 3). Interestingly, the antigenotoxicity of these compounds was not directly co-related to their antioxidant activity, which suggests different mechanisms for these two bioactivity types. Our results also suggested that flavonoids were better antigenotoxic compounds than terpenes, showing always the lowest CGI values among all the compounds studied.

We have recently revealed that the antigenotoxic effects against UV of the essential oils from *Lippia* species depend on

**Table 4** Some plant extracts or constituents with the reported antigenotoxic effect against UV-induced DNA damage

Plan extracts (constituents)	Models (assays used)	Effect detected [sources]
<i>Araucaria angustifolia</i> (biflavonoid fraction)	Calf thymus DNA (8-OHdG and CPD HPLC detection)	Inhibition of 8-OHdG and CPD formation <sup>45</sup>
<i>Artemisia herba alba</i>	<i>S. cerevisiae</i> (reversion and gene conversion assays)	Inhibition of UV mutagenesis <sup>46</sup>
<i>Camellia sinensis</i> (EGCG polyphenols)	Human skin cell DNA ( <sup>32</sup> P postlabeling technique)	Inhibition of UV-induced DNA damage (CPD) and stimulate nucleotide excision repair involving interleukin (IL)-12 <sup>13,47–52</sup>
<i>Bixa orellana</i> (norbixin)	Mice (CPD immunodetection)	
<i>Cinnamomum camphora</i>	Human skin cells (CPD immunodetection)	
<i>Colobanthis quitensis</i> (flavonoid and carotenoids)	<i>E. coli</i> (SOS Chromotest)	Inhibition of SOS functions <sup>53</sup>
	<i>S. cerevisiae</i> (reversion and gene conversion assays)	Inhibition of UV mutagenesis <sup>46</sup>
	<i>S. cerevisiae</i> (reversion assay)	Prevent CPD formation and stimulate DNA repair processes <sup>54</sup>
<i>Deschampsia antarctica</i> (flavonoid and carotenoids)	Hamster V79 fibroblasts cells (comet assay)	
	<i>H. aspersas</i> (comet assay)	
	<i>S. cerevisiae</i> (reversion assay)	Prevent CPD formation and stimulate DNA repair processes <sup>54</sup>
<i>Fragaria ananassa</i> (anthocyanins)	Hamster fibroblasts (comet assay)	
<i>Jatropha curcas</i>	<i>H. aspersas</i> (comet assay)	
<i>Lycopodium clavatum</i> (apigenin)	Human fibroblasts (comet assay)	Inhibition of UV-induced DNA damage <sup>55</sup>
	Human lymphocytes (comet assay)	
	Human keratinocytes (comet assay)	Inhibition of UV-induced DNA damage <sup>56</sup>
	Mice (chromosome aberrations and micronuclei assays)	Inhibition of UV-induced DNA damage and stimulate nucleotide excision repair <sup>57</sup>
	Mice skin cells (DNA repair qPCR expression analysis)	
<i>Origanum compactum</i>	<i>S. cerevisiae</i> (reversion and gene conversion assays)	Inhibition of UV mutagenesis <sup>46</sup>
<i>Phyllanthus orbicularis</i> (polyphenols)	(flow cytometry analysis)	Remove UV-induced CPD modulating DNA repair <sup>58</sup>
<i>Polytrichum juniperinum</i> (flavonoids and carotenoids)	Human skin cells (CPD immunodetection)	
	<i>S. cerevisiae</i> (reversion assay)	Prevent CPD formation and stimulate DNA repair processes <sup>54</sup>
Romanian propolis ( <i>p</i> -cumaric acid, ferulic acid)	Hamster fibroblasts (comet assay)	
<i>Prunella vulgaris</i> (rosmarinic acid)	<i>H. aspersas</i> (comet assay)	
<i>Salvia officinalis</i> (EO, thujone, cineole, camphor)	Mice (CPD immunodetection)	Inhibition of UV-induced CPD formation <sup>59</sup>
<i>Silybum marianum</i> (silymarin)	Human keratinocytes (comet assay)	Inhibition of UV-induced DNA damage <sup>60</sup>
	<i>E. coli</i> (reversion assay)	Inhibition of UV mutagenesis <sup>61–63</sup>
	<i>S. typhimurium</i> (reversion assay)	
	<i>S. cerevisiae</i> (reversion assay)	
	Human keratinocytes and fibroblasts (comet assay, CPD immunodetection, DNA repair qPCR expression analysis)	Inhibition of UV-induced CPD formation and stimulate DNA repair processes <sup>64</sup>
<i>Undaria crenata</i>	Human keratinocytes (comet assay)	Decrease UV-induced DNA fragmentation <sup>65</sup>
<i>Vitis vinifera</i> (proanthocyanidins)	Human fibroblasts (CPD immunodetection)	Inhibition of UV-induced CPD and stimulate nucleotide excision repair involving interleukin (IL)-12 <sup>66</sup>

CPD: cyclobutane pyrimidine dimer. 8-OHdG: 8-hydroxy-20-deoxyguanosine. HPLC: high performance liquid chromatography.

specimen chemotypes,<sup>67</sup> indicating that chemical characterization is critical for studying plant extract antigenotoxicity. Therefore, broad chemical characterization of promissory plant extracts from *B. nitida*, *S. crotonifolium*, *H. suaveolens* and *P. caerulea* is necessary to identify the major constituents and evaluate their antigenotoxicity against UV to support this hypothesis.

At least one of these promissory plant species, *i.e.*, *L. origanoides* (COL560259), have shown very broad-ranging therapeutic potential<sup>68</sup> and its extracts have been well characterized exhibiting pinocembrin and naringenin as the major compounds.<sup>33</sup> The genoprotective properties against UV (present work) widen its potential as a source of compounds

(*i.e.*, pinocembrin and naringenin) with possible application in photoprotection and skin cancer chemoprevention. In addition, we showed several plant extracts that inhibited constitutive AP synthesis (*i.e.*, *W. calycina*, *B. tomentosa*, *M. septentrionalis*, *C. moritzianus*, *L. origanoides*, *Satureja* ssp., *E. pendula*, and *E. punicifolia*). The antimicrobial potential of these plants against *E. coli* is under investigation.

## Concluding remarks

Our results demonstrated that the SOS Chromotest was highly sensitive to UV-induced DNA damage. This assay provided



useful information about the cytotoxic and antigenotoxic effects of plant extracts and their constituents. Therefore, this assay can be used for bio-prospecting of plant antigenotoxic agents. The extracts from *B. nitida*, *S. crotonifolium*, *H. suaveolens*, *P. caerulea*, and *L. origanoides* have antigenotoxic properties against UV. The *L. origanoides* (COL560259) extract antigenotoxicity was linked to its flavonoid constituents, pino-cembrin and naringenin.

## Acknowledgements

The authors thank Montserrat Llagostera Casal from Universidad Autónoma de Barcelona for the PQ37 *E. coli* strain supply. This work was supported by the “Vicerrectoría de Investigaciones y Extensión de la Universidad Industrial de Santander”, and by the “Patrimonio Autónomo Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación, Francisco José de Caldas” (Grant No. RC-0572-2012). The “Ministerio de Ambiente y Desarrollo Sostenible (MADS)” of Colombia supported the present project through access permits to genetic resources and derivatives for scientific research (Agreement No. 101, Resolution No. 0812).

## References

- 1 S. Gandini, F. Sera, M. S. Cattaruzza, P. Pasquini, O. Picconi, P. Boyle and S. F. Melchi, Meta-analysis of risk factors for cutaneous melanoma II: Sun exposure, *Eur. J. Cancer*, 2005, **41**, 45–60.
- 2 M. P. Purdue, L. E. Freeman, W. F. Anderson and M. A. Tucker, Recent trends in incidence of cutaneous melanoma among US Caucasian young adults, *J. Invest. Dermatol.*, 2008, **128**, 2905–2980.
- 3 E. Linos, S. M. Swetter, M. G. Cockburn, G. A. Colditz and C. A. Clarke, Increasing burden of melanoma in the United States, *J. Invest. Dermatol.*, 2009, **129**, 1666–1674.
- 4 G. P. Pfeifer, Formation and Processing of UV Photoproducts: Effects of DNA Sequence and Chromatin Environment, *Photochem. Photobiol.*, 1997, **65**, 270–283.
- 5 H. Ikehata and T. Ono, The mechanisms of UV mutagenesis, *J. Radiat. Res.*, 2011, **52**, 115–125.
- 6 A. A. Vink and L. Roza, Biological consequences of cyclobutane pyrimidine dimers, *J. Photochem. Photobiol. B*, 2001, **65**, 101–104.
- 7 N. M. Wikonkal and E. B. Douglas, Ultraviolet Radiation Induced Signature Mutations in Photocarcinogenesis, *J. Invest. Dermatol. Symp. Proc.*, 1999, **4**, 6–10.
- 8 M. R. Zaidi, C. P. Day and G. Merlino, From UVs to metastases: modeling melanoma initiation and progress in the mouse, *J. Invest. Dermatol.*, 2008, **128**, 2381–2391.
- 9 Y. Gilaberte and S. González, Update on photoprotection, *Actas Dermosifiliogr.*, 2010, **101**, 659–672.
- 10 F. Afaq, Natural agents: Cellular and molecular mechanisms of photoprotection, *Arch. Biochem. Biophys.*, 2011, **508**, 144–151.
- 11 W. J. Zhang and L. O. Björn, The effect of ultraviolet radiation on the accumulation of medicinal compounds in plants, *Fitoterapia*, 2009, **80**, 207–218.
- 12 J. A. Nichols and S. K. Katiyar, Skin photoprotection by natural polyphenols: anti-inflammatory, antioxidant and DNA repair mechanisms, *Arch. Dermatol. Res.*, 2010, **302**, 71–83.
- 13 F. Afaq and S. K. Katiyar, Polyphenols: Skin Photoprotection and Inhibition of Photocarcinogenesis, *Mini-Rev. Med. Chem.*, 2011, **11**, 1200–1215.
- 14 R. Jansen, S. Q. Wang, M. Burnett, U. Osterwalder and H. W. Lim, Photoprotection: Part I. Photoprotection by naturally occurring, physical, and systemic agents, *J. Am. Acad. Dermatol.*, 2013, **69**, 853.e1–12.
- 15 J. Parrish, K. F. Jaenicke and R. R. Anderson, Erythema and melanogenesis action spectra of normal human skin, *Photochem. Photobiol.*, 1982, **36**, 187–191.
- 16 H. Ikehata, N. Munakata and T. Ono, Skin can control solar UVR-induced mutations through the epidermis-specific response of mutation induction suppression, *Photochem. Photobiol. Sci.*, 2013, **12**, 2008–2015.
- 17 J. Stanfield, U. Osterwalder and B. Herzog, *In vitro* measurements of sunscreen protection, *Photochem. Photobiol. Sci.*, 2010, **9**, 489–494.
- 18 A. P. Schuch, J. Carvalhães-Lago, T. Yagura and C. F. M. Menck, DNA Dosimetry assessment for sunscreen genotoxic photoprotection, *PLoS One*, 2012, **7**, e40344.
- 19 S. De Flora, A. Camoirano, F. D'Agostini and R. Balansky, Modulation of the mutagenic response in prokaryotes, *Mutat. Res.*, 1992, **267**, 183–192.
- 20 P. Quillardet, O. Huisman, R. D'Ari and M. Hofnung, SOS chromotest, a direct assay of induction of an SOS function in *Escherichia coli*, K-12 to measure genotoxicity, *Proc. Natl. Acad. Sci. U. S. A.*, 1982, **79**, 5971–5975.
- 21 T. Ohta, M. Watanabe, R. Tsukamoto, Y. Shirasu and T. Kada, Antimutagenic effects of 5-fluorouracil and 5-fluorodeoxyuridine on UV-induced mutagenesis in *Escherichia coli*, *Mutat. Res.*, 1986, **173**, 19–24.
- 22 O. Huisman and R. D'Ari, An inducible DNA replication-cell division coupling mechanism in *Escherichia coli*, *Nature*, 1981, **290**, 797–799.
- 23 M. Rosenberg and H. Echols, Differential Recognition of Ultraviolet Lesions by RecA Protein, *J. Biol. Chem.*, 1990, **265**, 20641–20645.
- 24 J. Courcelle, J. R. Donaldson, K. H. Chow and C. T. Courcelle, DNA damage induced replication fork regression and processing in *Escherichia coli*, *Science*, 2003, **299**, 1064–1067.
- 25 P. Quillardet and M. Hofnung, Induction by UV light of the SOS function *sfia* in *Escherichia coli* strains deficient or proficient in excision repair, *J. Bacteriol.*, 1984, **157**, 35–38.
- 26 M. S. Eller, A. Asarch and B. A. Gilchrest, Photoprotection in human skin- A multifaceted SOS response, *Photochem. Photobiol.*, 2008, **84**, 339–349.

- 27 N. Bandyopadhyay, S. Gautam and A. Sharma, Suppression of SOS repair in *E. coli*: possible mechanism of antimutagenicity and protective effects of common vegetables, *Int. J. Food Sci. Nutr.*, 2014, **65**, 251–258.
- 28 C. A. Prada-Medina, E. T. Aristizabal-Tessmer, N. Quintero-Ruiz, J. Serment-Guerrero and J. L. Fuentes, Survival and SOS response induction in ultraviolet B irradiated *Escherichia coli* cells with defective repair mechanisms, *Int. J. Radiat. Biol.*, 2016, **92**, 321–328.
- 29 E. Zeiger, What is needed for an acceptable antimutagenicity manuscript?, *Mutat. Res.*, 2007, **626**, 1–3.
- 30 P. Quillardet and M. Hofnung, The SOS Chromotest, a colorimetric bacterial assay for genotoxins: procedures, *Mutat. Res.*, 1985, **147**, 65–78.
- 31 P. A. White and J. B. Rasmussen, SOS Chromotest results in a broader context: Empirical relationships between genotoxic potency, mutagenic potency, and carcinogenic potency, *Environ. Mol. Mutagen.*, 1996, **27**, 270–305.
- 32 P. A. White, J. B. Rasmussen and C. Blaise, A semi-automated, microplate version of the SOS Chromotest for the analysis of complex environmental extracts, *Mutat. Res.*, 1996, **360**, 51–74.
- 33 E. E. Stashenko, J. R. Martínez, M. P. Cala, D. C. Durán and D. Caballero, Chromatographic and mass spectrometric characterization of essential oils and extracts from *Lippia* (Verbenaceae) aromatic plants, *J. Sep. Sci.*, 2013, **36**, 192–202.
- 34 P. Cos, A. J. Vlietinck, D. Vanden-Berghe and L. Maes, Anti-infective potential of natural products: How to develop a stronger *in vitro* ‘proof-of-concept’, *J. Ethnopharmacol.*, 2006, **106**, 290–302.
- 35 N. Quintero, E. E. Stashenko and J. L. Fuentes, The influence of organic solvents on estimates of genotoxicity and antigenotoxicity in the SOS chromotest, *Genet. Mol. Biol.*, 2012, **35**, 503–514.
- 36 J. L. Fuentes, M. Vernhe, E. B. Cuetara, A. Sánchez-Lamar, J. L. Santana and M. Llagostera, Tannins from barks of *Pinus caribaea*, Morelet protect *Escherichia coli* cells against DNA damage induced by  $\gamma$ -rays, *Fitoterapia*, 2006, **77**, 116–120.
- 37 D. Huang, B. Ou, M. Hampsch-Woodill, J. A. Flanagan and E. K. Deemer, Development and validation of oxygen radical absorbance capacity assay for lipophilic antioxidants using randomly methylated  $\beta$ -cyclodextrin as the solubility enhancer, *J. Agric. Food Chem.*, 2002, **50**, 1815–1821.
- 38 D. Huang, B. Ou, M. Hampsch-Woodill, J. A. Flanagan and R. L. Prior, High-throughput assay of oxygen radical absorbance capacity (ORAC) using a multichannel liquid handling system coupled with a microplate fluorescence reader in 96-well format, *J. Agric. Food Chem.*, 2002, **50**, 4437–4444.
- 39 R Core Team, *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria, 2013, <http://www.R-project.org>.
- 40 J. D. Regan and H. Yoshida, DNA UVB dosimeters, *J. Photochem. Photobiol., B*, 1995, **31**, 57–61.
- 41 M. Lemay and K. A. Wood, Detection of DNA damage and identification of UV-induced photoproducts using the Comet assay™ Kit, *BioTechniques*, 1999, **27**, 846–851.
- 42 F. Hazane, S. Sauvaigo, T. Douki, A. Favier and J. C. Beani, Age-dependent DNA repair and cell cycle distribution of human skin fibroblasts in response to UVA irradiation, *J. Photochem. Photobiol., B*, 2006, **82**, 214–223.
- 43 A. P. Schuch and M. C. S. Menck, The genotoxic effects of DNA lesions induced by artificial UV-radiation and sunlight, *J. Photochem. Photobiol., B*, 2010, **99**, 111–116.
- 44 A. P. Schuch, M. C. S. Moraes, T. Yagura and C. F. M. Menck, Highly sensitive biological assay for determining the photoprotective efficacy of sunscreen, *Environ. Sci. Technol.*, 2014, **48**, 11584–11590.
- 45 L. F. Yamaguchi, M. J. Kato and P. Di Mascio, Biflavonoids from *Araucaria angustifolia* protect against DNA UV-induced damage, *Phytochemistry*, 2009, **70**, 615–620.
- 46 F. Bakkali, S. Averbeck, D. Averbeck, A. Zhiri, D. Baudoux and M. Idaomar, Antigenotoxic effects of three essential oils in diploid yeast (*Saccharomyces cerevisiae*) after treatments with UVC radiation, 8-MOP plus UVA and MMS, *Mutat. Res.*, 2006, **606**, 27–38.
- 47 C. A. Elmets, D. Singh, K. Tubesing, M. Matsui, S. Katiyar and H. Mukhtar, Cutaneous photoprotection from ultraviolet injury by green tea polyphenols, *J. Am. Acad. Dermatol.*, 2001, **44**, 425–432.
- 48 S. M. Meeran, S. Akhtar and S. K. Katiyar, Inhibition of UVB-induced skin tumor development by drinking green tea polyphenols is mediated through DNA repair and subsequent inhibition of inflammation, *J. Invest. Dermatol.*, 2009, **129**, 1258–1270.
- 49 A. Schwarz, A. Maeda, D. Gan, T. Mammone, M. S. Matsui and T. Schwarz, Green tea phenol extracts reduce UVB-induced DNA damage in human cells via interleukin-12, *Photochem. Photobiol.*, 2008, **84**, 350–355.
- 50 N. Morley, T. Clifford, L. Salter, S. Campbell, D. Gould and A. Curnow, The green tea polyphenol (–)-epigallocatechin gallate and green tea can protect human cellular DNA from ultraviolet and visible radiation-induced damage, *Photodermatol., Photoimmunol. Photomed.*, 2005, **21**, 15–22.
- 51 S. M. Meeran, S. K. Mantena and S. K. Katiyar, Prevention of ultraviolet radiation-induced immunosuppression by (–)-epigallocatechin-3-gallate in mice is mediated through interleukin 12-dependent DNA repair, *Clin. Cancer Res.*, 2006, **12**, 2272–2280.
- 52 S. K. Katiyar, Green tea prevents non-melanoma skin cancer by enhancing DNA repair, *Arch. Biochem. Biophys.*, 2011, **508**, 152–158.
- 53 A. C. T. S. Júnior, L. M. B. O. Asad, E. B. de Oliveira, K. Kovary, N. R. Asad and I. Felzenszwalb, Antigenotoxic and antimutagenic potential of an annatto pigment (norbixin) against oxidative stress, *Genet. Mol. Res.*, 2005, **4**, 94–99.
- 54 B. K. Pereira, R. M. Rosa, J. da Silva, T. N. Guecheva, I. M. de Oliveira, M. Ianistcki, V. C. Benvegnú, G. V. Furtado, A. Ferraz, M. F. Richter, N. Schroder,

- A. B. Pereira and J. A. P. Henriques, Protective effects of three extracts from Antarctic plants against ultraviolet radiation in several biological models, *J. Photochem. Photobiol. B*, 2009, **96**, 117–129.
- 55 F. Giampieri, J. M. Alvarez-Suarez, S. Tulipani, A. M. Gonz  les-Param  s, C. Santos-Buelga, S. Bompadre, J. L. Quiles, B. Mezzetti and M. Battino, Photoprotective potential of strawberry (*Fragaria*  $\times$  *Ananassa*) extract against UV-A irradiation damage on human fibroblasts, *J. Agric. Food Chem.*, 2012, **60**, 2322–2327.
- 56 J. Sundari, R. Selvaraj, N. Rajendra-Prasad and R. Elumalai, Jatropha curcas, leaf and bark fractions protect against ultraviolet radiation-B induced DNA damage in human peripheral blood lymphocytes, *Environ. Toxicol. Pharmacol.*, 2013, **36**, 875–882.
- 57 S. Das, J. Das, A. Paul, A. Samadder and A. R. Khuda-Bukhsh, Apigenin bioactive flavonoid from *Lycopodium clavatum*, stimulates nucleotide excision repair Genes to protect skin keratinocytes from ultraviolet B-Induced reactive oxygen species and DNA Damage, *J. Acupunct. Meridian. Stud.*, 2013, **6**, 252–262.
- 58 M. Vernhes, M. Gonz  lez-Pumari  ga, L. Andrade, A. P. Schuch, K. M. de Lima-Bessa, C. F. M. Menck and A. S  nchez-Lamar, Protective effect of a *Phyllanthus orbicularis*, aqueous extract against UVB light in human cells, *Pharm. Biol.*, 2013, **51**, 1–7.
- 59 P. Bolfa, R. Vidrighinescu, A. Petruta, D. Dezmirean, L. Stan, L. Vlase, G. Damian, C. Catoi, A. Filip and S. Clichici, Photoprotective effects of Romanian propolis on skin of mice exposed to UVB irradiation, *Food Chem. Toxicol.*, 2013, **62**, 329–342.
- 60 J. Psotova, A. Svobodova, H. Kolarova and D. Walterova, Photoprotective properties of *Prunella vulgaris* and rosmarinic acid on human keratinocytes, *J. Photochem. Photobiol. B*, 2006, **84**, 167–174.
- 61 D. Simi  , B. Vukovi  -Ga  i   and J. Kne  zevi  -Vuk  evi  , Detection of natural bioantimutagens and their mechanisms of action with bacterial assay-system, *Mutat. Res.*, 1998, **402**, 51–57.
- 62 J. Kne  zevi  -Vuk  evi  , B. Vukovi  -Ga  i  , T. Stevi  , J. Stanojevi  , B. Nikoli   and D. Simi  , Antimutagenic effect of essential oil of sage (*Salvia officinalis*, L.) and its fractions against UV-induced mutations in bacterial and yeast cells, *Arch. Biol. Sci.*, 2005, **57**, 163–172.
- 63 B. Vukovi  -Ga  i  , S. Nik  e  , T. Beri  -Bjedov, J. Kne  zevi  -Vuk  evi   and D. Simic, Antimutagenic effect of essential oil of sage (*Salvia officinalis*, L.) and its monoterpenes against UV-induced mutations in *Escherichia coli* and *Saccharomyces cerevisiae*, *Food Chem. Toxicol.*, 2006, **44**, 1730–1738.
- 64 S. K. Katiyar, S. K. Mantena and S. M. Meeran, Silymarin protects epidermal keratinocytes from ultraviolet radiation-induced apoptosis and DNA damage by nucleotide excision repair mechanism, *PLoS One*, 2011, **6**, e21410.
- 65 Y. J. Hyun, M. J. Piao, M. H. Ko, N. H. Lee, H. K. Kang, E. S. Yoo, Y. S. Koh and J. W. Hyun, Photoprotective effect of *Undaria crenata* against ultraviolet B-induced damage to keratinocytes, *J. Biosci. Bioeng.*, 2013, **116**, 256–264.
- 66 M. Vaid, S. D. Sharma and S. K. Katiyar, Proanthocyanidins inhibit photocarcinogenesis through enhancement of DNA Repair and Xeroderma Pigmentosum group A-dependent mechanism, *Cancer Prev. Res.*, 2010, **3**, 1621–1619.
- 67 N. Quintero-Ruiz, Y. Cordoba-Campo, E. E. Stashenko and J. L. Fuentes, Antigenotoxicity effect against ultraviolet radiation-induced DNA damage of the essential oils from *Lippia* species, *Photochem. Photobiol.*, 2017, **93**, 1063–1072.
- 68 D. R. Oliveira, G. G. Leit  o, P. D. Fernandes and S. G. Leit  o, Ethnopharmacological studies of *Lippia origanoides*, *Rev. Bras. Farmacogn.*, 2014, **24**, 206–214.