

Cat Food

Input file: **standard input**
Output file: **standard output**
Time limit: 1 second
Memory limit: 256 megabytes

One day, while Meow is drinking Milo, Meow sees a van fetching her favorite cat food pass by. Unfortunately for the manufacturer, the driver forgot to close the rear door and cat food is dropping out on to the road. Meow decides to collect and return the cat food to the driver because he is such a benevolent cat (though it is a bit dirty and will probably give you covid or something. Kids, don't try this at home.). What a purr-fect plan.



For each meter, depending on the smoothness of the road, the cat food may or may not drop from the van.

If Meow is at the start of i^{th} meter, there are 2 approaches Meow can move:

1. Walk 1 meter to the right. Meow will pick up the cat food if there's cat food at i^{th} meter. Now Meow at the start of $(i + 1)^{th}$ meter.
2. Jump x meter to the start of $(i + x)^{th}$ meter. If there's cat food at $(i + x)^{th}$ meter, Meow jumps beside it. Meow has to walk as mentioned in the first approach to collect the cat food.

Meow positions at the start of 1^{st} meter of the road (No movement made). There are n sequences of cat food on the road and Meow wants to collect all of them. The i^{th} sequence starts at the l_i^{th} meter and ends at the r_i^{th} meter inclusive. In other words, the i^{th} sequence of cat food has a total of $(r_i - l_i + 1)$ cat food and they are positioned at the l_i^{th} , $l_i^{th} + 1$, $l_i^{th} + 2$, \dots , r_i^{th} meter.

Find the minimum amount of distance Meow has to walk in meters to collect all the cat food.

Input

The first line contains two space-separated integers n , x ($1 \leq n \leq 10^5$, $1 \leq x \leq 10^5$) — the number of the cat food sequences and the value of x for the jump.

The next n lines each contain the information about the cat food, the i^{th} line contains two space-separated integers l_i , r_i ($1 \leq l_i \leq r_i \leq 10^5$) — the starting meter and the ending meter of the i^{th} sequence.

Output

Output a single number — The minimum amount of distance Meow has to walk in meters to collect all the cat food and return to the driver.

Examples

standard input	standard output
2 3 5 6 10 12	6
1 2 1 50000	50000
2 4 100 100 22 80	64

Note

In the first sample, Meow is initially at the start of 1st meter, since there is no cat food from the 1st meter to the 3rd meter, Meow jumps and lands at the start of 4th meter. Since there are cat foods at both 5th meter and 6th meter, Meow should not jump, therefore Meow walks from the start of 4th meter to the end of 4th meter, the end of 5th (collect cat food) and the end of 6th meter (collect cat food). Then, Meow jumps and skips 3 meters and lands at the start of 10th meter. Meow walks from there to the end of 12th meter to collect cat food from 10th meter to 12th meter. In total, Meow walks 6 meters to collect 5 cat food and returns to the driver. The image above illustrates this case.

In the second example, the road is full of cat food, so Meow will walk all the way until the end to collect all of them.