

# Using Numerical Methods to Explore Quantum Mechanics

Marc John Bordier Dam

April 30, 2017

## 1 Summary

WRITE ME!

## 2 Introduction

WRITE ME!

## 3 Schrödinger's Equation and its Solutions

The most fundamental equation in quantum mechanics is the Schrödinger equation. It governs which functions are worthy of study, that is, which functions are allowed wave functions in a given potential. The time-dependent Schrödinger equation in one dimension is

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V\Psi. \quad (1)$$

To make solving the time-dependent Schrödinger equation easier we use separation of variables to obtain the time-independent Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V\psi = E\psi, \quad (2)$$




Figure 1: Wave Functions in the Infinite Square well

where  $\Psi = \psi e^{-\frac{iEt}{\hbar}}$ .

The wave functions which are solutions to the time-independent Schrödinger equation represent stationary states. These stationary states have the nice property that any function in Hilbert space can be written as a linear combination of them. Thus it is in many cases sufficient to work with the stationary states and linear combinations of them. Below we will present the stationary states along with their allowed energies for the infinite and finite square well.

### 3.1 The Infinite Square Well

In the infinite square well the potential is given by

$$V = \begin{cases} 0, & 0 \leq x \leq a \\ \infty, & \text{otherwise} \end{cases}. \quad (3)$$

Outside the well the wave function is 0 and inside the well it is given by

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi}{a}x\right), \quad (4)$$

where  $n$  is any positive integer. In the infinite square well the energy spectrum is discrete with the  $n$ th stationary state having the energy

$$E_n = \frac{n^2\pi^2\hbar^2}{2ma^2}. \quad (5)$$

Some of the wave functions are shown in figure 1.




Figure 2: Wave Functions in the Finite Square well

### 3.2 The Finite Square Well

In the finite square well the potential is given by

$$V = \begin{cases} -V_0, & -a \leq x \leq a \\ 0, & \text{otherwise} \end{cases}, \quad (6)$$

where  $V_0$  is a positive number.

In the finite square well the wave function can be found both inside and outside the well. We also have both bound ( $E < 0$ ) and scattering states ( $E > 0$ ) in the finite square well.

The bound states come in two flavors: Even and odd. The even wave functions are given by

$$\psi_{\text{even}}(x) = \begin{cases} Ae^{-\kappa x}, & a < x \\ A \frac{e^{-\kappa a}}{\cos la} \cos lx, & -a \leq x \leq a \\ Ae^{\kappa x}, & x < -a \end{cases}, \quad (7)$$

where  $A$  is a normalization constant,  $\kappa = \frac{\sqrt{-2mE}}{\hbar}$  and  $l = \frac{\sqrt{2m(E+V_0)}}{\hbar}$ .

Similarly the odd wave functions are given by

$$\psi_{\text{odd}}(x) = \begin{cases} Ae^{-\kappa x}, & a < x \\ A \frac{e^{-\kappa a}}{\sin la} \sin lx, & -a \leq x \leq a \\ -Ae^{\kappa x}, & x < -a \end{cases}, \quad (8)$$

where again  $A$  is a normalization constant,  $\kappa = \frac{\sqrt{-2mE}}{\hbar}$  and  $l = \frac{\sqrt{2m(E+V_0)}}{\hbar}$ .

The energy spectrum of the bound states is discrete but can not be determined analytically. Defining  $z = la$  and  $z_0 = \frac{a}{\hbar}\sqrt{2mV_0}$  the allowed energies for the even wave functions can be found by solving the equation

$$\tan z = \sqrt{\left(\frac{z_0}{z}\right)^2 - 1}. \quad (9)$$

Likewise the allowed energies for the odd wave functions can be found by solving the equation

$$-\cot z = \sqrt{\left(\frac{z_0}{z}\right)^2 - 1}. \quad (10)$$

Some of the wave functions for the bound states are shown in figure 2.

The wave functions for the scattering states are given by

$$\psi(x) = \begin{cases} Fe^{ikx} & a < x \\ C \sin lx + D \cos lx & -a \leq x \leq a, \\ Ae^{ikx} + Be^{-ikx} & x < -a \end{cases} \quad (11)$$

where  $A$  is a normalization constant and

$$\begin{aligned} k &= \frac{\sqrt{2mE}}{\hbar}, \\ l &= \frac{\sqrt{2m(E+V_0)}}{\hbar}, \\ B &= i \frac{\sin 2la}{2kl} (l^2 - k^2) F, \\ C &= \left( \sin la + i \frac{k}{l} \cos la \right) e^{ika} F, \\ D &= \left( \cos la - i \frac{k}{l} \sin la \right) e^{ika} F, \\ F &= \frac{e^{-2ika}}{\cos 2la - i \frac{k^2 + l^2}{2kl} \sin 2la} A. \end{aligned}$$

For the scattering states we observe reflection and transmission through the well wall.

The energy spectrum of the scattering states is continuous and thus any positive energy is allowed.

Table 1: Reduced Units (subscript  $r$  denotes the quantity in real units)

Quantity (Q.)	Q. in R. U.	Characteristic Q.	Electron in $\pi$ -bond
Length, $x$	$\frac{x_r}{a}$	$a$	$2.8 \text{ \AA}$
Mass, $m$	$\frac{m_r}{m_0}$	$m_0$	$m_e = 9.109 \times 10^{-31} \text{ kg}$
Time, $t$	$\frac{t_r}{t_0}$	$t_0 = \frac{m_0 a^2}{\hbar}$	$6.77 \times 10^{-16} \text{ s}$
Energy, $E$	$\frac{E_r}{E_0}$	$E_0 = \frac{\hbar^2}{m_0 a^2}$	$1.56 \times 10^{-19} \text{ J}$
Velocity, $v$	$\frac{v_r}{v_0}$	$v_0 = \frac{\hbar}{m_0 a}$	$4.13 \times 10^5 \text{ m s}^{-1}$
Momentum, $p$	$\frac{p_r}{p_0}$	$p_0 = \frac{\hbar}{a}$	$3.77 \times 10^{-25} \text{ m kg s}^{-1}$

## 4 Quantum Mechanics in Reduced Units

When doing simulations we want to work with the physical quantities in reduced units. We do this for two reasons: We don't have to keep track of the units on everything and quantities in reduced units tend to be of a reasonable order of magnitude.

In our simulations we have three constants: The particle mass  $m_0$ , the well width  $a$  and Planck's constant  $\hbar$ . In our system of reduced units these will all be equal to 1. With these three constants we can derive the reduced units for the remaining relevant quantities in our simulation. The relevant quantities are shown in table 1, where the real values for an electron in a  $\pi$ -bond in an ethylene molecule is also shown.

## 5 Numerical Methods

## 6 Numerical Results and Discussions

## 7 Appendix

ALL THE CODE!!!!