

Properties of the dot product

$$(i) \vec{0} \cdot \vec{v} = \vec{v} \cdot \vec{0} = 0$$

$$(ii) \vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v} \quad (\text{Commutativity}).$$

$$(iii) (\lambda \vec{v}) \cdot \vec{w} = \vec{v} \cdot (\lambda \vec{w}) = \lambda (\vec{v} \cdot \vec{w})$$

$$(iv) \left. \begin{aligned} \vec{u} \cdot (\vec{v} + \vec{w}) &= \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w} \\ (\vec{v} + \vec{w}) \cdot \vec{u} &= \vec{v} \cdot \vec{u} + \vec{w} \cdot \vec{u} \end{aligned} \right\}$$

(Distributive law)

$$(v) \vec{v} \cdot \vec{v} = \|\vec{v}\|^2 \quad ; \text{ relation with length.}$$

The next theorem establishes the relation between the dot product & the angle between two vectors. Hence the angle between two vectors is chosen to satisfy $0 \leq \theta \leq \pi$.

Theorem (Dot product & the angle).

Let θ be the angle between two non-zero vectors \vec{v} & \vec{w} . Then

$$\vec{v} \cdot \vec{w} = \|\vec{v}\| \|\vec{w}\| \cos \theta \quad \text{or}$$

$$\cos \theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|}$$

Proof:



The laws of cosines gives

$$\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2 \cos \theta \|\vec{v}\| \|\vec{w}\| \quad (1)$$

$$\begin{aligned} \text{Now } \|\vec{v} - \vec{w}\|^2 &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) \\ &= \vec{v} \cdot \vec{v} - 2\vec{v} \cdot \vec{w} + \vec{w} \cdot \vec{w} \\ &= \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\vec{v} \cdot \vec{w} \end{aligned}$$

\therefore Therefore by (1) $=$

$$\boxed{\vec{v} \cdot \vec{w} = \|\vec{v}\| \|\vec{w}\| \cos \theta.}$$

Ex. Find the angle θ between

$$\vec{v} = \langle 3, 6, 2 \rangle \text{ \& } \vec{w} = \langle 4, 2, 4 \rangle.$$

Solⁿ

$$\|\vec{v}\| = \sqrt{49} = 7, \quad \|\vec{w}\| = \sqrt{36} = 6.$$

$$\cos \theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} = \frac{\langle 3, 6, 2 \rangle \cdot \langle 4, 2, 4 \rangle}{7 \cdot 6}$$

$$= \frac{3 \cdot 4 + 6 \cdot 2 + 2 \cdot 4}{42} = \frac{32}{42} = \frac{16}{21}$$

$$\therefore \theta = \cos^{-1}\left(\frac{16}{21}\right) //$$

Defⁿ (Perpendicular / Orthogonal).

Two non-zero vectors \vec{v} & \vec{w} are called perpendicular or orthogonal

if the angle between them is $\pi/2$.

We write $\vec{v} \perp \vec{w}$ if \vec{v} & \vec{w} are orthogonal. Thus we have

$$\vec{v} \perp \vec{w} \text{ if and only if } \vec{v} \cdot \vec{w} = 0$$

Ex. Determine if $\vec{v} = \langle 2, 6, 1 \rangle$ is orthogonal to $\vec{u} = \langle 2, -1, 1 \rangle$ or $\vec{w} = \langle -4, 1, 2 \rangle$

Solⁿ

$$\vec{v} \cdot \vec{u} = \langle 2, 6, 1 \rangle \cdot \langle 2, -1, 1 \rangle = 2 \cdot 2 + 6(-1) + 1 \cdot 1 \\ = -1 \quad (\text{not orthogonal}).$$

$$\vec{v} \cdot \vec{w} = \langle 2, 6, 1 \rangle \cdot \langle -4, 1, 2 \rangle = 2(-4) + 6(1) + 1(2) \\ = 0 \quad (\text{orthogonal})$$

⊛ The cross product or vector product is used in physics and engineering to describe quantities involving rotation such as torque and angular momentum. In electro-magnetic theory, magnetic forces, etc. . .

⊛ The cross product $\vec{v} \times \vec{w}$ is ~~again~~ again a vector.

⊛ It is defined using determinant,

For 2x2

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - cb.$$

For 3x3

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$$

$$- a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

Defⁿ (The Cross Product / Vector Product)

The cross product of vectors

$\vec{v} = \langle a_1, b_1, c_1 \rangle$ & $\vec{w} = \langle a_2, b_2, c_2 \rangle$ is

the vector

$$\vec{v} \times \vec{w} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix}$$

$$\vec{v} \times \vec{w} = \begin{vmatrix} b_1 & c_1 \\ b_2 & c_2 \end{vmatrix} \underline{i} - \begin{vmatrix} a_1 & c_1 \\ a_2 & c_2 \end{vmatrix} \underline{j} + \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} \underline{k}$$

Ex Calculate $\vec{v} \times \vec{w}$, where $\vec{v} = \langle -2, 1, 4 \rangle$ & $\vec{w} = \langle 3, 2, 5 \rangle$.

Solⁿ

$$\vec{v} \times \vec{w} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ -2 & 1 & 4 \\ 3 & 2 & 5 \end{vmatrix} = \begin{vmatrix} 1 & 4 \\ 2 & 5 \end{vmatrix} \underline{i} - \begin{vmatrix} -2 & 4 \\ 3 & 5 \end{vmatrix} \underline{j} + \begin{vmatrix} -2 & 1 \\ 3 & 2 \end{vmatrix} \underline{k}$$

$$= \begin{vmatrix} 1 & 4 \\ 2 & 5 \end{vmatrix} \underline{i} - \begin{vmatrix} -2 & 4 \\ 3 & 5 \end{vmatrix} \underline{j} + \begin{vmatrix} -2 & 1 \\ 3 & 2 \end{vmatrix} \underline{k}$$

~~$$= -3\underline{i} - (-22)\underline{j} + (-7)\underline{k}$$~~

~~$$\vec{v} \times \vec{w} = \langle -3, 22, -7 \rangle //$$~~

$$= \underline{i}(5-8) - \underline{j}(-10-12) + \underline{k}(-4-3)$$


$$= \underline{i}(-3) - \underline{j}(-22) + \underline{k}(-7)$$

$$= -3\underline{i} + 22\underline{j} - 7\underline{k}$$

$$= \langle -3, 22, -7 \rangle //$$

Note

(*) $\{\vec{v}, \vec{w}, \vec{v} \times \vec{w}\}$ forms a right handed system.



i.e. when the fingers of your right hand curl from \vec{v} to \vec{w} , your ~~thumb~~ thumb points to the same side of the plane spanned by \vec{v} & \vec{w} .

Theorem (Geometric Description of the Cross Product).

The cross product $\vec{v} \times \vec{w}$ is the unique vector with the following three properties:

(i) $\vec{v} \times \vec{w}$ is orthogonal to \vec{v} & \vec{w} .

(ii) $\vec{v} \times \vec{w}$ has length $\|\vec{v}\| \|\vec{w}\| |\sin \theta|$,
 $\theta = \text{angle between } \vec{v} \text{ \& } \vec{w}$.

(iii) $\{\vec{v}, \vec{w}, \vec{v} \times \vec{w}\}$ forms a right-handed system.

Basic Properties of the Cross Product

(i) $\vec{w} \times \vec{v} = -\vec{v} \times \vec{w}$

(ii) $\vec{v} \times \vec{v} = \vec{0}$

(iii) $\vec{v} \times \vec{w} = \vec{0}$ iff $\vec{w} = \lambda \vec{v}$ for
some scalar λ or $\vec{v} = \vec{0}$

(iv) $(\lambda \vec{v}) \times \vec{w} = \vec{v} \times (\lambda \vec{w}) = \lambda (\vec{v} \times \vec{w})$

(v) $(\vec{u} + \vec{v}) \times \vec{w} = (\vec{u} \times \vec{w}) + (\vec{v} \times \vec{w})$

$\vec{v} \times (\vec{u} + \vec{w}) = (\vec{v} \times \vec{u}) + (\vec{v} \times \vec{w})$

Ex

$$\underline{i} \times \underline{j} = \underline{k} \quad , \quad \underline{j} \times \underline{k} = \underline{i} \quad , \quad \underline{k} \times \underline{i} = \underline{j}$$

$$\underline{i} \times \underline{i} = \underline{j} \times \underline{j} = \underline{k} \times \underline{k} = \underline{0}.$$

Ex

Compute $(2\underline{i} + \underline{k}) \times (3\underline{j} + 5\underline{k})$.

Sol

$$\begin{aligned} & (2\underline{i} + \underline{k}) \times (3\underline{j} + 5\underline{k}) \\ &= (2\underline{i} \times 3\underline{j}) + (2\underline{i} \times 5\underline{k}) + (\underline{k} \times 3\underline{j}) \\ & \quad + (\underline{k} \times 5\underline{k}) \end{aligned}$$

$$= 6(\underline{i} \times \underline{j}) + 10(\underline{i} \times \underline{k}) + 3(\underline{k} \times \underline{j}) + 5(\underline{k} \times \underline{k})$$


$$= 6\underline{k} - 10\underline{j} - 3\underline{i} + 5\underline{0}$$

$$= \langle -3, -10, 6 \rangle //$$

Relation between the Cross Product, Area and Volume

(i) Area:

Consider the parallelogram P spanned by non-zero vectors \vec{v} & \vec{w} with a common basepoint.




The parallelogram has a base $b = \|\vec{v}\|$ and height $h = \|\vec{w}\| \sin \theta$,

$$\begin{aligned}\therefore \text{Therefore area of } P &= bh \\ &= \|\vec{v}\| \|\vec{w}\| \sin \theta \\ &= \|\vec{v} \times \vec{w}\|\end{aligned}$$

(II) Volume.

6

Consider the parallelepiped P spanned by three non-zero vectors $\vec{u}, \vec{v}, \vec{w}$ in \mathbb{R}^3 .



The base of P is the parallelogram spanned by \vec{v} & \vec{w} , so the area of the base is $\|\vec{v} \times \vec{w}\|$. The height of P is, $h = \|\vec{u}\| \cdot |\cos \theta|$ where θ is the angle between \vec{u} & $\vec{v} \times \vec{w}$.

$$\begin{aligned} \therefore \text{Volume of } P &= (\text{area of base}) \times \text{height} \\ &= \|\vec{v} \times \vec{w}\| \cdot \|\vec{u}\| \cdot |\cos \theta| \\ &= \left| \vec{u} \cdot (\vec{v} \times \vec{w}) \right| \end{aligned}$$

Note

(*) The quantity $\vec{u} \cdot (\vec{v} \times \vec{w})$,
called the vector triple product,

can be expressed as a following
way. Let $\vec{u} = \langle a_1, b_1, c_1 \rangle$, $\vec{v} = \langle a_2, b_2, c_2 \rangle$
& $\vec{w} = \langle a_3, b_3, c_3 \rangle$ - Then.

$$\vec{u} \cdot (\vec{v} \times \vec{w}) = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$