

Eigenvalues & Eigenvectors

Let A is an $n \times n$ matrix,
then a scalar λ is called
an eigenvalue of A , if
associated with it there
is a non-zero vector \underline{v} ,
called an eigenvector, such
that

$$A \underline{v} = \lambda \underline{v}$$

④ To find the eigenvalues solve
the characteristic equation.

$$|A - \lambda I| = 0$$

(*) To find the eigenvectors
solve

$$(A - \lambda I) \underline{v} = \underline{0}.$$

(*) Note that, scalar λ can be
real number or complex number.

Ex. Find the eigenvalues and
corresponding eigenvectors of

$$A = \begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix}_{2 \times 2}.$$

Solⁿ.

First find eigenvalues
using characteristic equation.

$$|A - \lambda I| = 0$$

12

$$\left| \begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right| = 0$$

$$\begin{vmatrix} 4-\lambda & -5 \\ 2 & -3-\lambda \end{vmatrix} = 0$$

$$(4-\lambda)(-3-\lambda) + 10 = 0$$

$$\lambda^2 - \lambda - 2 = 0$$

$$\lambda = -1, 2.$$

Next find corresponding eigenvectors for each eigen values.

For $\lambda = -1$

To find eigenvectors solve

$$(A - \lambda I)\underline{v} = \underline{0} \quad \text{or} \quad \lambda = -1$$

$$\text{i.e. } \begin{pmatrix} 5 & -5 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{let } \underline{v} = \begin{pmatrix} x \\ y \end{pmatrix}_{2 \times 1}$$

$$\begin{pmatrix} 5x - 5y \\ 2x - 2y \end{pmatrix}_{2 \times 1} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}_{2 \times 1}$$

$$\Rightarrow \begin{cases} 5x - 5y = 0 \\ 2x - 2y = 0 \end{cases} \quad x - y = 0$$

If we take t is any parameter and set $y=t$, then $x=y=t$.

Therefore corresponding eigen vectors for $\lambda = -1$,

$$E_{-1} = \underline{v} = \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} t \\ t \end{pmatrix} = t \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$E_{-1} = \left\{ t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \mid t \in \mathbb{R} \right\}$$

↑
eigen space for eigen value $\lambda = -1$.

For $\lambda = 2$

$$(A - \lambda I) \underline{v} = \underline{0}$$

$$\begin{pmatrix} 2 & -5 \\ 2 & -5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow 2x - 5y = 0$$

\Rightarrow If we set $y = t$, then

$$x = \frac{5y}{2} = \frac{5t}{2}$$

\therefore Corresponding eigenvector for $\lambda = 2$

is eigen space.

$$E_{\lambda=2} = \underline{v} = \begin{pmatrix} x \\ y \end{pmatrix} = \left\{ t \begin{pmatrix} 5/2 \\ 1 \end{pmatrix} \mid t \in \mathbb{R} \right\}.$$

Note,

We denote E_λ is to represent all the eigenvectors together with 0 vector

corresponding to eigenvalue λ .

E_λ called as eigenspace.

Ex

Find eigenvalues & eigenvectors of $\bar{A} = \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix}_{2 \times 2}$.

Solⁿ.

(4)

For eigenvalues $|A - \lambda I| = 0$

$$\begin{vmatrix} 4-\lambda & 0 \\ 0 & 4-\lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda = 4, 4 \quad (\text{i.e. one-eigenvalue})$$

For $\lambda = 4$

$$(A - \lambda I) \underline{V} = \underline{0}$$

$$\begin{pmatrix} 4-4 & 0 \\ 0 & 4-4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$\Rightarrow x$ & y can be any ~~vector~~ ^{value.}

\Rightarrow let s & t be any parameter. Then

$x=s$ & $y=t$ given eigen vectors.

\therefore Corresponding eigen ~~vectors~~ ^{space}.

for $\lambda = 4$,

$$E_4 = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} s \\ t \end{pmatrix} / s, t \in \mathbb{R} \right\}$$

$$E_4 = \left\{ s \begin{pmatrix} 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} 0 \\ 1 \end{pmatrix} / s, t \in \mathbb{R} \right\}$$



Example.

Find the eigenvalues & Eigenspace of the matrix

$$A = \begin{pmatrix} 1 & -3 & 3 \\ 3 & -5 & 3 \\ 6 & -6 & 4 \end{pmatrix}_{3 \times 3}$$

Solⁿ

15

Let us find values of λ which satisfy the characteristic eqⁿ of the matrix A.

$$[A - \lambda I] = 0$$

$$\begin{vmatrix} 1-\lambda & -3 & 3 \\ 3 & -5-\lambda & 3 \\ 6 & -6 & 4-\lambda \end{vmatrix} = 0$$

$$\Rightarrow 16 + 12\lambda - \lambda^3 = 0$$

$$\Rightarrow \lambda = -2, 4.$$

E₂ for $\lambda = 4$

$$(A - \lambda I)\underline{v} = \underline{0} \quad ; \text{ where } \underline{v} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

Here we have three equations with three variables. Let us solve this linear system by ~~CRA~~ Gauss - Elimination method.

$$(A - \lambda I) = \begin{pmatrix} -3 & -3 & 3 \\ 3 & -9 & 3 \\ 6 & -6 & 0 \end{pmatrix}.$$

$$(A - \lambda I)\underline{v} = \underline{0}$$

$$\left(\begin{array}{ccc|c} -3 & -3 & 3 & 0 \\ 3 & -9 & 3 & 0 \\ 6 & -6 & 0 & 0 \end{array} \right) \xrightarrow{\text{After elementary row operations}}$$

$$\left(\begin{array}{ccc|c} 1 & 0 & -1/2 & 0 \\ 0 & 1 & -1/2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right).$$

$$\Rightarrow x_1 - \frac{1}{2}x_3 = 0$$

$$x_2 - \frac{1}{2}x_3 = 0$$

If $x_3 = t$, then $x_1 = t/2$ and $x_2 = t/2$.

So eigen space corresponding to the eigenvalue $\lambda = 4$,

$$E_4 = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} t/2 \\ t/2 \\ t \end{pmatrix} = \left\{ t \begin{pmatrix} 1/2 \\ 1/2 \\ 1 \end{pmatrix} \mid t \in \mathbb{R} \right\}$$

E_1 for $\lambda = -2$

(6)

$$A + 2I = \begin{pmatrix} 3 & -3 & 3 \\ 3 & -3 & 3 \\ 6 & -6 & 6 \end{pmatrix}.$$

$$(A + 2I)V = \underline{0}.$$

$$\left(\begin{array}{ccc|c} 3 & -3 & 3 & 0 \\ 3 & -3 & 3 & 0 \\ 6 & -6 & 6 & 0 \end{array} \right) \longrightarrow \left(\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$$x_1 - x_2 + x_3 = 0$$

Then setting parameters s & t as
 $x_2 = s$, $x_3 = t$, ~~then~~ we have

$$x_1 = s + t = t + s - t$$

$$E_{-2} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} s+t \\ s \\ t \end{pmatrix} = \begin{pmatrix} t \\ 0 \\ t \end{pmatrix} + \begin{pmatrix} s \\ s \\ 0 \end{pmatrix}.$$

$$E_{-2} = \left\{ t \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} + s \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \mid s, t \in \mathbb{R} \right\}$$

Let $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$

be a polynomial in x . If A
is an $n \times n$ matrix then $f(A)$
is defined as

$$f(A) = a_n A^n + a_{n-1} A^{n-1} + \dots + a_1 A + a_0 I$$

where I is the identity matrix of
order n .