

Tutorial (02)  
Solutions

(01)

(01)

$$(a) \quad 3x + 7y - 4z = -46$$

$$5w + 4x + 8y + z = 7$$

$$8w + 4y - 2z = 0$$

$$-w + 6x + 2z = 13$$

$\therefore$  Augmented Matrix :

$$\left( \begin{array}{cccc|c} 0 & 3 & 7 & -4 & -46 \\ 5 & 4 & 8 & 1 & 7 \\ 8 & 0 & 4 & -2 & 0 \\ -1 & 6 & 0 & 2 & 13 \end{array} \right)$$

$\downarrow R_1 \leftrightarrow R_2$

$$\left( \begin{array}{cccc|c} 5 & 4 & 8 & 1 & 7 \\ 0 & 3 & 7 & -4 & -46 \\ 8 & 0 & 4 & -2 & 0 \\ -1 & 6 & 0 & 2 & 13 \end{array} \right)$$

$\downarrow R_1 / 5 \rightarrow R_1$

$$\left( \begin{array}{cccc|c} 1 & 0.8 & 1.6 & 0.2 & 1.4 \\ 0 & 3 & 7 & -4 & -46 \\ 8 & 0 & 4 & -2 & 0 \\ -1 & 6 & 0 & 2 & 13 \end{array} \right)$$

$$\begin{array}{l} \downarrow R_3 - 8R_1 \rightarrow R_3 \\ R_4 + R_1 \rightarrow R_4 \end{array}$$

$$\left( \begin{array}{cccc|c} 1 & 0.8 & 1.6 & 0.2 & 1.4 \\ 0 & 3 & 7 & -4 & -46 \\ 0 & -6.4 & -8.8 & -3.6 & -11.2 \\ 0 & 6.8 & 1.6 & 2.2 & 14.4 \end{array} \right)$$

$$\downarrow R_2 / 3 \rightarrow R_2$$

$$\left( \begin{array}{cccc|c} 1 & 0.8 & 1.6 & 0.2 & 1.4 \\ 0 & 1 & \frac{7}{3} & -\frac{4}{3} & -\frac{46}{3} \\ 0 & -6.4 & -8.8 & -3.6 & -11.2 \\ 0 & 6.8 & 1.6 & 2.2 & 14.4 \end{array} \right)$$

$$\begin{aligned}
 & \downarrow \begin{aligned} R_1 - 0.8 R_2 &\rightarrow R_1 \\ R_3 + 6.4 R_2 &\rightarrow R_3 \\ R_4 - 6.8 R_2 &\rightarrow R_4 \end{aligned}
 \end{aligned}$$

$$\left( \begin{array}{cccc|c} 1 & 0 & -\frac{4}{15} & \frac{19}{15} & \frac{41}{3} \\ 0 & 1 & \frac{7}{3} & -\frac{4}{3} & -\frac{46}{3} \\ 0 & 0 & \frac{92}{15} & -\frac{182}{15} & -\frac{328}{3} \\ 0 & 0 & -\frac{214}{15} & \frac{169}{15} & \frac{356}{3} \end{array} \right)$$

$$\downarrow R_3 / \frac{92}{15} \rightarrow R_3$$

$$\left( \begin{array}{cccc|c} 1 & 0 & -\frac{4}{15} & \frac{19}{15} & \frac{41}{3} \\ 0 & 1 & \frac{7}{3} & -\frac{4}{3} & -\frac{46}{3} \\ 0 & 0 & 1 & -\frac{91}{46} & -\frac{410}{23} \\ 0 & 0 & -\frac{214}{15} & \frac{169}{15} & \frac{356}{3} \end{array} \right)$$

$$\begin{aligned}
 & \downarrow \begin{aligned} R_1 + \frac{4}{15} R_3 &\rightarrow R_1 \\ R_2 - \frac{7}{3} R_3 &\rightarrow R_2 \\ R_4 + \frac{214}{15} R_3 &\rightarrow R_4 \end{aligned}
 \end{aligned}$$

(04)

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & \frac{17}{23} \\ 0 & 1 & 0 & \frac{151}{46} \\ 0 & 0 & 1 & -\frac{91}{46} \\ 0 & 0 & 0 & -\frac{390}{23} \end{array} \right) \begin{array}{c} \frac{205}{23} \\ \frac{604}{23} \\ -\frac{410}{23} \\ -\frac{3120}{23} \end{array}$$

$$\downarrow R_4 / -\frac{390}{23} \rightarrow R_4$$

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & \frac{17}{23} \\ 0 & 1 & 0 & \frac{151}{46} \\ 0 & 0 & 1 & -\frac{91}{46} \\ 0 & 0 & 0 & 1 \end{array} \right) \begin{array}{c} \frac{205}{23} \\ \frac{604}{23} \\ -\frac{410}{23} \\ 8 \end{array}$$

$$\downarrow \begin{array}{l} R_1 - \frac{17}{23} R_4 \rightarrow R_1 \\ R_2 - \frac{151}{46} R_4 \rightarrow R_2 \\ R_3 + \frac{91}{46} R_4 \rightarrow R_3 \end{array}$$

(05)

$$\begin{pmatrix} 1 & 0 & 0 & 0 & | & 3 \\ 0 & 1 & 0 & 0 & | & 0 \\ 0 & 0 & 1 & 0 & | & -2 \\ 0 & 0 & 0 & 1 & | & 8 \end{pmatrix}$$

$$\therefore \begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \\ 8 \end{pmatrix}.$$

(02)

(06)

(a) Let  $x, y, z$  represent the number of small, medium, and large T-shirts, respectively.

$$x + y + z = 30$$

$$4x + 5y + 6z = 154$$

$$4x - 6z = -40$$

$\therefore$  the corresponding Augmented Matrix :

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 30 \\ 4 & 5 & 6 & 154 \\ 4 & 0 & -6 & -40 \end{array} \right)$$

then using Gauss - Jordan elimination, we can obtain the following row reduced-echelon form.

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & 8 \\ 0 & 1 & 0 & 10 \\ 0 & 0 & 1 & 12 \end{array} \right)$$

$$\therefore x = 8, \quad y = 10, \quad z = 12$$

(02) (b) (ii)

(07)

$$\begin{aligned}x - y + 2z &= 4 \\ -x + 3y + z &= -6 \\ x + y + 5z &= 3\end{aligned}$$

Augmented Matrix:

$$\left( \begin{array}{ccc|c} 1 & -1 & 2 & 4 \\ -1 & 3 & 1 & -6 \\ 1 & 1 & 5 & 3 \end{array} \right)$$

$$\downarrow R_1 + R_2 \rightarrow R_2$$

$$\left( \begin{array}{ccc|c} 1 & -1 & 2 & 4 \\ 0 & 2 & 3 & -2 \\ 1 & 1 & 5 & 3 \end{array} \right)$$

$$\downarrow -R_1 + R_3 \rightarrow R_3$$

$$\left( \begin{array}{ccc|c} 1 & -1 & 2 & 4 \\ 0 & 2 & 3 & -2 \\ 0 & 2 & 3 & -1 \end{array} \right) \xrightarrow{-R_2 + R_3 \rightarrow R_3}$$

$$\left( \begin{array}{ccc|c} 1 & -1 & 2 & 4 \\ 0 & 2 & 3 & -2 \\ 0 & 0 & 0 & 1 \end{array} \right)$$

$\therefore$  the last row represents

$$0 \cdot x + 0 \cdot y + 0 \cdot z = 1$$

$$\therefore 0 = 1$$

Contradiction

$\therefore$  the system is inconsistent and has no solution.



(02)  
(c)

(09)

$$(i) \quad x + 2y + 3z = 10$$

$$x + y + z = 7$$

$$3x + 2y + z = 18$$

$\therefore$  the augmented matrix :

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 10 \\ 1 & 1 & 1 & 7 \\ 3 & 2 & 1 & 18 \end{array} \right)$$

$$\downarrow -R_1 + R_2 \rightarrow R_2$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 10 \\ 0 & -1 & -2 & -3 \\ 3 & 2 & 1 & 18 \end{array} \right)$$

$$\downarrow -3R_1 + R_3 \rightarrow R_3$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 10 \\ 0 & -1 & -2 & -3 \\ 0 & -4 & -8 & -12 \end{array} \right) \xrightarrow{-R_2 \rightarrow R_2}$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 10 \\ 0 & 1 & 2 & 3 \\ 0 & -4 & -8 & -12 \end{array} \right)$$

$$\downarrow (-2)R_2 + R_1 \rightarrow R_1$$

$$\left( \begin{array}{ccc|c} 1 & 0 & -1 & 4 \\ 0 & 1 & 2 & 3 \\ 0 & -4 & -8 & -12 \end{array} \right)$$

$$\downarrow 4R_2 + R_3 \rightarrow R_3$$

$$\left( \begin{array}{ccc|c} 1 & 0 & -1 & 4 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$\therefore$  the last row corresponds to the identity  $0=0$  which is true for every value of  $x, y$ , and  $z$ .

from the first two equations,

$$x - z = 4$$

$$y + 2z = 3$$

$$\therefore x = z + 4$$

(11)

$$y = 3 - 2z$$

$\therefore$  the solutions of the system

$$(x, y, z) = (z + 4, 3 - 2z, z)$$

$\therefore$  the solution set is

$$\left\{ (z + 4, 3 - 2z, z) \mid z \in \mathbb{R} \right\}.$$

(02)

(12)

(c) (ii)

$$x + y + z = 1$$

$$2x - 2y + 6z = 10$$

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 2 & -2 & 6 & 10 \end{array} \right)$$

$$\downarrow -2R_1 + R_2 \rightarrow R_2$$

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 0 & -4 & 4 & 8 \end{array} \right)$$

$$\downarrow \left(-\frac{1}{4}\right)R_2 \rightarrow R_2$$

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 0 & 1 & -1 & -2 \end{array} \right)$$

$$\downarrow (-1)R_2 + R_1 \rightarrow R_1$$

$$\left( \begin{array}{ccc|c} 1 & 0 & 2 & 3 \\ 0 & 1 & -1 & -2 \end{array} \right)$$

(13)

$$x + 2z = 3 \quad \therefore x = 3 - 2z$$

$$y - z = -2 \quad y = z - 2$$

$$(x, y, z) = (3 - 2z, z - 2, z).$$

# Homogeneous System of Linear Equations

(14)

a homogeneous system of linear equations is as follows:

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n &= 0 \\a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n &= 0 \\&\vdots\end{aligned}$$

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n = 0$$

Where

$a_{ij}$  ;  $i = 1, 2, \dots, m$  ;  $j = 1, 2, \dots, n$  are

constants.

$\therefore$  the augmented matrix

$$\left( \begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & 0 \\ a_{21} & a_{22} & \dots & a_{2n} & 0 \\ & \vdots & & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & 0 \end{array} \right)$$

$$(A | b)$$

The above system is always satisfied (15)  
by  $x_1=0, x_2=0, \dots, x_n=0$

This solution is called the trivial solution.

\*: Any system of homogeneous linear equations is always consistent.

Case I :-  
if  $|A| \neq 0$  then the <sup>homogeneous</sup> system of linear equations has a unique solution and hence the only <sup>†</sup> solution to the system is the trivial solution.

Case II :-  
If  $|A| = 0$  then the homogeneous system has a non-trivial solution.

(03)  
(a)  
(1)

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 3 & 4 & 4 & 0 \\ 7 & 10 & 12 & 0 \end{array} \right)$$

$$\begin{array}{l} \downarrow R_2 \rightarrow R_2 - 3R_1 \\ \downarrow R_3 \rightarrow R_3 - 7R_1 \end{array}$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & -2 & -5 & 0 \\ 0 & -4 & -9 & 0 \end{array} \right)$$

$$\begin{array}{l} \downarrow R_2 \rightarrow R_2 / (-1) \\ \downarrow R_3 \rightarrow R_3 / (-1) \end{array}$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 2 & 5 & 0 \\ 0 & 4 & 9 & 0 \end{array} \right)$$

$$\downarrow R_3 \rightarrow R_3 - 2R_2$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 2 & 5 & 0 \\ 0 & 0 & -1 & 0 \end{array} \right) \xrightarrow{R_3 \rightarrow R_3 / (-1)}$$



$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 2 & 5 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right)$$

(17)

$$\therefore x = y = z = 0$$

$\therefore$  the system has a unique solution.

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 3 & 4 & 4 \\ 7 & 10 & 12 \end{vmatrix}$$

$$= -2 \neq 0$$

$\therefore$  the system has only the trivial solution.

(04)

(18)

(ii)

$$x + 3y - 2z = 0$$

$$2x - y + 4z = 0$$

$$x - 11y + 14z = 0$$

$$\left( \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 2 & -1 & 4 & 0 \\ 1 & -11 & 14 & 0 \end{array} \right)$$

$$\downarrow R_2 \rightarrow R_2 - 2R_1$$

$$\left( \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & -7 & 8 & 0 \\ 0 & -14 & 16 & 0 \end{array} \right)$$

$$\downarrow R_2 \rightarrow R_2 / (-7)$$

$$\left( \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 1 & -\frac{8}{7} & 0 \\ 0 & -14 & 16 & 0 \end{array} \right)$$

$$\downarrow R_1 \rightarrow R_1 - 3R_2$$

$$\left( \begin{array}{ccc|c} 1 & 0 & \frac{10}{7} & 0 \\ 0 & 1 & -\frac{8}{7} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$\therefore$  the system has an infinite number of solutions. (19)

$$x + \frac{10}{7}z = 0 \Rightarrow x = -\frac{10z}{7}$$

$$y - \frac{8}{7}z = 0 \Rightarrow y = \frac{8z}{7}$$

$\therefore$  the solution set:

$$\left\{ (x, y, z) \mid x = -\frac{10z}{7} ; y = \frac{8z}{7} ; z \in \mathbb{R} \right\}$$

$$= \left\{ \left( -\frac{10z}{7}, \frac{8z}{7}, z \right) \mid z \in \mathbb{R} \right\}$$

(05)

(20)

$$kx + y + z = 1$$

$$x + ky + z = 1$$

$$x + y + kz = 1$$

$$\therefore AX = b \quad \text{where}$$

$$A = \begin{pmatrix} k & 1 & 1 \\ 1 & k & 1 \\ 1 & 1 & k \end{pmatrix} ; X = \begin{pmatrix} x \\ y \\ z \end{pmatrix} ; b = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

(i) For the system of linear equations to have a unique solution,

$$|A| \neq 0$$

$$\therefore \begin{vmatrix} k & 1 & 1 \\ 1 & k & 1 \\ 1 & 1 & k \end{vmatrix} \neq 0$$

$$k(k^2 - 1) - 1(k - 1) + 1(1 - k) \neq 0$$

$$k(k - 1)(k + 1) - 2(k - 1) \neq 0$$

$$(k - 1)(k^2 + k - 2) \neq 0$$

$$(k - 1)^2(k + 2) \neq 0$$

$$\therefore k \neq 1 \quad \text{or} \quad k \neq -2$$

$\therefore |A| = 0$  When  $k = 1$  or  $k = -2$  (21)

$\therefore$  When  $k = 1$  or  $k = -2$  the system of linear equations are inconsistent. or (dependent with) infinitely many solutions.

(iii)  
When  $k = 1$ , the augmented matrix:

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{array} \right)$$

$$\begin{array}{l} \downarrow R_2 \rightarrow R_2 - R_1 \\ \downarrow R_3 \rightarrow R_3 - R_1 \end{array}$$

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$\therefore$  When  $k = 1$  ; the system has infinitely many solutions, and the solution set:

$$\left\{ (x, y, z) \mid x + y + z = 1 \quad ; \quad x, y, z \in \mathbb{R} \right\}$$

(iii) When  $k = -2$ ; the augmented matrix: (22)

$$\left( \begin{array}{ccc|c} -2 & 1 & 1 & 1 \\ 1 & -2 & 1 & 1 \\ 1 & 1 & -2 & 1 \end{array} \right)$$

$$\downarrow R_1 \rightarrow R_1 / (-2)$$

$$\left( \begin{array}{ccc|c} 1 & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 1 & -2 & 1 & 1 \\ 1 & 1 & -2 & 1 \end{array} \right)$$

$$\begin{array}{l} \downarrow R_2 \rightarrow R_2 - R_1 \\ \downarrow R_3 \rightarrow R_3 - R_1 \end{array}$$

$$\left( \begin{array}{ccc|c} 1 & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & -\frac{3}{2} & \frac{3}{2} & \frac{3}{2} \\ 0 & \frac{3}{2} & -\frac{3}{2} & \frac{3}{2} \end{array} \right)$$

$$\downarrow R_2 \rightarrow R_2 / \frac{3}{2}$$

$$\left( \begin{array}{ccc|c} 1 & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 1 & -1 & -1 \\ 0 & \frac{3}{2} & -\frac{3}{2} & \frac{3}{2} \end{array} \right)$$

$$\begin{array}{l} \downarrow R_1 \rightarrow R_1 + \frac{1}{2} R_2 \\ R_3 \rightarrow R_3 - \frac{3}{2} R_2 \end{array}$$

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 3 \end{array} \right)$$

$\therefore$  the last row yields the following

Contradiction:

$$0 = 3$$

$$\text{but } 0 \neq 3$$

$\therefore$  the system of linear equations is inconsistent or has no solutions when

$$k = -2$$